

CONSOLIDATED REPORT

AVIATION 1988

709
66.1
1988

17751

JICA LIBRARY



1066202113

REPUBLIC OF PERU
REPORT ON MINERAL EXPLORATION
IN
COTAHUASI AREA
CONSOLIDATED REPORT

JANUARY 1988

JAPAN INTERNATIONAL COOPERATION AGENCY
METAL MINING AGENCY OF JAPAN

PREFACE

At the request of the Government of the Republic of Peru, the Japanese Government planned and carried out a geological survey concerning mineral exploration to examine the possibility of the existence of mineral resources in the Cotahuasi Area located in the southern part of Peru. The Japan International Cooperation Agency was entrusted with the execution of the general plan. The Japan International Cooperation Agency in turn entrusted the execution of this survey to the Metal Mining Agency of Japan since this survey was essentially a professional survey of geology and mineral resources.

The survey was conducted for three years from fiscal 1985 to 1987 and accomplished as schedule with the cooperation of the Peru Government, and particularly the Instituto Geologico Minero y Metalurgico (INGEMMET).

This report is the compilation of the results of the whole survey during these three years.

Lastly, we would like to express our heartfelt gratitude to the members concerned of the Government of the Republic of Peru, the Ministry of Foreign Affairs of Japan, and the Embassy of Japan in Peru, and to all those who extended their kind cooperation to us in executing above-mentioned survey.

January, 1988



Kensuke Yanagiya
President
Japan International Cooperation Agency



Junichiro Sato
President
Metal Mining Agency of Japan




Fig. I - I Location Map of the Project Area

ABSTRACT

The Government of the Republic of Peru had requested Japanese Government for an assistance and co-operation in the mineral exploration of the Cotahuasi Area located in the southern Peru with an intention to activate exploitation of non-ferrous metal industries.

In response to the request, Japanese Government conducted a mineral exploration of the proposed area for the periods of 3 years from 1985 to 1987, by having entrusted the exploration work to the Japan International Cooperation Agency (JICA) and the Metal Mining Agency of Japan (MMAJ).

This report summarizes all the work carried out during the periods and the results of the work.

The 1st year's work comprises a geological interpretation of landsat images for an area of 30,000 km² including the proposed area, and a regional geological and geochemical (stream-sediment sampling) survey for an area of 6,000 km² followed by a follow-up geological prospecting for the selected 3 areas of 800 km² in total.

The above work resulted in having uncovered the regional geology in relation with mineral occurrences in the Cotahuasi area.

The area is distributed by the Jurassic and the Cretaceous systems mainly consisting of sedimentary rocks over the basement of Precambian granite and diorite.

The Tertiary and Quarternary systems mainly consisting of volcanic rocks unconformably overlie these rocks.

Batholiths or stocks of diorite and dikes of andesite intrude the rocks of Miocene age or older.

Mineralization and associated alteration are hosted by the rocks of Miocene age or older, and consist mainly of quartz veins, quartz vein networks and silicified or oxidized zone along fractures or fractures zones, associated with Au and Ag minerals in places.

The 1st year work outlined 4 promising target areas for mineralization, namely (1) Mina Pararapa, (2) Tanisca alteration zone, (3) Mina Luicho and (4) Pirca alteration zone. However, the 3 areas except the Pirca alteration zone were covered by private mining concessions and discarded for further exploration work.

By having re-examined the results of the 1st year's work, the Marcabamba (80 km²) and the Pirca (90 km²) areas were selected for the 2nd year targets. The Pirca area was divided into two sub areas, the eastern and the western Pirca.

Geological mapping and geochemical prospecting (soil sampling) were carried out in both the Marcabamba and the Pirca areas, and 10 holes of diamond core drilling with a total length of 1,000 m were put in the eastern Pirca sub-area.

In the Marcabamba area, a number of mineralization and alteration zones were found either in the Tertiary Tacaza Formation (Tc) or in the Cretaceous Hualhuani (Yu) and Murco (Mu) Formations. Of the zones, two zones, the Colpar and the Marcamalata, were of particular interest because of their association with geochemical anomalies of notable sizes and values.

In the Pirca area, geochemical anomalies were small in sizes and low in values of elements of interest, in spite of extensive alteration zones.

The drill holes in the eastern Pirca sub area failed to intersect sections of mineralization of any values though alteration was prominent.

Based on the results above described, the Colpar (5 km²) and Marcamalata (2 km²) of the Marcabamba area were selected for the 3rd year targets.

The 3rd year's work comprises detailed prospecting with an aid of trending and drilling of 5 holes, 3 250 m-hole for the Colpar and 2 200 m-holes for the Marcamalata.

In the course of the detailed prospecting a number of abandoned old workings and mineralized outcrops were located in the Colpar alteration zone and the geochemical anomaly adjacent to the south which had been

outlined in the previous year. These old workings, without any record remained, had been unknown for years even by local residents.

The 3rd year work located two mineralized zones in the Colpar, the northern mineralized zone including 4 mineralized veins and the southern mineralized zone including 9 mineralized veins, both hosted by the Tacaza Formation (Tc).

These veins are mostly developed along NE-SW trending fractures or fractured zones, and consist of quartz veins, quartz vein networks and silicified zones. Major ore minerals are electrum, argentite, polybasite, pearceite, galena, sphalerite, chalcopyrite and pyrite. An identified alteration mineral assemblage is quartz-potash feldspar (aduralio)-sericite with or without chlorite.

A surface sample of the N3 vein in the northern mineralized zone indicated assay results of 5.70 g/t Au and 640 g/t Ag for a width of 1 m.

The holes MJP-11 and -12 intersected a number of quartz veins, quartz vein networks and silicified zones including the N1 and N2 veins with appreciable values of Au and Ag but were too short to reach the N3 and N4 veins.

The northern mineralized zone may continue to the northern silicified zone to the northeast and form a sizable unit of the mineralization-alteration zone with a length of about 1.5 km and a width of 0.3 km.

Surface samples of the S3 and the S7 veins in the southern mineralized zone indicated assay results of 21.50 g/t Au and 410 g/t Ag for a width of 0.15 m of the S3 vein and 20.10 g/t Au and 1200 g/t Ag for a width of 0.3 m of the S7 vein.

The hole MJP-13 intersected S3, S4 and S5 veins, a part of which indicated appreciable values in Au and Ag with subordinate Cu, Pb and Zn values.

The southern mineralized zone is possible to continue southwestwards under the scree cover to the Auebrada Querumahuaco alteration zone and may form a sizable unit of a mineralization-alteration zone with a length of about 0.9 km and a width of 0.2 km.

The mineralization in the Marcamalata is hosted by sedimentary rocks of the Hualhuani and Murco Formations of Cretaceous age and yields some Ag values with minor Au and other elements. The mineralized veins are discontinuous and limited in extensions in general.

The holes MJP-14 and -15 intersected a number of quartz veins and quartz vein networks but without any Au-Ag values of commercial interest.

The work failed to find any evidences suggesting potential for commercial mineralization.

In conclusion, the 3rd year work resulted in outlining the two mineralization-alteration zones in the Colpar.

However, no promising evidences were formed for the Marcamalata.

CONTENTS

PREFACE
LOCATION MAP
ABSTRACT

PART I SUMMARY REPORT

	Page
CHAPTER 1 OUTLINE OF THE 3 YEAR PROJECT	1
1-1 Location	1
1-2 Purposes	1
1-3 Methods and Techniques	1
1-4 The Project Areas, the Quantities of the Work and Project Periods	1
1-5 Organization of the Survey Team	1
CHAPTER 2 PREVIOUS WORK FOR THE GENERAL AREA	9
2-1 Report of Previous Work	9
2-2 Regional Geology and Mineralization of the Ayacucho-Arequipa Region	9
CHAPTER 3 GENERAL GEOLOGY OF THE COTAHUASI AREA	15
3-1 Geology	15
3-2 Alteration and Mineralization	16
CHAPTER 4 GENERAL REMARKS OF THE COTAHUASI AREA	19
4-1 Introductory Notes to the Republic of Peru	19
4-2 Access to the Project Area	20
4-3 Geography	22
CHAPTER 5 CONCLUSION AND RECOMMENDATION	25
5-1 Conclusion	25
5-2 Recommendation	28

PART II DETAILED REPORT

CHAPTER 1 INTERPRETATION OF LANDSAT IMAGERIES	31
1-1 The Area for the Landsat Imagery Analysis	31
1-2 Result of the Analysis	31
CHAPTER 2 REGIONAL GEOLOGICAL INVESTIGATION OF THE COTAHUAI AREA	35
2-1 Regional Geology and Geological Structure	35
2-2 Alteration-Mineralization	44
2-3 Geochemical Exploration	48
2-4 Conclusion and Recommendation of the 1st Year's Work	52
CHAPTER 3 PIRCA AREA	57
3-1 Geology and Geological Structures	57
3-2 Alteration and Mineralization	58
3-3 Geochemical Exploration	70
3-4 Drilling Operation and Results	84
3-5 Conclusion and Recommendation for the Pirca Area	90
CHAPTER 4 MARCABAMBA AREA	93
4-1 Geology and Geological Structures	93
4-2 Alteration and Mineralization	94
4-3 Geochemical Exploration	102
4-4 Conclusion and Recommendation for the Marcabamba Area	111
CHAPTER 5 COLPAR AREA	115
5-1 Geology and Geological Structures	115
5-2 Alteration and Mineralization	120
5-3 Drilling Operation and Results	131
5-4 Conclusion and Recommendation for the Colpar	149

CHAPTER 6	MARCAMALATA AREA	151
6-1	Geology and Geological Structures	151
6-2	Alteration and Mineralization	153
6-3	Drilling Operation and Results	161
6-4	Conclusion and Recommendation for the Marcamalata	173

PART III CONCLUSION AND RECOMMENDATION

CHAPTER 1	CONCLUSION	175
-----------	------------------	-----

CHAPTER 2	RECOMMENDATION	179
-----------	----------------------	-----

References

LIST OF TABLES

Table I-1	Project Summary
Table I-2	Organization of Negotiation Team and Field Survey Team in Phase I-III.
Table I-3	Principal Mineral Deposits of Ayacucho-Arequipa District Southeastern Peru
Table II-1	List of Alteration and Mineralization Zones (Main)
Table II-2	List of Geochemical Anomaly Zones
Table II-3	Principal Mineralization Zones (Cotahuasi Area)
Table II-4	Principal Geochemical Anomaly Zones (Cotahuasi Area)
Table II-5	Recommendation for Follow-up Surveys (Cotahuasi Area)
Table II-6	List of Alteration and Mineralization Zones of the Pirca Area
Table II-7	Chemical Analyses of Altered Rocks and Ore Samples of the Pirca Area
Table II-8	List of Geochemical Anomaly Zones in the Pirca Area
Table II-9	Comparison of Geochemical Anomaly Zones with Mineralization Zones in the Pirca Area
Table II-10	Principal Alteration and Mineralization Zones of the Pirca Area
Table II-11	Relation with Geochemical Anomaly Zones and Alteration Zones in the Pirca Area
Table II-12	Chemical Analyses of Altered Rocks and Quartz Vein of Drilling Cores
Table II-13	Drilling Results
Table II-14	List of Alteration and Mineralization of Drilling Cores in Pirca Eastern Area
Table II-15	List of Alteration and Mineralization Zones of the Marcabamba Area
Table II-16	Chemical Analyses of Altered Rocks and Ore Samples of the Marcabamba Area

- Table II-17 List of Geochemical Anomaly Zones in the Marcabamba Area
- Table II-18 Comparison of Geochemical Anomaly Zones with Mineralization Zones in the Marcabamba Area
- Table II-19 Comparison of the Pirca Area with the Marcabamba Area on Geochemical Statistic Parameters
- Table II-20 Principal Alteration and Mineralization Zones of the Marcabamba Area
- Table II-21 Relation with Geochemical Anomaly Zones and Alteration Zones in the Marcabamba Area
- Table II-22 Important Mineralized Vein in the Colpar Area
- Table II-23 List of Mineralization Zones of the Colpar Area
- Table II-24 Holes of the Colpar Area
- Table II-25 Mineralization Zones of Drilling Holes in the Colpar Area
- Table II-26 Alteration Zones of Drilling Holes in the Marcamalata Area

LIST OF ILLUSTRATIONS

- Fig. I-1 Location Map of the Project Area
- Fig. I-2 Flowchart of Survey in Cotahuasi Area
- Fig. I-3 Flowchart of Selection of Promising or Deposit Areas
- Fig. I-4 Metalogenic Map of Ayacucho-Arequipa Region Southeastern Peru
- Fig. I-5 Accessibility Map to the Servey Area
- Fig. II-1 Interpretation Map of Landsat False Color Images
- Fig. II-2 Geological Map of the Cotahuasi Area
- Fig. II-3 Generalized Stratigraphic Column of the Cotahuasi Area
- Fig. II-4 $\text{SiO}_2 - (\text{Na}_2\text{O} + \text{K}_2\text{O})$ Variation Diagram of Valcanic Rocks
- Fig. II-5 $\text{K}_2\text{O} - \text{MgO}$ Variation diagram of Valcanic Rocks
- Fig. II-6 Normative Quartz-Orthoclase - (Albite+Anorthite) Triangular Diagram of Some Igneous Rocks
- Fig. II-7 Location Map of Alteration and Mineralized Zones
- Fig. II-8 Geochemical Interpretation Map (Composite Data)
- Fig. II-9 Geological Map of the Pirca Eastern Area
- Fig. II-10 Geological Map of the Pirca Western Area
- Fig. II-11 Location Map of Alteration and Mineralization Zone of the Pirca Eastern Area
- Fig. II-12 Location Map of Alteration and Mineralization Zone of the Pirca Western Area
- Fig. II-13 Geochemical Interpretation Map of the Pirca Eastern Area (Composite Data)
- Fig. II-14 Geochemical Interpretation Map of the Pirca Western Area (Composite Data)
- Fig. II-15 Interpretation Map of the Pirca Eastern Area
- Fig. II-16 Interpretation Map of the Pirca Western Area
- Fig. II-17 Location Map of the Drilling Sites
- Fig. II-18 Drilling Progress of the Pirca Eastern Area (MJP-1 - 10)

- Fig. II-19 Geological Map of the Marcabamba Area
- Fig. II-20 Location Map of Alteration and Mineralization Zone of the Marcabamba Area
- Fig. II-21 Geochemical Interpretation Map of the Marcabamba Area (Composite Data)
- Fig. II-22 Interpretation Map of the Marcabamba Area
- Fig. II-23 Geological Map and Section of the Colpar Area
- Fig. II-24 Stratigraphic Column of the Colpar Area
- Fig. II-25 Location Map of Alteration and Mineralization Zone of the Colpar Area
- Fig. II-26 Location Map of Mineralization Zones of the Northern Part of the Colpar Area
- Fig. II-27 Location Map of Mineralization Zones of the Southern Part of the Colpar Area
- Fig. II-28 Location Map of the Drilling Sites
- Fig. II-29 Location Map of the Drilling Sites in the Colpar Area
- Fig. II-30 Drilling Progress of the Colpar Area (MJP-11 - 13)
- Fig. II-31 Geological Section of the Drilling Holes (MJP-11, 12, 13) in the Colpar Area (Scale 1:5,000)
- Fig. II-32 Geological Section of Drilling Hole MJP-11 in the Colpar Area
- Fig. II-33 Geological Section of Drilling Hole MJP-12 in the Colpar Area
- Fig. II-34 Geological Section of Drilling Hole MJP-13 in the Colpar Area
- Fig. II-35 Geological Map and Section of the Marcamalata Area
- Fig. II-36 Stratigraphic Column of the Marcamalata Area
- Fig. II-37 Location Map of Alteration and Mineralization Zone of the Marcamalata Area
- Fig. II-38 Location Map of Mineralization Zones of the Marcamalata Area

- Fig. II-39 Location Map of the Drilling Sites
- Fig. II-40 Location Map of the Drilling Sites in the Marcamalata Area
- Fig. II-41 Drilling Progress of the Marcamalata Area (MJP-14 - 15)
- Fig. II-42 Geological Section of the Drilling Holes (MJP-14, 15) in the Marcamalata Area (Scale 1:5,000)
- Fig. II-43 Geological Section of the Drilling Hole MJP-14 in the Marcamalata Area
- Fig. II-44 Geological Section of Drilling Hole MJP-15 in the Marcamalata Area
- Fig. III-1 Interpretation Map of the Colpar Area

PART I SUMMARY REPORT

PART 1 SUMMARY REPORT

CHAPTER 1 OUTLINE OF THE 3 YEAR PROJECT

1-1 Location

The project area is situated in the southern part of the Republic of Peru. The center of the area is about 200 km northwest of Arequipa, the second largest city in the country, at the latitude of about 15°15'S and the longitude of about 73°00'W (Fig. I-1).

1-2 Purposes

The project was intended to evaluate the potential of the area for possible commercial mineral occurrences by comprehending the geology in relation with mineralization and associated alteration.

1-3 Methods and Techniques

The applied methods and techniques are summarized in Table 1-1 and Fig. 1-2 in accordance with the progress of the 1st year's through the 3rd year's project. Fig. 1-3 provides the flows of the decision-makes and the actions taken based on the results of each year's work.

1-4 The Project Areas, the Quantities of the Work and Project Periods

The areas investigated and the quantities of the work are summarized in Table 1-1 and Fig. 1-1 for each year.

The time required for each year's programme is also summarized in Table 1-1.

1-5 Organization of the Survey Team

Organization of negotiating team and field survey team in phase I to III are summarized in Table I-2.

Table I-1 Project Summary

Year	Method	Area	Quantity	Field Work	Laboratory Work	Period
1st year	LANDSAT data analysis	Area covered by LANDSAT data analysis (Lat. 14°30'S - 16°00'S and Long. 72°00'W - 74°00'W)	30,000 km ²	Interpretation of geological structure		From 25th Sept. '85 to 28th Feb. '86
	Geological survey and geochemical exploration	Regional survey area (Lat. 15°00'S - 15°30'S and Long. 72°30'W - 73°30'W) Detailed survey areas: Areas A, B and C	Area: 5,200 km ² No. of geochemical samples: 1,309 Total area of 3 areas (A, B and C): 800 km ²	Exploration density: 0.29 km/km ² (1,439 km/5,200 km ²) Sampling density: 0.25/km ² (1,309/5,200 km ²) Geochemical samples: Riverbed deposits Indicator element: Au, Ag, Cu, Pb, Zn and As Exploration density: 0.80 km/km ² (500 km/800 km ²) Sampling density: 1.02/km ² (815/800 km ²) Geochemical samples: Riverbed deposits Indicator element: Au, Ag, Cu, Pb, Zn and As K - Ar method	Microscopic observation of rocks and ores X-ray diffraction (powder method) analysis Chemical analysis of rocks (13 components) Chemical analysis of ores (Au, Ag, Cu, Pb and Zn)	
		Orcopampa Mine and entire surrounding area	No. of geochemical samples: 50 No. of samples for absolute age determination: 5			
2nd year	Geological survey and geochemical exploration	Marcabamba Area	Area: 80 km ²	Exploration density: 2.1 km/km ² (164 km/80 km ²) Sampling density: 7.8/km ² (620/80 km ²) Geochemical samples: Soil samples (620/80 km ²) Indicator element: Au, Ag, Cu, Pb, Zn and As	Microscopic observation of rocks and ores X-ray diffraction (powder method analysis)	From 11th Jul. '86 to 10th Feb. '87
		Pirca Area	Area: 90 km ² No. of geochemical samples: 1,408	Exploration density: 2.9 km/km ² (261 km/90 km ²) Sampling density: 15.6/km ² (1,408/90 km ²) Geochemical samples: Soil samples (1,408/90 km ²) Indicator element: Au, Ag, Cu, Pb, Zn and As	Chemical analysis of rocks (13 components) Chemical analysis of ores (Au, Ag, Cu, Pb and Zn)	
3rd year	Drilling operation	Pirca Area	10 holes Total drilling length: 1,000 m	Drilling length 100 m x 10 holes (vertical drilling)		
	Geological survey (Detailed survey including trench survey)	Colpar Area Marcamalata Area	Area: 5.0 km ² Trench survey: Total length 179.5 m Area: 2.0 km ² Trench survey: Total length 153 m	Exploration density: 5.2 km/km ² (26.1 km/5.0 km ²) Exploration density: 8.1 km/km ² (16.2 km/2.0 km ²)	Microscopic observation of rocks and ores X-ray diffraction analysis (powder method) Chemical analysis of ores (Au, Ag, and partly Cu, Pb and Zn)	From 26th Jun. '87 to 30th Jan. '88
	Drilling operation	Colpar Area Marcamalata Area	3 holes Total drilling length: 750 m 2 holes Total drilling length: 400 m	Drilling length 250 m x 3 holes (-45° inclined drilling) Drilling length 200 m x 2 holes (-45° inclined drilling)		

Table I-2 Organization of Negotiating Team and Field Survey Team in Phase I-III

	Negotiating team					Field survey team						
	Japanese side			Peruvian side		Japanese side			Peruvian side			
1985	Makoto Ishida	leader	Metal Mining Agency of Japan	Juan Zegarra W.	leader	Instituto Geologico Minero y Metalurgico	Hiroshi Miyajima	leader	Sumiko Consultants Co., Ltd.	Cesar Vilca Neira	leader	Instituto Geologico Minero y Metalurgico
	Yoshikazu Taketomi		Ministry of International Trade and Industry; Agency of Natural Resources and Energy	Gregorio Flores N.		ditto	Kiyoharu Nakashima	member	ditto	Hector Zarate Olazagal	member	ditto
	Sumihiro Fure		Metal Mining Agency of Japan; Lima Office	Erick Soriano B.		ditto	Masatsugu Sakai	ditto	ditto	Walter Rodriguez Olaete	ditto	ditto
	Tadaaki Ezawa		Metal Mining Agency of Japan				Seiji Tsuchida	ditto	ditto	Carlos Jimenez Velasco	ditto	ditto
	Takashi Kamiki		ditto				Jiro Natori	ditto	ditto	Emilio Rojas Rivera	ditto	ditto
1986	Makoto Ishida	leader	Metal Mining Agency of Japan	Juan Zegarra Wuest	leader	Instituto Geologico Minero y Metalurgico	Kiyoharu Nakashima	leader	Sumiko Consultants Co., Ltd.	Cesar Vilca Neira	leader	Instituto Geologico Minero y Metalurgico
	Sumihiro Fure		ditto (Lima Office)	Gregorio Flores Nãnes		ditto	Masayoshi Matsugi	member	ditto	Hector Zarate Olazagal	member	ditto
	Takashi Kamiki		ditto	Cesar Vilca Neira		ditto	Jiro Natori	ditto	ditto	Emilio Rojas Rivera	ditto	ditto
				Walter Rodriguez Olarte		ditto	Akihiko Murase	ditto	ditto	Daniel Carpio T. and others	contractor	Geotec S.A.
				Emilio Rojas Rivera		ditto						
1987	Kazunori Kano	leader	Metal Mining Agency of Japan	Juan Zegarra Wuest	leader	Instituto Geologico Minero y Metalurgico	Kiyoharu Nakashima	leader	Sumiko Consultants Co., Ltd.	Cesar Vilca Neira	leader	Instituto Geologico Minero y Metalurgico
	Kenzo Hagio		ditto (Lima Office)	Gregorio Flores Nãnes		ditto	Atsunu Nonami	member	ditto	Hector Zarate Olazagal	member	ditto
	Yozo BABA		ditto	Cesar Vilca Neira		ditto	Akihiko Murase	ditto	ditto	Walter Rodriguez Olarte	ditto	ditto
	Naotaka Adachi		ditto	Hector Zarate Olazagal		ditto				Pedro Samame I. and others	contractor	Geotec S.A.

Fig. I-2 Flow Chart of Survey in Cotahuasi Area






Fig. I-3 Flow Chart of Selection of Promising Ore Deposit Areas





CHAPTER 2 PREVIOUS WORK

2-1 Report of Previous Work

The reports of the previous work are listed in the appendices of this report.

Fornari, M and Vilca, C.N. (1977) describe the general geology of the Ayacucho-Arequipa region in their report on argentiferous mineral occurrences in the southern Peru.

Metallogenic maps covering the whole area of Peru have been prepared by Bellido, D. et al (1972) and published by the Minister of Energy and Mines of the Republic of Peru.

Local geology is described by Victor, P.G. for Pausa and Caraveli (1983) and by Carlos G.R. and David, D.M. for Coracora Pacapausa (1983).

Results of geochemical investigation of the Andes zone, southern Peru, are available in papers collected by Onuma, N. (1985).

The following explanation of the regional geology and the mineralization are summarized from the reports of the previous work above mentioned.

2-2 Regional Geology and Mineralization of the Ayacucho-Arequipa Region

(1) Geology

The region occupies a large of the southern Peru include the Cotahuasi area, the area for the 3 year programme, outlined by the latitudes of 14°00' and 16°30'S and the longitudes of 71°30' and 74°30'S (Fig. I-4).

The west Andes Range in this region forms a great plateau about 150 km wide with an average elevation of about 4,000 m. The plateau is mildly relieved with occasional high peaks, the highest of which reaches 6,426 m above sea level.

The Andes Range, since the cessation of uplifting, have been subject to retrogressive erosion having initiated from the west coast facing the Pacific Ocean. However, most of the area have not been deeply dissected as yet except the Cotahuasi area where many deep valleys with about 2,000 m elevation differences are well developed.

Signs of glaciation are observed at a number places on the pleau and thick morrains or glacial soils and cirques are well developed. The rocks distributed in this area comprise metamorphics of Precambrian era and sedimentaries and volcanics of Palaeozoic, Cesozoic and Cenozoic eras.

The Precambrian metamorphics are distributed along the Oceanic coasts or along a few rivers at their bottoms, and consist mainly of gneisses or gneissic granites.

The Palaeozoic sedimentary rocks unconformably overly the Precambrian basement. This area emerged above sea level towards the end of the Carboniferous and, supplied clastics to surrounding sedimentary basins in the Permian.

These Permian sedimentary basins become sites for volcanisms and were deposited thick volcanic piles interbedded with marine sediments in Triassic age.

The area started emerging again in the Cretaceous. The tectonic of Cretaceous is characterized by an extensive Orogensis which formed the west Andes Range of the NW-SE trending folding mountains at the end of the Cretaceous. Granodioritic batholiths along the Oceanic coast intruded mostly during Cretaceous age.

Red beds, terrestrial clastic sediments, deposited in the late Cretaceous and the early Tertiary.

Oligocene to miocene volcanics and pyroclastics unconformably overly older rocks. At the base of the volcanics and pyroclastics, there are developed volcanic breccias containing agglomerates and tuffs.

The upper section consists mainly of andesitic volcanics with subordinate rhyolites and is co-related to the Tacaza formation.

The Tacaza formation changes its lithology laterally in the area of argentiferous mineral deposits. The western part, the Pugio zone, is dominated by pyroclastics and sediments interbedded with layers or lenses of shales, gypsum and carbonates. These lithological features indicate lagoonal natures of the sedimentary basins. These characteristic sediments is absent in the eastern part or developed only to a limited extent.

The Oligocene-Miocene pyroclastic are gently folded forming anticlines and synclines with bedding planes dipping upto 60 degrees.

Though individual tectonic elements are variable in their direction, the major folding belts trend in the NW-SE direction parallel to the Andean mountains. The trend may be a remnant of the major tectonic element of the late Cretaceous.

Ignimbrites of the Pliocene to Quarternary is widely distributed covering eroded surfaces.

The tuffs are not influenced by the folding structures observed in the Tacaza formation.

In addition, there are andesitic lavas, the activity for which initiated in early pliocene and have continued to the present time.

A number of strato volcanics in the southern Peru formed mostly in the Pleistocene. Andesitic lavas extruded from these volcan are extensively distributed to form lava plateaus or fill valley bottoms.

(2) Mineralization

Known mineral occurrences in the Ayacucho-Arequipa region are shown in Fig. 1-4. Table I-3 presents particulars of major occurrences of these.

The metallogenic map of Peru divides the region into 4 metallogenic sub provinces, namely the highland polymetallic, the Oceanic Copper, the Oceanic Iron, and the Precambrian Metamorphic. The highland polymetallic sub province comprises the Andahuaylos-Yauri and the Puquio-Caylloma blocks. The northwestern part of the Oceanic Copper sub-province is named of the Nazca-Ocona block.

The Andahuaylos-Yauri subprovince, being located in the northeast of the region, is mainly distributed by Iron-Copper mineral occurrences of a pyrometasomatic type. The Puquio-Caylloma subprovince occupies the central part of the region including the Cotahuasi project area and is distributed by hydrothermal or pneumatritic Au and/or Ag mineral occurrences. Hydrothermal or pneumatitic Cu and Au mineral occurrences are known in the Nazca-Oconia block occupying the southwestern part.

All of the mineralization is believed to have been formed in the periods of late Cretaceous or Tertiary age.

The Cotahuasi project area, being located in the central part of the region, occupies parts of the Puquio-Caylloma subprovince in the northeast and of the Nazca-Ocona block in the southwest.

The Colpar which have been selected for a potential area for Au-Ag mineralization, belongs to the Nazca-Ocona Cu or Ag metallogenic area in the Oceanic Copper subprovince.




Table I - 3 Principal Mineral Deposits of Ayacucho - Arequipa District Southeastern Peru

	PRINCIPAL MINERAL DEPOSITS	MINERAL ASSOCIATIONS	METALLOGENIC PROVINCES	METALLOGENIC EPOCH	TYPE OF MINERAL DEPOSITS	DIMENSIONES
95	Chalcobamba	Cu	HIGHLAND POLYMETALIC SUB-PROVINCE Andahuallas - Yauri Area	Upper Cretaceous - Tertiary	Contact metasomatic, Irregular	Medium
96	Yuringa	Cu	ditto	ditto	Hydrothermal or Pneumatolytic, Vein or Seam	Small
97	Charcas	Cu	ditto	ditto	Contact metasomatic, Irregular	Small
98	Sulfobamba	Cu	ditto	ditto	ditto	Small
99	Ferrobamba	Cu	ditto	ditto	ditto	Medium
103	Livitaca	Fe	ditto	ditto	ditto	
104	Colquamarca	Au	ditto	ditto	Hydrothermal or Pneumatolytic, Mantle or Stratiform	Occurrences or Prospects
106	Katange	Cu	ditto	ditto	Hydrothermal or Pneumatolytic, Irregular	Medium
108	San Juan de Lucanas	Au, Ag, Pb, Zn	HIGHLAND POLYMETALIC SUB-PROVINCE Pucalloma - Caylloma Area	ditto	Hydrothermal or Pneumatolytic, Vein or Seam	Medium
111	Arcata	Au, Ag	ditto	ditto	ditto	Small
114	Suykutambo	Ag, Pb, Zn	ditto	ditto	ditto	Medium
116	Orcopampa	Au, Ag, Cu, Pb	ditto	ditto	ditto	Small
117	Caylloma	Au	ditto	ditto	ditto	Medium
123	San Luis	Au	PACIFIC CUPRIFEROUS SUB-PROVINCE Nazca - Ocona Area	ditto	ditto	Small
124	La Capitana	Cu	ditto	ditto	ditto	Small
128	Madrigal	Au	HIGHLAND POLYMETALIC SUB-PROVINCE	ditto	ditto	Big
129	Calpa	Au	PACIFIC CUPRIFEROUS SUB-PROVINCE Nazca - Ocona Area	ditto	ditto	Small
130	Alpacay		ditto	ditto	ditto	Small
132	Andaray	Cu	ditto	ditto	Contact metasomatic, Irregular	Small
133	Pasco	Au, Ag, Cu	ditto	ditto	Hydrothermal or Pneumatolytic, Vein or Seam	Small

CHAPTER 3 GENERAL GEOLOGY OF THE COTAHUASI AREA

3-1 Geology

The Precambrian basement complex comprises mainly gneiss and diorite and is located in the southern part of the area for the field work of the project. (The name Cotahuasi specifies the area for the field work)

The Jurassic system, unconformably overlying the Precambrian basement, consists of green or brown andesitic volcanics in the lower part with a development of foliated structures at its bottom and of grey - black limestone, dark - greenish grey shales and thick sandstones in the middle and the upper parts.

The Cretaceous system is composed of light grey - grey thick silicious sandstone in the lower part and of shales with subordinate sandstones and grey - dark grey limestones containing fossils.

Intensive faulting and folding structures are extensively developed in the Jurassic and the Cretaceous systems which are unconformably overlain by the tertiary system.

The lower part of the Tertiary system consists of sedimentary rocks mainly of sandstones and conglomerates and is overlain by andesitic or dacitic volcanics with highly variable facies, which are in turn overlain by rhyolitic pyroclastics.

Folding structures are locally developed in the sedimentary and andesitic volcanic rocks of the lower part.

The Quarternary system includes volcanoes such as Nevado Solimana, which exceed 5,000 m in elevations, and is composed of andesitic volcanics of the lower most formation widely distributing over the Archiplano plateau, and of overlying glacial deposits with local development of andesites and basalts. Alluvials are developed along rivers, on gentle slopes or in basins such as the Archiplano basin, and consists generally of taffaceous materials.

Major intrusives are Cretaceous granitic batholiths or stocks belonging to the Coast batholith, and dioritic and andesitic stocks or dikes of Tertiary age.

The arrangement of the Cretaceous intrusive bodies generally trends WNW-ESE, parallel to the trend of the Coast batholith.

The arrangement of the Tertiary intrusive bodies are not well defined due to the Quarternary covers but there appear to be two directions NE-SW and NW-SE.

The rocks of the early Miocene or earlier (the Tacaza formation or lower) have been subject to folding and faulting which have formed a number of geological blocks.

No major folding or faulting is developed in the rocks of the late Miocene or later (the Alfabamba formation or upper)

Axes of anticlines and synclines run generally in the direction of NW-SE or WNW-ESE.

NW-SE trending faults are predominated in this area with subordinate occurrences of E-W, NE-SW or N-S trending faults.

3-2 Alteration and Mineralization

Alteration and mineralization are not well developed in the area distributed mainly by sedimentary rocks of Cretaceous age due to development of limestones and siliceous sandstones.

Alteration zones of sizable scales associated with reddish brown stains are often observed in the area distributed by volcanic rocks of Miocene age or earlier.

Alteration zones are located at Oyolo and Mina Lucho of the Area A, at Tanisca and Mina Picha of the Area B, and at Mina Pararapa of the Area C in

the area for the detailed work of Phase I, as well as at Sara Sara, Pirca and Maran in the area for the regional work.

Of these alteration zone, the Tanisca zone is the most extensive continuing in the direction of N-S for approximately 10 km with widths ranging from 1 to 3 km. The alteration is argillization in the andesitic volcanics of the Tacaza formation, associated with weak silicification in part.

Oxide clay veins and quartz veins occur in the underlying silicious sandstones of the Yura formation and have been once meagrely prospected.

The Mina Pararapa alteration zone is formed along a NNW-SSE trending porphyritic andesite dike and in the surrounding andesitic volcanics of the Tacaza formation and shows brown colour and outcrops due to iron oxide stains.

A quartz vein with some Au-Ag values is located in association with alteration zone, and strikes N20°E dipping 80°NE.

The vein had been explored in underground until a recent year. Au values, variable from place to place, are averaged at 4.61 g/t for approximately 200 m along the strike, though occasionally as high as 40 g/t according to data at the Mine (Mina Pararapa).

The vein can be traced for approximately 1.3 km with variable widths ranging from 0.5 to 1.5 m. There are good potential for parallel veins nearby.

The alteration zone is approximately 1 km wide and 2.5 km long.

The Mina Luicho and the Maran alteration zones are also associated with Au-Ag mineralization but smaller in dimensions than those above described.

The Mina Picha alteration zone is located at the contact between limestones of the Cretaceous Arcurquia formation and a diorite stock, and contains skarn minerals in association massive Cu·Pb·Zn ores.

The mineralization is apparently of pyrometasomatic origin.

Other alteration zones than above described, showing brown colour due to oxide iron stains, are associated with only poor mineralization.

Major alteration zones identified by the landsat imagery analysis are located to the southeast of Nevado Gororupa, nearby Nasama and to the Northeast of Nevado Firura and appear to consist of white clay alteration.

CHAPTER 4 GENERAL REMARKS OF THE COTAHUASI AREA

4-1 Introductory Notes to the Republic of Peru.

The Republic of Peru is situated in the Pacific Ocean Side of the South American Continent and occupies an area of 1,285,215 km² extending from the Equator to the latitude 18°20'S.

The total population is estimated at approximately 19.2 millions (1984), of which natives (Indios) account for 47%, mestizos (mixed bloods between natives and white) for 40% and whites for 12%. The Country, together with Bolivia, is highly populated with the natives among the South American countries.

The capital of the Country is Lima.

The official language is Spanish which is widely used all over the country. However, Quechua is also popular among the natives and Aimava is locally used in the part near the boarder to Bolivia.

The Country's gross national product amounted to approximately 18 billion dollars in U.S. (1984) or 789\$ per capita.

The domestic production of primary energy totals 11,080,000 tons in crude oil equivalent, and exceeds the domestic consumption of 7,660,000 tons.

The Country, being abundant in mineral resources was ranked at the 2nd place in the world for the silver production (1,739 tons), at the 4th for the zinc (580,000 tons), at the 5th for the lead (210,000 tons), and at the 8th for the copper, according to the 1983 records. Petroleum and natural gas are also plentiful in the country.

According to the 1983 statistics, the Country's exports and imports amounted to 3,015 million and 2,680 million dollars in U.S. respectively, and the trade ballance was surplus by 327 million dollars in U.S.

The Nation's official external debt amounted to 11,484 million dollars in U.S.

The country exported zinc, silver and copper ores totalling US\$330 millions to Japan, and in turn imported automobiles, steels, machinery and other processed materials totalling US\$140 millions in 1985. This resulted in the Pere's trade surplus by US\$190 millions.

Galloping inflation brought about a serious economic dislocation in Peru before 1984. After coming to power, the present Peruvian regime has implemented a series of measures aimed at the stabilization of the national economy, such as the fixing of the exchange rate of Soles against the US dollar and price control. The present bank exchange rate of Soles against the US dollar stands at 17,300 Soles to US\$1.00 and the official rate at 14,000 Soles to US\$1.00. The Peruvian Government has announced its intention to freeze the exchange rates and oil price until the end of the first six-month period of 1986.

4-2 Access to the Project Area

The survey areas are located in the southern part of the Republic of Peru.

The principal communities in the survey area which are served by a road passable in vehicles are Cotahuasi and Pausa, both located almost in the center of the area. The access road to the two communities starts in Arequipa, Peru's second largest city, but there is no roadway connecting the communities directly (Fig. I-5).

Cotahuasi can be reached by following the National Highway 1 from Arequipa to Tambillo and then taking Departmental Road 103 running northwestward and passing Pta Colorado, Aplao and Chuquibamba. The distance between Arequipa and Cotahuasi is about 410 km and can be covered in eight to ten hours by car. That section of the access road running from Arequipa to Pta Colorado (nearly 180 km) is paved, but the remaining section (about 230 km) leading to Cotahuasi is a gravel road.




Fig. I-5 Accessibility Map to the Project Area

Pausa can be reached by following the national Highway 1 along the coastline from Arequipa to Pta Tinaja, then proceeding northward to Craveli and from there taking the Departmental Road 100. The distance between Arequipa and Pausa is about 490 km and can be covered in 10 to 12 hours by car. The Arequipa-Pta Tinaja section of about 280 km in length is paved and the remaining section of about 210 km leading to Pausa is a gravel road. The roadway condition between Arequipa and Pausa is worse than those of the Arequipa-Cotahuasi route.

Pausa can also be reached by following the National Highway 100 from Atica and then taking the Departmental Road 115 from Chala, but this route takes much longer time.

The roads passable by vehicle in the survey area, with the exception of the above-mentioned access roads from Arequipa, include a gravel road of about 20 km connecting Alca and Cotahuasi, a second gravel road of about 15 km running from Pausa to Lampa along a river and another gravel road running about 8 km to the south. There are no other roadways and horses provide the only means of transportation of men and goods.

4-3 Geography

(1) Topography

The survey area is located in the southwest of the Altiplano, a plateau 4,000 to 5,000 m above sea level, in the Andes. The Altiplano is deeply incised by two Grand Canyon-shaped valleys of relative height ranging from 1,000 to 2,500 m which are believed to have been formed by dissection after the middle stage of the Tertiary. These are the rivers of Rio Maran and Rio Cotahuasi which meet together into the Rio Ocona which pours in the south into the Pacific.

Rising above the Altiplano are Quaternary volcanos 5,500 to 6,300 m above sea level which range from WNW to ESE. These include Nevado Sara Sara (5,505 m), Nevado Solimana (6,093 m), Nevado Firura (5,498 m) and Nevado Coropuna (6,377 m). In the ESE direction other Quaternary volcanos of comparable height including Nevado Anpato (6,288 m), stretch into outside the survey area.

The Altiplano generally consists of gently sloping hills and stretches of flatland among them. Marshlands or extensive grasslands called "pampa" and marshes or lagoons called "laguna" are formed in the basins.

The valleys shaped like the Grand Canyon are mostly sloped at a greater angle than 40° and they often form precipitous (sloped at 60° to 70°) or nearly perpendicular cliffs. There are intermittent stretches of alluvial flatland of limited area along major rivers. The breast of the steep slopes are dotted with gently sloping flatlands of limited area.

(2) Climate and Vegetation

The region extending from the southwestern end of the survey area to the Pacific Coast belongs to the arid zone of the South American Pacific Coasts which is characterized by the desert type climate which permits of almost no vegetation. On the other hand, the Altiplano district including the survey area belongs to the semiarid climate and is generally called a zone of grass-covered cold highlands and basins.

In the survey area, the difference between the lowest and highest elevation is approximately 5,200 m, the lowest being the vicinity (1,150 m) of the southern bed of Rio Cotahuasi and Rio Maran and the highest being Nevado Coropuna (6,377 m). As a result, the climate and vegetation of the survey area vary widely with elevation.

In the areas along the Rio Cotahuasi and Rio Maran which are at a lower elevation than 2,300 m, the temperature is higher and papayas, bananas, grapes, pacays and other fruits are raised and a variety of trees and plants grow.

The areas lying between 2,300 and 3,500 m in height have a mild semiarid climate and maize, wheat, other cereals and broad beans are raised there, but fruits are almost non-existent. In the hill district several species of cactus and shrubs grow rather densely. Most of the communities, including Cotahuasi and Pausa where a base camp was set up, are concentrated in this mild semiarid climate zone.

The climate of the mountain district ranging in elevation between approximately 3,500 m and 5,000 m is a cool to cold semiarid climate. The midslopes of the mountains of up to about 4,000 m in height are dotted with a growth of several species of cactus and shrubs, but in the Altiplano district ranging in elevation from 4,000 to 5,000 m the growth of cactus and shrubs is very limited and a greater part of this district forms a desert except a wide variety of moss growing in the pampas and scatterings of low shrubs.

The mountains of 5,000 m or greater height are crowned with perpetual snow and in this mountain district the weather is extremely cold and no vegetation is found.

In the Altiplano district, rainfall is generally small and the dry season lasts from April to September and the wet season from October to March. In the survey area, the wet season sets in earlier in its eastern part and rainfall starts in October, but in the western part the wet season tends to begin half a month to a month later. During the wet season, snow falls in the mountains of 4,000 m or more in height and rain falls in the lower mountains.

CHAPTER 5. CONCLUSION AND RECOMMENDATION

5-1 Conclusion

The exclusive conclusions of the 3 year-project are summarized as follows.

- (1) The basement of the project area consists of gneissic granites or diorites of Precambrian age.

The Jurassic and Cretaceous systems, mainly consisting of sedimentary rocks, unconformably overlies the Precambrian basement and are unconformably underlain by volcanic rocks of the Tertiary and the Quaternary systems. Intrusions comprise granitic batholiths or stocks of Cretaceous age and andesitic and dioritic stocks or dikes of Tertiary age.

- (2) Folding and NW-SE or NE-SW trending fault structures are well developed in the Jurassic and the Cretaceous systems as a result of the Andean Orogeny.

NW-SE or NE-SW trending faults are also observed in the Miocene volcanics, particularly the Tacaza formation (Tc).

These structural features are well expressed in the Landsat imagery.

- (3) Mineralization and alteration occur mainly in the Tacaza formation of Miocene age or lower formations.

Most of the known deposits in the project area are of Au-Ag vein type and consist of quartz veins, quartz vein networks, or silicified or oxidized fractures.

A pyrometamorphic deposit of a small scale has been located at the contact between a diorite stock and limestones of the Cretaceous Arcurquina formation.

- (4) Alteration (silicification and argillization) zones of sizable scales were outlined in the Pirca area in the course of the 1st year's field work.

However, the 2nd year's geochemical investigation indicated that geochemical anomalies associated with the alteration zones were limited in their extensions and low in values of elements of interest.

The results of the 10 holes of drilling in the eastern Pirca area intersected intensive alteration zones but with only minor mineralization.

- (5) Alteration and mineralization zones were outlined in the Tacaza formation (Tc) of Miocene age and in the Hualhuani (Yu) and the Murco (Mu) formation of the Cretaceous age in the Marcabamba area.

Of a number of the alteration (silicification and argillization) and mineralization zones outlined in the course of the 2nd year's investigation, the Colpar and the Marcamalata alteration zones were associated with promising geochemical anomalies.

The 2nd year's work resulted in selection of two target areas, the Colpar (5 km²) and the Marcamalata (2 km²), including these alteration zones, for the 3rd year's programme.

- (6) A number of abandoned old workings, which had been unknown, were located in the Colpar in the course of the 3rd year's work.

Two mineralized zones, the northern and the southern mineralized zones, were outlined in the Miocene Tacaza formation (Tc) by the detailed prospecting with aids of trending.

These mineralized zones include mineralized veins, 4 major veins in the northern zone and 9 in the southern zone, in association with surrounding silicification and minor parallel veins.

All the vein except for two in the southern zone trend in the general direction of NE-SW, and comprise quartz veins, quartz vein network and silicified fracture zones.

Major ore minerals are electrum, argentite, polybasite, pearceite, galena, sphalerite and pyrite in association with alteration minerals of quartz, potash feldspar and sericite with or without chlorite.

- (7) The N3 vein of the northern vein yielded a surface sample of the best assay results, 5,79 g/t Au and 640 g/t Ag for an 1 m width.

The drill holes, MJP11 and 12, intersected the N1 and N2 veins but were too short to reach the N3 and N4 veins. The mineralized intersections yielded appreciable values in Au and Ag, which would suggest the mineralized zone to be continuous for a substantial distances.

The northern silicified zone, located to the northeast of the northern mineralized zone, may continue to the northern mineralized zone. If the assumption is true, the incorporated alteration mineralization zone would form a sizeable area approximately 0.3 km wide and 1.5 km long.

- (8) In the southern mineralized zone, the S3 and the S7 veins yielded surface samples of high assay values, 21.50 g/t Au and 410 g/t Ag for, a 0.15 m width and 20.10 g/t Au and 1,200 g/t Ag for a 0.3 m width respectively.

The drill hole, MJP-13, intersected the S3, the S4 and S5 veins and the quartz vein network of the S4 vein indicated assay values of 5.04 g/t Au, 45.0 g/t Ag, 0.79 % Cu, 1.37 % Pb and 1.30 % Zn. The mineralization contains appreciable values of Cu, Pb and Zn in addition to Au and Ag, and is considered to be promising.

The southern mineralized zone may continue to the Quebrad Quermahuaico alteration zone. If the assumption is true, the incorporated alteration mineralization zone would form a sizable area approximately 0.2 km wide and 0.9 km long.

- (9) The mineralization-alteration in the Marcamalata occurs in the Cretaceous Hualhuani (Yu) and Murco (Mu) formation and is associated with a minor amount of Ag.

All the observed veins are limited in extentions and discontinuous.

The two holes, having been drilled beneath the surface mineralized veins, intersected a number of quartz veins and quartz vein networks but without mineralization of any values.

- (10) In conclusion, the two mineralization-alteration zones, the northern (0.3 km wide and 1.5 km long) and the Southern (0.2 km wide and 0.9 km long), seem to be promising judging from the exclusive results of the 3 year work, and would be worthwhile for further exploration.

5-2 Recommendation

The fiscal year 1987, closing at the end of March, 1988, is the last year of the 3 year technical co-operation project in the Cotahuasi area by Japanese Government.

However, the 3 year work for the project have outlined two promising targets for potentially commercial mineralization in the Colpar, the northern mineralization-alteration zone (0.3 km wide and 1.5 km long), and the southern mineralization-alteration zones (0.2 km wide and 0.9 km long).

It would be recommended that these two zones in the Colpar be followed up by further detailed exploration including diamond core drilling.

Methods and purposes of the recommended work are summarized in the table below.

Method	Purposes
Detailed Prospecting	To clarify occurrences, extents, and grades of the mineralization on the surface by investigating further in detail mineralized outcrops and abandoned old workings of the mineralization-alteration zones.
Drilling	To define extents and grades of veins in strike sides and dip sides, and also to explore other veins parallel to the known ones.

PART II DETAILED REPORT

PART II DETAILED REPORT

CHAPTER 1 LANDSAT IMAGERY ANALYSIS

1-1 The Area for the Landsat Imagery Analysis

An area of 30,000 km² was selected for the Landsat Imagery Analysis, including the Cotahuasi Area. (Fig. I-1)

1-2 Result of the Analysis

The Landsat Imageries covering the above area were interpreted with an emphasis placed on comprehending regional features of the geological structure.

The results of the analysis are shown in the Interpretation map of the Landsat Imagery. (Fig. II-1)






Fig. II-1 Interp Lands



0 10 20km

LEGEND

- | | | |
|---------------------------------------|-------|--------------------------------------|
| | Or-ol | Sediment |
| | Or-2 | Lava flow |
| Quaternary | Or-1 | Pyroclastic rocks, Sedimentary rocks |
| | Or-3 | Pyroclastic rocks and lava (basic) |
| | Or-4 | Pyroclastic rocks |
| | Or-5 | Pyroclastic rocks |
| | Or-6 | Pyroclastic rocks and lava |
| Tertiary | Or-7 | Pyroclastic rocks and lava |
| | T-1 | Sedimentary rocks |
| | T-2 | Sedimentary rocks |
| | T-3 | Sedimentary rocks |
| Cretaceous
Jurassic
Precambrian | PJC | Sedimentary rocks, Intrusive rocks |
-
- | | |
|--|----------------------|
| | Alteration zone |
| | Lineament |
| | Curvicular structure |
| | Known mine site |
| | Strike and dip |
| | Anticline |
| | Pond |
| | Remaining snow |

Fig. II-1 Interpretation Map of Landsat False Color Images


CHAPTER 2 GEOLOGICAL INVESTIGATION IN THE COTAHUASI AREA



2-1 Geology and Geological Structure (Fig. II-2)

The geology of the project area consists of Precambrian basement comprising gneissic granite and diorite, the thick Jurassic and Cretaceous systems composed of clastic and calcareous rocks, and widespread Tertiary and Quaternary systems composed mainly of volcanic rocks and intrusive rocks.

Intrusive rocks occur as batholiths of diorite and numerous stocks and dikes of diorite and andesite. These are partly accompanied by skarns and alteration zones.

The formations of Miocene or earlier are heavily folded in general and block faulted. However, neither prominent nor fault is observed in the formations later than Miocene.






LEGEND

Quaternary	Holocene	Alluvium	al	Sand, mud and gravel		
		Mollebamba Volcanic Rocks	Vm	Andesite lava and volcanic ash		
		Volcanic Sediment of Pausa	Vsp	Volcanic ash, sand and gravel		
		Lampa Volcanic Rocks	Vla	Andesite (basaltic), volcanic breccia		
		Moraine Sediment	Mo	Gravel, sand and mud		
			Moraine Sediment	Mo	Gravel, sand and mud	
		Pleistocene	Barroso Group	Upper	Vbu	Acidic tuff
			Barroso Group	Lower	Vbl	Andesite lava and pyroclastic rocks
		Tertiary	Pliocene	Sencca Volcanic Rocks	Vse	Hornblende-biotite dacite lava, welded tuff and tuff
				Huayllillas Formation	Hy	Dacitic tuff (partly pumice bearing)
Alpabamba Formation	Alf			Dacitic tuff, lapilli tuff, tuff breccia and welded tuff (partly with dacite lava or andesite lava)		
Tacaza Group	Tacaza			Tc	Andesitic tuff breccia, Andesite, tuff and dacitic tuff breccia (greenish grey)	
	Huanca Formation			Hc	Andesitic volcanic conglomerate, tuff breccia and tuffaceous sandstone (greenish grey)	
Mesozoic	Cretaceous			Arcurquina Formation	Arf	Limestone and marl with sandstone and chert nodule
		Murca Formation	Mf	Red shale and sandstone with gypsum bearing conglomerate		
		Yura Group	Yu	Quartzite, siliceous sandstone, black shale and alternation of quartzite and shale		
		Socsoni Formation	Sf	Black shale, limestone with sandstone and tuff		
		Chocolate volcanic rocks	Chv	Andesitic tuff breccia, tuff, andesite and tuffaceous sandstone		
Precambrian		Gn	Gneiss, gneissose granite and diorite			
Tertiary	Intrusive Rocks	Stock and Dyke	St	Hornblende andesite, andesite		
		Accha Stock	Ac	Diorite and quartz diorite		
		La Costa Batholith	Lc	Quartz diorite and granodiorite		
Structural Features	Fault			Fault		
	Inferred fault			Inferred fault		
	Anticline			Anticline		
	Syncline			Syncline		
	Geological boundary			Geological boundary		
	Strike and dip of bedding			Strike and dip of bedding		
	Strike and dip of foliation			Strike and dip of foliation		
	Strike and dip of flow structure			Strike and dip of flow structure		
	Mine (working or closed)			Mine (working or closed)		
	Hot spring			Hot spring		
Phase II Survey Area			Phase II Survey Area			
Phase III Survey Area			Phase III Survey Area			

Fig. II-2 Geological Map of the Cotahuasi Area



- Abbreviations
- | | | | | | |
|-----------|---------------------------|------------|--------------------------|-------|-------------------------|
| gn | gneiss, gneissose diorite | qzt | quartzite | ba-an | basaltic andesite lava |
| an-tf-bre | andesitic tuff breccia | vol-cg | volcanic conglomerate | tf-cg | tuffaceous conglomerate |
| an-tf | andesitic tuff | cg | conglomerate | silt | silt |
| tf-ss | tuffaceous sandstone | dc-vol-bre | dacitic volcanic breccia | grv | gravel |
| ls | limestone | an-tf-bre | andesitic tuff breccia | | |
| on | andesite lava | dc-wtf | dacitic welded tuff | | |
| sh | shale | dc-tf | dacitic tuff | | |
| tf | tuff | dc | dacitic lava | | |
| ss | sandstone | rho-tf | rhyolitic tuff | | |

Fig. II-3 Generalized Stratigraphic Column of the Cotahusai Area

(1) Stratigraphy (Fig. II-3)

Rocks of the survey area comprises the Precambrian Basement, the Jurassic, Cretaceous, and Tertiary systems, intrusive rocks through these Basement and systems, and the Quaternary System which is widespread in the Altiplano highlands.

(2) Intrusives

The intrusive rocks occurring in the survey area comprise Coast batholith, (CB), Accha stock (Di) and andesite stocks and dikes (An).

(3) Chemical Composition of Rocks

Whole rock chemical analysis (covering 13 constituents) was made for 20 rock samples collected during the present field work and normative calculations were also performed under CIPW classification. The results of the analysis and normative calculations are presented in Fig. II-4, Fig. II-5, Fig. II-6.

Described below are the results of analysis of volcanic rocks found in the Survey area according to a simple method of classification of the type of volcanic rocks on the basis of the SiO_2 and $Na_2O + K_2O$ contents (classification method of Middlemost, E.A.K. (1972)) (Fig. II-4).




Fig. II-4 $SiO_2 - (Na_2O + K_2O)$ Variation Diagram of Volcanic Rocks

Fig. II-5 illustrates the relationship between K_2O and MgO based on the results of the chemical analysis of rocks (13 constituents). The diagram of Fig. II-5 shows that the MgO content tends to decrease with an increase of K_2O , but the rhyolitic rocks of the Alapabamba Formation (Al) have a lower MgO content than other rocks and a correlation between the MgO content and K_2O content cannot be established. A sample (Gb-56) from the Senca Formation (Vse) is a rhyolitic rock and exhibits a facies similar to that of the rhyolitic rocks of the Alapabamba Formation, but it shows a higher MgO content than the rhyolitic rock of the Alapabamba Formation.




Fig. II-5 K_2O - MgO Variation Diagram of Volcanic Rocks

With respect to intrusive rocks, whole rock chemical analysis was made for two samples of the Coast batholith (CB) and two samples of the Accha stock. All these samples are granitic in composition and a triangular diagram of quartz - (albite + anorthite) - orthoclase was prepared on the basis of the classification of Bateman et al., (1963) selected from among the various classifications of felsic igneous rocks (Fig. II-6).

The four samples can be classified as granodiorites. The Coast batholith in the south of the survey area tends to contain a slightly larger quantity of orthoclase.




Fig. II-6 Normative Quartz - Orthoclase - (Albite + Anorthite) Triangular Digram of Some Igneous Rocks

(4) Geological Structures

The geological structure of the Survey Area is highly characterized by the remarkable folds and faults formed in the Jurassic and Cretaceous rocks by the orogeny of the Andes. In the Tertiary formations strata show a gentle dip locally and small faults are observable. Judging from their scale, the strata in the Tertiary System do not seem to have undergone the effects of a vehement tectonic movement (Fig. II-2).

The folds found in the Jurassic and Cretaceous rocks include a large-scale syncline having a $N70^{\circ} W$ axis in the Socosani Formation occurring in the south of Mt. Tanisca, an E-W syncline found in the

Yura Group west of the village of Velinga on the Cotahuasi River, NW-SE anticline and syncline, N-S anticlines and NE-SW anticlines in that part of the Arcurquina Formation extending from Lancacolla west of the village of Cotahuasi to Taurisma to the northeast.

The folds observed in the Tertiary rocks include small-scale anticlines and synclines with a NNW-SSE axis found in the Tacaza Group west of the village of Carcana in the north of detailed survey area B. No other folds are found in the strata formed after Cretaceous age.

Faults are developed in the formation of the middle Miocene or earlier. The major faults trend, generally in the NW-SE direction including a NW-SE trending fault passes by the village of Saima in the south of the detailed survey area A, an E-W fault in the valley of Palmadeas to the southeast, an NW-SE fault extending from a point east of Lamapa via the south of San Jose de Ushua to the vicinity of Mt. Tanisca and an NW-SE fault running along a valley south of Mina Picha. These faults have throws estimated to be over 1,000 m. Substantially parallel to these large-scale faults are other faults, such as those running across the northeast of Bitama, Mina Luicho and Puca Puca, all found in the west of the survey area, and those passing the valley of Ccolapa and the Chuquibamba River.

NNW-SSE faults are found to the west of Colta, to the east of Pomacocha and in the valley of Anirca northeast of Mina de Huayllura. NE-SW faults are located in a valley east of Quechualla on the Cotahuasi River, east of the village of Cotahuasi, near Puica in the northeast of the survey area and around Sumana east of Puica. All these NNW-SSE and NE-SW faults are on a relatively small scale and some of these have cut the Alfabamba Formation. The NW-SE and NE-SW orientations of fold axes and faults are regionally in substantially consistent with the NW-SE and NE-SW lineaments and the directions of arrangement of circular structures extracted from LANDSAT images. This presumably reflects the basement structure of the survey area.

2-2 Mineralization and Alteration

The mineralization and alteration in the Cotahuasi area as follows.


The alteration zones and mineralized zones in the project area are shown in Fig. II-7. These zones are distributed mainly in the Jurassic and Cretaceous rocks and in the Tacaza Group of the Tertiary. Major among them are a) Mina Pararapa, b) Minas de Huayllura (East of Tanisca), c) West of Tanisca, d) Mina Luicho, e) Mina Picha, f) South of Maran, g) Oyolo and h) Pirca.



The alteration zone can be grouped according to the type of alteration into 1) white alteration with silicification, 2) brown alteration with silicification and argillization and 3) brown alteration contaminated with ferrous oxide.

Mineralization includes vein type mineralization of quartz veins bearing gold and silver; mineralization of gold and silver found in the zones of contamination with ferrous oxide along fracture zones and joints; contact metasomatic mineralization of gold, silver, copper, lead, zinc and magnetite occurring in skarns found in the area of contact between intrusive rocks and limestones; and pyritization of dissemination type observable in and around intrusive rock bodies.

The mineralized zones with confirmed outcrops of gold-silver bearing quartz veins include a) Mina Pararapa, Copacahuana Mine in the Minas de Huayllura b) and f) the south of Maran. In Mina Luicho the mineralized zone is locally accompanied by quartz veinlets which lack continuity.

Table II-1 show the scale, host rock, and characteristics of alteration and mineralization of each zone.





LEGEND

Alteration zone		minerals	
	mainly silicification (qz ± al ± kn)	qz	quartz
	silicification + argillization	pl	plagioclase
	others (iron oxides stained zone)	or	orthoclase
	skarn	al	alunite
	samples	kn	koolinite
	mineralization zone	hal	halloysite
	mine	sr	sericite
	hot spring	mon	montmorillonite
	Batholith granitic rocks	sr/mon	sericite-montmorillonite mixed layer
	Stock dioritic rocks	dia	diaspore
	Stock and dyke andesitic rocks	mar	mordenite
	lineament (Landsat)	and	andradite
	fault	cpx	clinopyroxene
	lineament (aerial photograph)	ca	calcite
	Curvicular structure (Landsat)	py	pyrite
	Curvicular structure (aerial photograph)	mg	magnetite
		goe	goethite
		scs	scorodite
		jar	jarosite

Fig. II-7 Location Map of Alteration and Mineralized Zones


Table II-1 List of Alteration and Mineralization Zones (Main)



Name	Location	Scale	Host rock	Alteration	Mineralization
Mina Pararapa	Approx. 20 km North-northeast of Cotahuasi	1 km x 2.5 km	Andesite dikes (An) and Andesitic volcanic rocks (Tacaza Group)	<ul style="list-style-type: none"> Brown altered zone contaminated by ferrous oxide Hydrothermal alteration consisting primarily of silicification (quartz + halloysite, cb-11 quartz + montmorillonite cb-13) The andesite dikes are chloritized and contaminated by pyrites. 	<ul style="list-style-type: none"> Mineralization of gold and silver in N70°W and 80°NE gold-silver bearing quartz veins Quartz vein being prospected by tunnels is 0.5 to 1.5 m wide and 1.3 km long. According to the mine data, the grade of gold ore is Au, max. 40 g/ton and average 4.6 g/ton. The average grades of gold and silver ores contained in quartz dikes of 80 cm in width are Au 4.6 g/ton and Ag 288 g/ton.
Minas de Huayllura (east of Tonisca)	Approx. 30 km west-southwest of Cotahuasi	1 ~ 2 km x 10 km	Quartzite (Yura Group)	<ul style="list-style-type: none"> Brown altered zone contaminated by ferrous oxide Hydrothermal alteration accompanied by quartz veinlets (quartz + sericite Bb-10 quartz + diaspare + sericite + jarosite Bb-9) 	<ul style="list-style-type: none"> Mineralization exists in oxidized zones occurring along fracture zones and joints and in gold-silver bearing quartz veins Abandoned small-scale adits are found at several locations Analysis of the quartz veins of Mine Copacahuana shows the grades of Au and Ag are 4.6 g/ton and 288 g/ton.
West of Tonisca	Approx. 33 km west-southwest of Cotahuasi and approx. 23 km east-southeast of Pausa	1.5 km x 4 km	Andesitic volcanic rocks (Tacaza Group)	<ul style="list-style-type: none"> Essentially hydrothermal alteration of quartz + alunite and quartz + sericite + montmorillonite Relationship between both types of alteration is not clear. 	<ul style="list-style-type: none"> Conspicuous mineralization not observable. Analysis of altered rocks containing clay minerals revealed the content of Au 0.06 g/ton and Ag 0.31 g/ton.
Mina Luicho	Approx 5 km northeast of Pausa	1 km x 2 km	Quartzite (Yura Group)	<ul style="list-style-type: none"> Silicification around quartz veinlets and contamination with ferrous oxide. (quartz + scorodite (Ae-22)) 	<ul style="list-style-type: none"> Gold-silver bearing quartz veinlets and contamination with ferrous oxide observable. Orientation of the veinlets is not clear. A dozen or so small-scale stopes are scattered. Spot samples from the stopes showed Au 26.0 g/ton, Ag 116.1 g/ton and Au 9.7 g/ton, Ag 30.2 g/ton.
Pirca	Approx. 15 km south of Pirca	2 km x 5 km	Andesite and pyroclastic rocks (Tacaza Group and lower part of Barroso Group)	<ul style="list-style-type: none"> Mostly hydrothermal alteration of quartz + alunite. 	<ul style="list-style-type: none"> Contaminated with goethite near Pirca. Vein-type mineralization not observable. Analysis of samples from the zone contaminated with goethite showed the grade of Ag 0.6 to 1.6 g/ton.
South of Maran	Approx. 15 km south-southeast of Pausa	1.5 km x 3.5 km	Andesitic tuff to tuff breccia (Chocolate Formation)	<ul style="list-style-type: none"> Silicification around quartz veinlets, veinlike silicification and contamination with ferrous oxide. 	<ul style="list-style-type: none"> Gold-silver bearing quartz veins about 10 cm wide and contamination with limonite around fracture zones are observable at several locations. Analysis of samples from Pyrite quartz veins (5 to 10 cm wide) showed the grades of Au 1.1 g/ton and Ag 0.3 g/ton. There are many abandoned old adits. In downstream areas river sediments are washed for gold.
Oyolo	Approx. 20 to 25 km northeast of Pausa	2 km x 8 km	Dacitic pyroclastic rocks (Tacaza Formation to Alpabamba Formation)	<ul style="list-style-type: none"> Hydrothermal alteration (primarily argillization). In the northeast Cristobalite + silualte and in the southwest quartz + sericite 	<ul style="list-style-type: none"> Locally accompanied by contamination with ferrous oxide (grade: Ag 0.3 g/ton) Conspicuous mineralization not observable.
Mina Picha	Approx. 22 km west-southwest of Cotahuasi	Small ore bodies scattered in area of approx. 1 km x 2 km	Limestone (Arcurquina Formation) and diorite (stock)	<ul style="list-style-type: none"> Skarn type metamorphism (garnet (andradite) - calcite-quartz) 	<ul style="list-style-type: none"> Skarn type Mineralization of Cu, Pb, Zn, Ag and Au. Ore bodies are small lens-shaped bodies. Analysis of mineral ores consisting mainly of galena and sphalerite showed the following grades: Au 7.7 g/ton, Ag 170 g/ton, Cu 1.38%, Pb 23.4%, Zn 21.6%

2-3 Geochemical Prospecting

Geochemical prospecting in the area was conducted with the purpose of detecting geochemical anomalies caused by mineralization and of obtaining basic data for mineral exploration in the next phase of the project.

In the prospecting, stream sediments were collected and six elements, gold (Au), silver (Ag), arsenic (As), copper (Cu), lead (Pb) and zinc (Zn) were analyzed as indicators of mineralization. The results are shown in Fig. II-8 and Table II-2.





- LEGEND**
- Au Ag Anomaly Zone of Elements and Anomalous Elements
 - Z1, Z2 Anomaly Zone of Principal Components and Anomalous Components
 - Maran : Name of Anomaly Zone

Fig. II-8 Geochemical Interpretation Map (Composite Data)

Table II-2 List of Geochemical Anomaly Zones

Anomalous zone	Location	Scale (km × km)	Indicator element						Remarks	
			Au	Ag	As	Cu	Pb	Zn		
- Cotahuashi Area -										
1	Mina Pararapa	Approx. 20 km NNW of Cotahuashi	8 × 9	++	++	+	-		Au>Ag	
2	Quechualla	Along Cotahuashi River in the Center of the Survey Area	11 × 14							
	Picha	Northern part of Anomalous zone		++	++	+	-	+	-	Ag>Au
	Quechualla	Along Cotahuashi River		++	++	-	+	-		Au>Ag
	Abandonada	South of Quechualla		+	+	-	+	-	++	
	Tanisea	Western part of Anomalous zone		++	+			+		
3	Huayjo	Approx. 10 km south of Quechualla	2 × 10	++	+	-	++	+	-	
4	Maran	Approx. 10 km south-east of Pausa		++	-	+	++	+	-	Cu>Au
5	Mina Luicho	Approx. 4 km north-east of Pausa	4 × 4	++	++		++	+	-	Au>Ag, Cu
6	Marcabamba	Marcabamba in the north-east of the survey area	3 × 6	+	++		-	-	+	
7	Velinga	Approx. 15 km west-southwest of Cotahuashi	4 × 7	+		+	-		-	Au>As
8	Alca	Approx. 15 km northeast of Cotahuashi	4 × 10	+	+				-	
9	Huaynacotas	Approx. 6 km northwest of Alca	3 × 5	+					++	
10	Colta	Approx. 15 km north-northeast of Colta	3 × 6	+	+				-	Ag>Au
11	Pirca	Approx. 13 km south of Pausa	4 × 4						+	
12	Taurisma	Approx. 3 km north of Cotahuashi	1.5 × 2	-	-				-	
13	Cerro Kiura	Northeastern margine of the Survey Area	1.5 × 4	+						
14	Pausacocha	Approx. 10 km west of Quechualla	2.5 × 2.5	+						
15	Oyolo	Approx. 20 km northeast of Pausa	1.5 × 2	+						
16	San Sebastian Sacraca	Approx. 7 km north of Pausa	1.5 × 2	+						
17	Salamanca	Approx. 10 km southeast of Nevada Sollman	1.5 × 2	+						
- Orcopampa Area -										
18	Orcopampa	Central part of the Orcopampa Area	10 × 6	(++)	(++)				-	+
19	Orcopampa North	Approx. 5 km northeast of Orcopampa	4 × 6	(++)	(++)					