

巻末資料 3 ホーリングコアスケッチ (縮尺1:200)

MJVB-1 (1)

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
0		Yellowish gray, weathered, fine-banded, broken schist (0.00 - 7.15 m), broken quartz contained in some place.	Limonite in cleavage and schistosity.
< 70			
< 60		Several white/light gray quartz veins/veinlets (7.15 - 10.70 m, thickness 1 - 10 cm) in light gray fine-grain sandstone and gray schist, lower sandstone is cut by some quartz veinlets.	Limonite & pyrite disseminated in quartz veins/veinlets.
< 40			
< 75			
< 40			
10		Fine-grain light gray sandstone (10.70 - 14.70 m), cut by several quartz veinlets.	Disseminated sulfide in some place, chloritization and sericitization.
< 65			
< 60			
< 35		Sandstone (14.70 - 17.00 m), containing some quartz veins/veinlets (thickness 0.5 - 10 cm).	Chloritization, some sulfide.
< 40			
< 70			
< 45			
< 45		Quartz zone (17.00 - 17.90 m): sandstone with quartz network and quartz breccia.	Disseminated sulfide, chloritization.
< 70			
< 45		Sandstone (17.90 - 19.20 m), containing quartz veinlets (18.80 m, thickness 1 cm).	Disseminated sulfide, chloritization.
< 70			
< 65		Dark gray schist (19.20 - 21.80 m), containing thin layer of sandstone (19.80 - 20.20 m), and injected by several quartz veinlets.	Pyrite disseminated.
20		Fine-grain light gray sandstone (21.80 - 26.30 m), cut by several quartz veinlets.	Disseminated sulfide, some place chloritization and sericitization.
< 40			
< 75-85		Dark gray schist (26.30 - 28.10 m), containing white quartz vein/veinlet at 26.30 & 27.15 m.	Pyritization, sericitization and chloritization.
< 70			
< 75		Quartz zone (28.10 - 28.55 m): mixture of quartz and schist, and white quartz vein (28.40 - 28.55 m).	Chloritization, pyritization (locally).
30		Dark gray schist (28.55 - 35.90 m), injected by several quartz veinlets.	Pyritization, silicification, and some place chloritization.
< 65			
< 60			
< 60		Quartz zone in dark gray schist (35.90 - 37.00 m): quartz, quartz breccia and dark gray schist.	Weak chloritization and sericitization.
< 80		Dark gray/black schist (37.00 - 42.00 m), injected by quartz veinlets (41.40 m, 2 cm).	Weak chloritization and sericitization, rarely pyritization.
40		Light gray quartzitic sandstone (42.00 m -), injected by several quartz veinlets (1 - 5 mm). Two quartz veinlets of 1.5 and 1.0 cm thick at 42.15 m and 44.70 m.	In shear zone and quartz veins: pyritization, chloritization and sericitization. In quartzitic sandstone: weak alteration.
< 45			
< 50			
< 65		Quartz zone (42.56 - 43.00 m): mixture of quartzitic sandstone, quartz and quartz breccia.	
50			

MJVB-1 (2)


Depth (m)

Drill Log

Geological Description

Mineralization & Alteration

50




< 45
< 45
< 37
< 45-60
< 45
< 30

Light gray quartzitic sandstone (- 60.25 m), containing several white/light gray quartz veins/veinlets (50.23 m, thickness 3 cm; 50.70 and 52.60 m, 1 cm; 53.40 m, 2 quartz veinlets thickness 0.1 cm/1 cm; 56.37 m, 1-3 cm; 59.07 m, 1 cm and 59.58 - 59.63 m, thickness 5 cm).

Pyritization, sericitization and chloritization.

60



< 50
< 60

Alternation of fine-grain light gray sandstone and black schist (60.25 - 64.85 m), cut by quartz veinlets (60.42 m, 4 cm).


Pyritization, sericitization, chloritization and silicification.

< 50
< 40
< 40
< 60
< 60

Dark gray/black locally dark green schist (64.85 - 67.20 m), containing some white/light gray quartz veinlets (64.85 and 65.50 m, thickness 1 cm and 65.80 m, 5 cm).

Pyritization, sericitization, chloritization and silicification.

70




< 30-65

Quartz zone (67.20 - 70.45 m): mixture of quartz, quartz network, quartz breccia (white/light gray) and quartzitic sandstone.

Sericitization, chloritization and silicification. Pyrite disseminated in quartz zone.

80



< 45
< 47
< 60

Dark gray schist (70.45 - 82.45 m), injected by several quartz veinlets in schistosity. (71.25 m quartz network thickness 5 cm in schistosity).

Sericitization, silicification, strong pyritization and some place chloritization.

< 55



< 77
< 50
< 45
< 70

Dark gray schist (82.45 - 86.00 m), containing several white/light gray quartz veins/veinlets and quartz zone (82.45 m, quartz zone thickness 25 cm; mixture of quartz, schist and quartz breccia; 83.43, 84.21, 84.60 and 85.15 m, quartz veinlets 5 cm).

Sericitization, silicification, and some place chloritization. Pyrite disseminated in quartz veins/veinlets.

< 60




< 60

Alternation of fine-grain light gray sandstone and black schist (86.00 - 87.85 m), cut by quartz veinlets (1 cm).

Weak chloritization and sericitization, rarely pyritization.

< 30-60



< 30-60

Dark gray/black (some place greenish gray) schist (87.85 - 91.50 m), injected by quartz veinlets.

Chloritization, sericitization, pyritization and silicification.

< 45
< 25-75




< 30-70
< 65

Dark gray/black (some place greenish gray) schist (91.50 - 99.00 m), injected by quartz veinlets and containing several white/light gray quartz veins and quartz zones (91.60, 92.15 and 92.45 m - quartz veins of complicated form, thickness 10 cm; 93.20 - 93.45 m quartz zone 25 cm. 94.15 - 94.37 m quartz zone, 94.50 - 94.76 m quartz zone, 95.52 - 95.58 m quartz veinlet, 95.65 m quartz veinlet 3 cm, 96.00 - 96.15 m quartz zone, 96.40 - 96.54 m quartz veinlet 3 cm, 97.10 - 97.85 m quartz zone, 98.30 - 98.94 m quartz zone).

Strong chloritization, sericitization, pyritization and silicification. Pyrite & arsenopyrite in quartz veins & quartz zones.

< 85-90
< 80-90
< 70-90















< 70-90

Quartz zone (99.00 - 100.00 m): mixture of psammitic sandstone, quartz, quartz network and quartz breccia.








Strong chloritization, sericitization, pyritization and silicification.

MJVB-1 (3)

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
100		< 45 Gray/greenish gray psammitic sandstone (100.00 - 102.10 m), containing 2 quartz veinlets (101.80 m, thickness 2 - 4 cm).	Pyritization, chloritization, sericitization and silicification.
		< 35 < 65 < 55 < 70-80 < 60-70 < 50-60 Gray/greenish gray/dark gray/black schist (102.10 - 110.80 m), some place psammitic, containing several white/light gray quartz zones, quartz veins/veinlets (102.92, 103.23 and 112.55 m veinlets 2 - 5 cm; 104.88 - 105.35 m quartz vein 20 cm, 105.55 - 105.85 m, 105.90 - 106.02, 106.30 - 106.80, 107.62 - 108.40, 108.58 - 109.00, and 110.35 - 110.80 m quartz zones 10 - 80 cm; mixture of quartz, quartz veins, quartz network, schist and quartz breccia).	Strong pyritization, sericitization and chloritization. Pyrite disseminated in quartz.
110		< 40 Dark gray psammitic sandstone (110.80 - 112.37 m) cut by several quartz veinlets.	Pyritization, sericitization, chloritization and silicification.
		< 60 Dark gray/green schist (112.37 - 113.20 m), injected by several quartz veinlets (112.55 m thickness 5 cm).	Pyritization, sericitization, chloritization and silicification.
120		< 65 Dark gray/green psammite (113.20 - 125.00 m), containing some white/light gray quartz vein/veinlet (121.25 m thickness 10 cm), 124.80 m calcite veinlet.	Pyritization, sericitization, chloritization and silicification.
		< 70 Dark gray/green schist (125.00 - 128.60 m), injected by several quartz veinlets in schistosity.	Sericitization, silicification, strong pyritization and some place chloritization.
130		< 60 Quartz zone (128.60 - 130.00 m): mixture of quartz, quartz network, quartz breccia (white/light gray) and black schist.	Strong chloritization, sericitization and silicification. Pyrite disseminated.
		< 60 Brecciated dark gray/green/ black schist (130.00 - 131.65 m), some place containing brecciated light/gray quartz.	Silicification, weak pyritization and chloritization
		< 70 Dark gray/black quartzitic sandstone/psammite (131.65 - 136.50 m) injected by quartz veinlets.	Weak sericitization, silicification, pyritization and chloritization.
		< 60-70 Quartz zone (136.50 - 137.50 m): mixture of sandstone, quartz, quartz network and quartz breccia.	Pyritization, sericitization and chloritization.
140		< 25 Dark gray/black (some place grayish green) schist (137.50 - 139.40 m) injected by quartz veinlets.	Weak chloritization and sericitization, nearly pyritization.
150		< 25 Gray/light gray quartzitic sandstone/psammite (139.40 m -), injected by quartz veinlets (141.20 m 0.5 cm).	Weak chloritization, sericitization, pyritization and silicification.

MJVB-1 (4)

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
150		<p>Gray/light gray quartzitic sandstone/psammite (- 170.60 m), injected by some white/light gray quartz vein/veinlets (153.60 and 155.75 m veinlets 1 cm and 5 cm, 155.17 - 155.52 m quartz vein 35 cm; 155.63 and 156.40 m veinlets 3 cm and 2 cm).</p>	<p>Weak pyritization, chloritization, sericitization and silicification.</p>
160		<p>Mainly gray psammitic sandstone (170.60 - 172.00 m), some place quartzitic sandstone with black schist, cut by some gray/light gray quartz veinlets.</p>	<p>Weak pyritization, sericitization, chloritization and silicification.</p>
170		<p>Dark gray/black quartzitic sandstone/psammite (172.00 m -) injected by several quartz veinlets (181.30, 181.58, 195.50 and 198.40 m thickness 1 cm each, 190.30 m 5 cm, 191.55 m 3 cm and 199.20 m 0.7 cm)</p>	<p>Weak sericitization, silicification, pyritization and chloritization.</p>
180			
190			
200			

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
200	 <p data-bbox="446 268 526 313">< 40-45</p>		
210	 <p data-bbox="446 716 526 761">< 20 < 60</p>		
220	 <p data-bbox="446 940 526 1030">< 30 < 25-30 < 40 < 45</p>	<p data-bbox="590 851 1005 1030">Gray/light gray quartzitic sandstone/psammite (- 238.70 m), injected by some white/light gray quartz veins/veinlets (213.40, 213.65, 216.30, 219.70, 220.00, 220.10, 220.40, 220.60, 220.90, 221.14 and 221.35 m, 1- 5 cm each, 230.85 - 230.98 m thickness 13 cm).</p>	<p data-bbox="1133 896 1420 963">Weak pyritization, chloritization, sericitization and silicification.</p>
230	 <p data-bbox="446 1321 526 1366">< 40 < 40</p>		
240	 <p data-bbox="446 1635 526 1680">< 45</p>	<p data-bbox="590 1590 1053 1657">Dark gray schist (238.70 - 239.53 m), injected by some gray/light gray quartz veinlets.</p>	<p data-bbox="1133 1590 1436 1657">Weak sericitization, silicification, pyritization and chloritization.</p>
250	 <p data-bbox="446 1724 526 1769">< 45 < 50 < 45</p>	<p data-bbox="590 1747 1037 1836">Gray/light gray quartzitic sandstone/psammite (239.53 - 249.53 m), injected by some white/light gray quartz veinlets (241.80, 246.20 and 246.90 m, 0.5- 1.5 cm each).</p>	<p data-bbox="1133 1769 1436 1836">Weak sericitization, silicification, pyritization and chloritization.</p>
250	 <p data-bbox="446 2016 526 2060">< 60</p>	<p data-bbox="590 1971 1037 2038">Dark gray schist (249.53 m -) injected by several white/light gray quartz veinlets.</p>	<p data-bbox="1133 1971 1436 2038">Weak sericitization, silicification, pyritization and chloritization.</p>

MJVB-1 (6)

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
250			
	< 60	Dark gray schist (- 251.20 m) injected by several white/light gray quartz veinlets.	Weak pyritization, chloritization, sericitization and silicification.
		Gray/light gray quartzitic sandstone/psammite (251.20 - 255.20 m), some place containing schist, injected by some white/light gray quartz vein/veinlets.	Weak pyritization, chloritization, sericitization and silicification.
	< 50	Dark gray schist (255.20 - 256.80 m), cut by several white/light gray quartz veinlets.	Weak pyritization, chloritization, sericitization and silicification.
		Mainly gray fine-grain psammitic/quartzitic sandstone (256.80 - 258.95 m), injected by some white/light gray quartz veinlets.	Weak pyritization, chloritization, sericitization and silicification.
260			
	< 65-70		
	< 40	Dark gray schist (258.95 - 266.10 m), cut by several white/light gray quartz veinlets (261.80 m thickness 1cm).	Weak pyritization, chloritization, sericitization; strong silicification.
	< 60-70		
		Mainly gray psammitic sandstone (266.10-267.45 m), some place quartzitic sandstone with schist.	Weak pyritization, chloritization, sericitization; strong silicification.
270			
	< 45		
	< 60-80	Dark gray schist (267.45 - 275.50 m), cut by several white/light gray quartz veinlets (270.20 m thickness 2 cm, 273.45 and 273.80 m thickness 1 and 1.5 cm).	Weak pyritization, sericitization, chloritization and silicification.
	< 65		
	< 60		
	< 35-40		
280			
	< 60		
	< 55		
	< 40		
	< 45		
		Mainly gray fine-grain psammitic/quartzitic sandstone (275.50 - 300.00 m), some place with dark gray schist (277.00 - 278.70, 280.30 - 280.95, 281.75 - 282.30 and 289.90 - 290.60 m) and injected by several white/light gray quartz veinlets (280.85 m thickness 5 cm, 283.80, 285.40 and 288.95 m thickness 1 cm).	Weak sericitization, silicification, pyritization and chloritization.
290			
	< 50		
300			

MJVB-2 (1)

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
0		Mixture of yellow/light brown/gray broken weathered sandstone/schist (0.00 - 16.00 m), some place with broken quartz.	Limonite in cleavages.
10			Mainly light gray psammite sandstone (16.00 - 25.00 m), some place with schist, contain quartz veinlets.
20			Mainly gray fine-grain psammite/quartzitic sandstone (25.00 m -), some place with dark gray schist (25.00 - 26.20, 44.20 - 44.80 m), injected by several light gray/gray quartz veins/veinlets (29.60 m thickness 1.5 cm; 40.40 - 40.50 m thickness 0.5cm; 42.30, 47.25, 48.15 and 48.40 m thickness 2cm; 47.58 and 47.90 m thickness 5 cm; 44.90 and 45.30 m thickness 10 cm; 48.95 m thickness 1 cm) .
30			
40			
50			

MJVB-2 (2)

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
50		<p>Mainly light gray psammitic/quartzitic sandstone (- 64.20 m), some place with schist, cut by several gray quartz veins/veinlets (51.24 - 51.52 m thickness 28 cm; 51.84, 51.99, 53.47, 53.48, and 56.00 m thickness 1 - 3 cm; 53.40, 54.84, 59.40 and 60.45 m thickness 4 - 6 cm, 58.45 m thickness 8 cm).</p>	<p>Weak pyritization, chloritization and sericitization; strong silicification. Pyrite disseminated in quartz veins/veinlets.</p>
60		<p>Dark gray schist (64.20 - 66.50 m).</p>	<p>Silicification</p>
70		<p>Mainly gray/light gray quartzitic sandstone/psammite (66.50 - 76.88 m), some place with schist, contain some gray quartz veinlets (73.30 m thickness 3 cm, 76.05 m thickness 1 cm).</p>	<p>Weak pyritization, chloritization, sericitization; strong silicification.</p>
80		<p>Quartz zone (76.88 - 77.43 m): mixture of light gray/gray quartz, sandstone, quartz breccia and quartz veinlets.</p>	<p>Chloritization and sericitization; strong silicification. Pyrite and arsenopyrite disseminated in quartz.</p>
90		<p>Mainly gray fine-grain psammitic/quartzitic sandstone (77.43 m -), some place with siliceous schist, injected by several light gray/gray quartz veins/veinlets (77.80 m thickness 1 cm; 79.84 - 79.97 and 86.78 - 86.87 m thickness 10 cm; 81.13 - 81.33 m thickness 20 cm; 80.80, 81.77 - 81.83, 82.12, 82.68, 84.30, 86.52, 94.25 - 94.40, 95.20 - 95.40, and 97.30 m thickness 1 - 15cm).</p>	<p>Silicification, pyritization and chloritization; strong sericitization around quartz zone. Pyrite and arsenopyrite disseminated in quartz veins/veinlets.</p>
100			

MJVB-2 (3)







Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
100	<p>< 35 < 45-60 < 40 < 60-70 < 40 < 40 < 40 < 25 < 45 < 55 < 40-55 < 40 < 40 < 30</p>	<p>Mainly light gray psammitic/quartzitic sandstone (< 114.00 m), some place with schist, cut by several gray quartz veins/veinlets (100.55 m thickness 5 cm; 101.48, 101.95, 102.72, 105.87, 106.30, 107.55, 107.90, 108.64, 109.02, 109.55 - 109.70, 110.20, 110.83, 110.90, 110.20, 111.40, and 111.48 m thickness 1 - 2 cm; 104.75 - 104.90 m thickness 15 cm; 109.37 - 109.44 m thickness 7 cm).</p>	<p>Weak pyritization, chloritization and sericitization; strong silicification.</p>
110	<p>< 25-35 < 40 < 30 < 40-45 < 45 < 30-45 < 30 < 50 < 40-60 < 30-40 < 35 < 15</p>	<p>Mainly light gray psammitic/quartzitic sandstone (114.00 - 118.00 m), some place with schist, cut by several gray quartz veins/veinlets (115.12, 115.45, 115.55, 115.70 - 115.85, 116.00, 117.30, 117.40 and 117.85 m quartz veinlets/networks thickness 1 - 2 cm; 114.40 - 114.50 and 117.00 - 117.10 m quartz veins thickness 10 cm).</p> <p>Mainly gray/light gray quartzitic sandstone/psammite (118.00 - 122.00 m), containing gray quartz veinlets/quartz zones (118.02 - 118.62, 119.13 - 119.42, 120.15 - 120.36 and 120.81 - 121.09 m quartz zones 60, 25, 18 and 25 cm; mixture of quartz, quartz breccias, quartz veins/veinlets and sandstone; 118.77, 119.56 - 119.75 and 121.46 m quartz veinlets/networks 1 - 5 cm).</p>	<p>Weak pyritization, chloritization and sericitization; strong silicification.</p> <p>Pyritization, chloritization, sericitization; strong silicification.</p>
120	<p>< 10 < 15-45 < 30 < 35 < 30</p>	<p>Mainly gray fine-grain psammitic/quartzitic sandstone (122.00 - 134.70 m), injected by several gray quartz veins/veinlets (122.66, 123.05 - 123.15, 126.25, 129.68, 131.00 and 133.80 - 133.88 m quartz veinlets and networks 1 - 2.5 cm; 124.30 - 124.41 m quartz vein 11 cm).</p>	<p>Pyritization, chloritization and sericitization; strong silicification.</p>
130	<p>< 30 < 45</p>	<p>Quartzitic sandstone (134.70 - 139.30 m) with quartz zone (134.70 - 135.40 m and 136.59 - 139.30 m); mixture of gray quartz, sandstone, quartz breccia and quartz network.</p>	<p>Strong silicification, pyritization and chloritization.</p>
140	<p>< 40 < 40 < 50 < 40-60 < 55-60 < 40</p>	<p>Mainly gray fine-grain psammitic/quartzitic sandstone (139.30 - 146.30 m), injected by some gray/light gray quartz veins/veinlets (140.70, 142.40 quartz veinlets 2 cm and 3.5 cm; 141.30 - 141.46 m quartz vein thickness 15 cm).</p> <p>Quartzitic sandstone (146.30 m -) with quartz zone and quartz veinlets (146.45 - 146.66 m and 148.20 m - quartz zones; mixture of gray quartz, sandstone, quartz breccia and quartz network; 146.86, 147.15, 147.28, 147.55 and 147.82 m quartz veinlets thickness 1 - 6 cm).</p>	<p>Silicification, pyritization and chloritization; strong sericitization.</p> <p>Strong silicification, pyritization and chloritization.</p>
150			


MJVB-2 (4)

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
150		Quartz zone (- 152.10 m) in quartzitic sandstone: mixture of gray/white quartz, sandstone, quartz breccia and quartz network.	Strong silicification, pyritization and chloritization.
160	<p>< 50-60</p> <p>< 50</p> <p>< 20</p> <p>< 55</p> <p>< 40</p> <p>< 25-30</p> <p>< 40</p>	<p>Mainly light gray psammitic/quartzitic sandstone (152.10-170.00m), some place with psammitic schist, contain several gray quartz zones, quartz veins/veinlets (154.40 - 155.85 m quartz zone: quartz, quartz breccia, quartz network and sandstone; 156.30, 157.30, 158.40, 159.63, 158.80, 158.90 - 159.00, 162.10, 162.52, 165.70, 167.00, 167.25, 167.40, 168.48 and 169.29 m thickness 1 - 3 cm, 159.00 - 159.60 m quartz vein thickness 60 cm).</p>	<p>Pyritization, chloritization, sericitization; strong silicification. Pyrite disseminated in quartz.</p>
170	<p>< 35</p> <p>< 40</p> <p>< 40</p> <p>< 45</p> <p>< 50</p> <p>< 60</p> <p>< 25-30</p> <p>< 25-30</p>	<p>Mainly gray/dark gray or greenish gray fine-grain psammitic schist (170.00 - 190.00 m), some place quartzitic sandstone, injected by several gray quartz veins/veinlets and networks (181.00, 181.11 - 181.22, 181.22 - 181.32, 181.40, 181.57 - 181.62, 183.30 - 183.35, 184.80 - 184.90, 185.80 - 185.90, 186.10, 187.70, 188.17, 188.60, and 189.75 m thickness 1 - 8 cm).</p>	<p>Pyritization, chloritization and sericitization; strong silicification. 181.25m there is 2 grains Au in quartz veinlet.</p>
180	<p>< 45-50</p> <p>< 35</p> <p>< 60</p> <p>< 55</p> <p>< 45</p> <p>< 45</p> <p>< 50</p>	<p>Mainly gray fine-grain psammitic/quartzitic sandstone (190.00 m -), injected by some gray/light gray quartz veinlets and networks (190.40, 190.90, 195.10, 195.40, 195.95, 197.30 and 199.45 m thickness 1-2 cm; 199.60 and 199.92 m networks thickness 10 cm).</p>	<p>Silicification, pyritization and chloritization; strong sericitization. Pyrite disseminated in quartz.</p>
190	<p>< 30</p> <p>< 30-35</p> <p>< 30-35</p> <p>< 30</p> <p>< 40</p>	<p>Mainly gray fine-grain psammitic/quartzitic sandstone (190.00 m -), injected by some gray/light gray quartz veinlets and networks (190.40, 190.90, 195.10, 195.40, 195.95, 197.30 and 199.45 m thickness 1-2 cm; 199.60 and 199.92 m networks thickness 10 cm).</p>	<p>Silicification, pyritization and chloritization; strong sericitization. Pyrite disseminated in quartz.</p>
200			


Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
200		<p>Mainly light gray psammitic/quartzitic sandstone (- 216.60 m), some place with psammitic schist, containing several gray/white quartz zones, quartz veins /veinlets. (200.20 - 200.30, 201.35 - 201.67, 202.10 - 202.30, 207.35 - 208.20 and 208.40 - 209.08 m quartz zones; quartz, quartz breccia, quartz network and sandstone, 204.05, 205.85, 206.50, 206.70, 206.90, 213.47, 213.53 and 215.30 m veinlets sickness 1 - 3 cm, 207.20 - 207.35 and 216.50 - 216.60 m networkks, 215.00, 215.35 and 216.32 m quartz veins thickness 10 cm).</p>	<p>Pyritization; strong chloritization, sericitization, silicification. Pyrite, arsenopyrite and pyrrhotite disseminated in quartz.</p>
210		<p>Mainly gray/dark gray fine-grain schist (216.60 - 240.00 m), some place quartzitic sandstone, injected by several gray quartz zones, quartz veins/veinlets and quartz networks (231.60 - 231.85 and 232.70 - 233.00 m networks, 233.55, 233.76, 234.15 and 235.35 m veinlets thickness 2-5 cm, 234.90 - 235.00 m quartz zone; quartz network, quartz breccia and quartzitic/psammitic sandstone).</p>	<p>Pyritization, chloritization and sericitization; strong silicification.</p>
220		<p>Mainly dark gray schist (240.00 - 247.70 m), some place psammitic/quartzitic sandstone, injected by some gray/light gray quartz veinlets (thickness 1-3 cm).</p>	<p>Silicification, pyritization and chloritization.</p>
230		<p>Mainly psammite (247.70 m -), some place quartzitic/psammitic sandstone, injected by quartz veinlets (250.00 m thickness 1 cm).</p>	<p>Pyritization, chloritization and sericitization; strong silicification.</p>
240		250	


MJVB-2 (6)

Depth (m)	Drill Log	Geological Description	Mineralization & Alteration
250	 <p>< 40 < 50-55 < 45 < 40 < 45 < 50-60 < 40 < 30-40 < 45</p>	<p>Mainly light gray psammitic/quartzitic sandstone (- 261.50 m), some place with psammite, containing several gray/white quartz zones, quartz veinlets (251.30 - 251.60, 252.28 - 252.60, 254.16 - 254.80 and 256.67 - 256.79 m quartz zone; quartz, quartz breccia, quartz network, parallel quartz veinlets and psammite, 251.95, 252.00 and 253.76 m veinlets thickness 1 - 3 cm).</p>	<p>Weak pyritization, strong chloritization, sericitization and silicification.</p>
260	 <p>< 35 < 40 < 30</p>	<p>Mainly gray/dark gray fine-grain schist (261.50 - 267.45 m), some place quartzitic sandstone, injected by quartz veinlets (264.35 m thickness 2 cm).</p>	<p>Weak pyritization, chloritization, sericitization, strong silicification.</p>
270	 <p>< 40-60 < 40-50 < 40 < 45 < 45</p>	<p>Mainly light gray psammitic/quartzitic sandstone (267.45 - 273.70 m), some place with psammite, contain gray/white quartz veinlets (268.23 m veinlets thickness 1 cm).</p>	<p>Weak pyritization, chloritization, sericitization, strong silicification.</p>
280	 <p>< 30 < 40 < 40 < 40-50 < 35-40</p>	<p>Alternation of gray/dark gray fine-grain quartzitic/psammitic sandstone and psammite (273.70 - 282.80 m), containing several white/gray quartz veins/veinlets and quartz network (277.05, 277.34, 277.70, 278.37, 280.06, 280.05 and 281.40 m veinlets thickness 1 - 6 cm, 278.90 - 279.10 m quartz vein thickness 15 cm).</p>	<p>Pyritization, chloritization and sericitization, strong silicification.</p>
290	 <p>< 40 < 60 < 25-30 < 30 < 40-50 < 35</p>	<p>Mainly dark gray schist (282.80 - 296.35 m), some place psammitic/quartzitic sandstone, containing several gray/light gray quartz zones, quartz veinlets and networks (283.33 - 283.52, 290.00 - 290.72 and 291.45 - 292.05 m quartz zones; quartz, quartz breccia, quartz veinlets, quartz networks and schist, 288.10 - 288.15 and 288.35 - 288.40 m networks 0.5 - 1 cm, 292.41, 292.87, 293.05, 293.40 and 294.28 m veinlets thickness 1 - 7 cm).</p>	<p>Silicification, pyritization and chloritization. Pyrite and chalcocopyrite disseminated in quartz.</p>
300	 <p>< 35</p>	<p>Mainly psammite and quartzitic sandstone (296.35 - 300.00 m), some place schist, injected by quartz veinlets (297.45 m veinlet thickness 1 cm).</p>	<p>Pyritization, chloritization and sericitization; strong silicification.</p>





A — A'










A — A'




- LEGEND**
- STRATIGRAPHY**
- Quaternary Q Alluvial deposits
 - Devonian Dm1 Mia Le Formation
 - D-bb Roc Bun Formation
 - C3ts1 Lower Than So Formation
 - Cambrian Cmd Ma Dong Formation
 - 1: alternation of schist and sandstone
 - 2: mainly composed of sandstone
- OTHERS**
- Fault
 - Dip and strike of bedding / schistosity
 - Dip and strike of quartz vein
 - Anticlinal axis
 - Synclinal axis
 - Adit / Inclined shaft
 - Drill hole
 - A — A' Geologic section line



A — A'





Scale 1:10,000
 0 200 400 600 800 1000





- LEGEND**
- STRATIGRAPHY**
- C3ts1 Lower Than Sa Formation
 - Ma Dang Formation
 - 1: alternation of schist and sandstone
 - 2: mainly composed of sandstone
 - Cmd
- OTHERS**
- Fault
 - Dip and strike of bedding / schistosity
 - Dip and strike of quartz vein
 - Anticlinal axis
 - Adit / inclined shaft
 - Shaft / Pit
 - A—A Geologic section line

A — A'













LEGEND

- A002 Location and number of ore samples
- A003 Location and number of rock-chip samples






- LEGEND**
- Location and number of rock samples
 - A : Ore sample
 - P : Ore microscopy
 - T : Thin section
 - F : Fluid inclusion
 - X : X-ray diffraction analysis



LEGEND

- A002 Location and number of ore samples
- A003 Location and number of rock chip samples



- LEGEND
- Location and number of rock samples
 - A : Ore sample
 - P : Ore microscopy
 - T : Thin section
 - F : Fluid inclusion
 - X : X-ray diffraction analysis

