

Apc.22 Diagraphie géologique des trous de forages à circulation

inverse (RC) dans le Secteur de Kékoro

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-5.0m: Surface soil yellowish brown on the top, brown on the bottom, this layer contains white clay and some quartz fragments	51-1	46	-	-
		51-2	42	-	-
		51-3	40	-	-
		51-4	31	-	-
		51-5	42	-	-
6.0	5.0-6.0m: Crust and Carapace reddish brown colored, with iron nodules	51-6	204	-	-
10	6.0-12.0m: Saprolite A greenish grey colored saprolite	51-7	221	-	-
		51-8	404	-	-
		51-9	33	-	-
15	12.0-36.0m: Saprolite B greenish grey colored saprolite, including a lot of chips of weathered meta-greywacke	51-10	11	6	-
		51-11	7	-	-
		51-12	7	-	-
		51-13	7	-	-
		51-14	6	-	-
		51-15	5	-	-
		51-16	3	-	-
		51-17	6	-	-
		51-18	5	-	-
		51-19	5	-	-
		51-20	7	7	-
		51-21	29	-	-
20		51-22	5	-	-
		51-23	5	-	-
		51-24	3	-	-
		51-25	4	-	-
		51-26	21	-	-
		51-27	6	-	-
		51-28	3	-	-
		51-29	8	-	-
		51-30	5	6	-
		51-31	8	-	-
25		51-32	3	-	-
		51-33	2	-	-
		51-34	7	-	-
		51-35	11	-	-
		51-36	5	-	-
36.0	36.0-47.0m: Meta-sandstone dark grey colored meta-greywacke, with disseminated pyrite, with joint filling pyrite, total amount of pyrite is 1 % to 2 %	51-37	4	-	-
40		51-38	4	-	-
		51-39	2	-	-
		51-40	4	2	-
		51-41	9	-	-
		51-42	10	-	-
		51-43	3	-	-
		51-44	7	-	-
		51-45	3	-	-
47.0	47.0-51.0m: Peritic schist dark grey colored peritic schist, with disseminated pyrite, with schistosity filling pyrite total amount of pyrite is 2 %	51-46	7	-	-
50		51-47	13	-	-
		51-48	12	-	-
		51-49	16	-	-
51.0	51.0-57.0m: Meta-sandstone dark grey colored meta-greywacke, with disseminated pyrite, total amount of pyrite is 0 % to 1 %	51-50	11	10	-
55		51-51	7	-	-
		51-52	5	-	-
		51-53	7	-	-
		51-54	5	-	-
57.0	57.0-60.0m: Peritic schist dark grey colored peritic schist, with disseminated pyrite, with schistosity filling pyrite total amount of pyrite is 1 % to 2 %	51-55	6	-	-
60		51-56	8	-	-
		51-57	10	-	-
		51-58	12	-	-
		51-59	10	-	-
		51-60	14	7	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0	0.0-1.0m: Surface soil	52-1	26	-	-
3.0	brown colored, with laterite nodules	52-2	96	-	-
5	2.0-3.0m: Carapace reddish brown colored, with a lot of lateritic nodules	52-3	31	-	-
		52-4	28	-	-
		52-5	11	-	-
		52-6	385	-	-
10	3.0-14.0m: Saprolite A greenish grey colored saprolite	52-7	19	-	-
		52-8	32	-	-
		52-9	19	-	-
		52-10	10	7	-
		52-11	302	-	-
15	14.0-22.0m: Saprolite B greenish grey colored saprolite, including chips of weathered meta-greywacke	52-12	23	-	-
		52-13	27	-	-
		52-14	22	-	-
		52-15	14	-	-
		52-16	11	-	-
		52-17	9	-	-
20	22.0-26.0m: Saprolite B greenish grey colored saprolite, including chips of silicified? meta-greywacke (quartz schist?)	52-18	10	-	-
		52-19	6	-	-
		52-20	16	8	-
		52-21	5	-	-
		52-22	19	-	-
25	26.0-34.0m: Saprolite B greenish grey colored saprolite, including chips of weathered meta-greywacke	52-23	8	-	-
		52-24	5	-	-
		52-25	24	-	-
		52-26	11	-	-
		52-27	17	-	-
		52-28	8	-	-
30	34.0-38.0m: Alternation beds of meta-sandstone and peritic schist dark grey colored rock, including 1 % to 4 % of disseminated pyrite	52-29	6	-	-
		52-30	7	12	-
		52-31	23	-	-
		52-32	50	-	-
		52-33	30	-	-
		52-34	55	-	-
35	38.0-45.0m: Meta-sandstone meta-sandstone with 1 % to 4 % of disseminated pyrite	52-35	23	-	-
		52-36	15	-	-
		52-37	8	-	-
		52-38	18	-	-
40	water	52-39	10	-	-
		52-40	7	21	-
		52-41	37	-	-
		52-42	218	-	-
		52-43	0	-	-
45	45.0-52.0m: Alternation beds of meta-sandstone and peritic schist dark grey colored rock, with schistosity filling and disseminated pyrite, 1% to 4% in peritic schist, 1% in meta-sandstone	52-44	42	-	-
		52-45	1	-	-
		52-46	5	-	-
50	52.0-56.0m: Meta-sandstone meta-sandstone with great amount of disseminated pyrite, more than 4 % ?	52-47	31	-	-
		52-48	5	-	-
		52-49	4	-	-
		52-50	8	1	-
		52-51	1	-	-
		52-52	0	-	-
55	56.0-60.0m: Peritic schist peritic schist with schistosity filling pyrite and disseminated pyrite, more than 4 % ?	52-53	0	-	-
		52-54	0	-	-
		52-55	0	-	-
60		52-56	0	-	-
		52-57	0	-	-
		52-58	2	-	-
		52-59	0	-	-
		52-60	1	2	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
0.0-3.0	0.0-3.0m: Carapace reddish brown colored soil with laterite nodules	53-1	17	-	-
		53-2	14	-	-
		53-3	17	-	-
3.0-20.0	3.0-20.0m: Saprolite A greenish brown colored saprolite	53-4	23	-	-
		53-5	14	-	-
		53-6	19	-	-
		53-7	13	-	-
		53-8	15	-	-
		53-9	61	-	-
		53-10	37	28	-
		53-11	32	-	-
		53-12	6	-	-
		53-13	5	-	-
		53-14	7	-	-
		53-15	10	-	-
		53-16	4	-	-
		53-17	40	-	-
		53-18	36	-	-
		53-19	71	-	-
		20.0-39.0	20.0-39.0m: Saprolite B greenish grey colored saprolite, including chips of meta-greywacke and of perititic schist	53-20	16
53-21	3			-	-
53-22	4			-	-
53-23	8			-	-
53-24	14			-	-
53-25	19			-	-
53-26	13			-	-
53-27	12			-	-
53-28	8			-	-
53-29	14			-	-
53-30	30			4	-
53-31	1			-	-
53-32	6			-	-
53-33	2			-	-
53-34	6			-	-
53-35	4			-	-
53-36	5			-	-
53-37	8			-	-
53-38	7			-	-
53-39	2			-	-
39.0-57.0	39.0-57.0m: Perititic schist dark grey colored schist, with schistosity filling and disseminated fine grained pyrite, 1% to 3% ?	53-40	4	4	-
		53-41	20	-	-
		53-42	19	-	-
		53-43	14	-	-
		53-44	21	-	-
		53-45	17	-	-
		53-46	20	-	-
		53-47	17	-	-
		53-48	19	-	-
		53-49	41	-	-
		53-50	17	18	-
57.0-58.0	57.0-58.0m: Meta-sandstone grey colored rock, with 1 % to 3 % ? pyrite	53-51	25	-	-
		53-52	30	-	-
		53-53	66	-	-
		53-54	78	-	-
		53-55	37	-	-
		53-56	47	-	-
58.0-60.0	58.0-60.0m: Perititic schist dark grey colored schist, with schistosity filling and disseminated fine grained pyrite, 1% to 3% ?	53-57	36	-	-
		53-58	8	-	-
		53-59	11	-	-
		53-60	13	14	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-2.0m: Carapace reddish brown colored soil with laterite nodules	54-1	15	-	-
		54-2	5	-	-
10	2.0-4.0m: Mottled clay brown colored clay 4.0-18.0m: Saprolite A yellowish brown colored powder	54-3	13	-	-
		54-4	12	-	-
		54-5	6	-	-
		54-6	3	-	-
		54-7	4	-	-
		54-8	3	-	-
		54-9	6	-	-
		54-10	9	7	-
		54-11	38	-	-
		54-12	15	-	-
		54-13	12	-	-
		54-14	4	-	-
		54-15	22	-	-
		54-16	24	-	-
15	18.0-30.0m: Saprolite B greenish grey to yellowish brown colored saprolite, including chips of greenish grey colored meta-sediment	54-17	10	-	-
		54-18	5	-	-
		54-19	5	-	-
		54-20	3	3	-
		54-21	2	-	-
		54-22	1	-	-
		54-23	3	-	-
		54-24	30	-	-
		54-25	0	-	-
		54-26	11	-	-
20	30.0-32.0m: Meta-sandstone grey colored rock, with disseminated pyrite, 1 % to 4 %	54-27	89	-	-
		54-28	9	-	-
		54-29	3	-	-
		54-30	11	6	-
		54-31	167	-	-
		54-32	6	-	-
		54-33	6	-	-
		54-34	11	-	-
		54-35	4	-	-
		54-36	4	-	-
25	32.0-33.0m: Quartz schist dark grey colored rock, with strong dissemination of pyrite	54-37	9	-	-
		54-38	2	-	-
		54-39	5	-	-
		54-40	4	6	-
		54-41	19	-	-
		54-42	3	-	-
		54-43	4	-	-
		54-44	3	-	-
		54-45	12	-	-
		54-46	3	-	-
30	33.0-38.0m: Meta-sandstone grey colored, weakly silicified, with 1% to 4% of disseminated pyrite, with many fragments of quartz	54-47	0	-	-
		54-48	3	-	-
		54-49	1	-	-
		54-50	0	-	-
		54-51	6	-	-
		54-52	33	-	-
		54-53	13	-	-
		54-54	14	-	-
		54-55	4	-	-
		54-56	2	-	-
35	38.0-45.0m: Peritic schist dark grey colored schist, with schistosity filling and disseminated fine grained pyrite	54-57	4	-	-
		54-58	9	-	-
		54-59	2	-	-
		54-60	2	6	-
		54-61	2	6	-
		54-62	2	6	-
		54-63	2	6	-
		54-64	2	6	-
		54-65	2	6	-
		54-66	2	6	-
40	45.0-50.0m: Dioritic intrusion ? dark grey colored diorite, with dissemination of pyrite and pyrrhotite	54-67	2	6	-
		54-68	2	6	-
		54-69	2	6	-
		54-70	2	6	-
		54-71	2	6	-
		54-72	2	6	-
		54-73	2	6	-
		54-74	2	6	-
		54-75	2	6	-
		54-76	2	6	-
45	50.0-60.0m: Meta-sandstone grey colored rock, with strong dissemination of pyrite, with siliceous peritic schist	54-77	2	6	-
		54-78	2	6	-
		54-79	2	6	-
		54-80	2	6	-
		54-81	2	6	-
		54-82	2	6	-
		54-83	2	6	-
		54-84	2	6	-
		54-85	2	6	-
		54-86	2	6	-
50		54-87	2	6	-
		54-88	2	6	-
		54-89	2	6	-
		54-90	2	6	-
		54-91	2	6	-
		54-92	2	6	-
		54-93	2	6	-
		54-94	2	6	-
		54-95	2	6	-
		54-96	2	6	-
55		54-97	2	6	-
		54-98	2	6	-
		54-99	2	6	-
		54-100	2	6	-
		54-101	2	6	-
		54-102	2	6	-
		54-103	2	6	-
		54-104	2	6	-
		54-105	2	6	-
		54-106	2	6	-
60		54-107	2	6	-
		54-108	2	6	-
		54-109	2	6	-
		54-110	2	6	-
		54-111	2	6	-
		54-112	2	6	-
		54-113	2	6	-
		54-114	2	6	-
		54-115	2	6	-
		54-116	2	6	-

Column (m)	Description	Sample No.	Au				
			ppb	ppb	ppb		
5	0.0-5.0m: Carapace reddish brown colored soil with laterite nodules at the depth of 1m to 2m	55-1	0	-	-		
		55-2	16	-	-		
		55-3	17	-	-		
		55-4	18	-	-		
		55-5	19	-	-		
10	5.0-32.0m: Saprolite A 5.0m to 11.0m: brown colored silty soil	55-6	32	-	-		
		55-7	83	-	-		
		55-8	180	-	-		
		55-9	329	-	-		
		55-10	229	493	-		
		55-11	241	-	-		
		15	11.0m to 19.0m: yellowish light grey colored silty soil	55-12	40	-	-
				55-13	26	-	-
				55-14	39	-	-
				55-15	58	-	-
55-16	55			-	-		
55-17	134			-	-		
55-18	33			-	-		
55-19	21			-	-		
20	19.0m to 27.0m: grey to brown colored silty soil	55-20	7	39	-		
		55-21	32	-	-		
		55-22	25	-	-		
		55-23	25	-	-		
		55-24	53	-	-		
		55-25	27	-	-		
		55-26	42	-	-		
		55-27	36	-	-		
		25	27.0m to 32.0m: greenish grey to light grey to brown colored silty soil	55-28	30	-	-
				55-29	27	-	-
55-30	62			50	-		
55-31	95			-	-		
55-32	33			-	-		
30	32.0-47.0m: Saprolite B greenish light grey colored fine grained soil, with chips of rock fragments (diameter: 5mm)			55-33	36	-	-
				55-34	37	-	-
				55-35	46	-	-
				55-36	21	-	-
				55-37	27	-	-
		55-38	251	-	-		
		55-39	539	-	-		
		55-40	210	193	-		
		55-41	69	-	-		
		55-42	44	-	-		
35	47.0-54.0m: Peritic schist dark grey colored schist, with schistosity filling and disseminated fine grained pyrite, 1% to 3% with small amount of Fe-oxides	55-43	29	-	-		
		55-44	27	-	-		
		55-45	11	-	-		
		55-46	71	-	-		
		55-47	10	-	-		
		55-48	112	-	-		
		55-49	478	-	-		
		55-50	525	714	-		
		55-51	1,607	71	69		
		55-52	494	-	-		
40	54.0-60.0m: Alternation beds dark grey colored to greenish light grey colored alternation beds of meta-sandstone and peritic schist with schistosity filling and disseminated fine grained pyrite, 2% to 3% or more, including arsenopyrite with quartz veinlets	55-53	69	-	-		
		55-54	144	-	-		
		55-55	95	-	-		
		55-56	1,538	78	103		
		55-57	33	-	-		
		55-58	41	-	-		
		55-59	315	-	-		
		55-60	166	158	-		
		45					
50							
55							
60							

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 15 20 25 30 35 40 45 50 55 60	0.0-4.0m: Crust and carapace reddish brown colored, hard material with lateritic nodules	56-1	55	-	-
		56-2	35	-	-
		56-3	512	-	-
		56-4	83	-	-
	4.0-8.0m: Carapace reddish brown colored carapace including white clay and lateritic nodules	56-5	158	-	-
		56-6	889	-	-
		56-7	117	-	-
		56-8	120	-	-
	8.0-32.0m: Mottled clay brown to yellowish brown colored soft material, with no lateritic nodules	56-9	137	-	-
		56-10	50	91	-
		56-11	63	-	-
		56-12	68	-	-
		56-13	103	-	-
		56-14	177	-	-
		56-15	190	-	-
		56-16	330	-	-
		56-17	95	-	-
		56-18	109	-	-
		56-19	76	-	-
		56-20	253	614	-
		56-21	166	-	-
		56-22	405	-	-
		56-23	2,550	2,854	3,189
		56-24	117	-	-
		56-25	61	-	-
		56-26	16,470	11,451	8,709
		56-27	53	-	-
		56-28	172	-	-
		56-29	1,505	2,306	1,303
		56-30	1,052	1,016	-
		56-31	123	-	-
		56-32	613	-	-
32.0-38.0m: Saprolite A greenish grey colored fine to midium grained saprolite	56-33	135	-	-	
	56-34	359	-	-	
	56-35	172	-	-	
	56-36	304	-	-	
	56-37	210	-	-	
	56-38	130	-	-	
38.0-48.0m: Saprolite B greenish grey to bluish grey colored midium grained material including rock chips	56-39	259	-	-	
	56-40	122	-	-	
	56-41	433	-	-	
	56-42	230	-	-	
	56-43	328	-	-	
	56-44	48	-	-	
	56-45	137	-	-	
	56-46	138	-	-	
	56-47	174	-	-	
	56-48	333	-	-	
48.0-60.0m: Granodiorite light grey to greenish grey colored granodiorite, with medium grained facies and fine grained facies, partly chloritized, with dissemination of pyrite and traces of arsenopyrite, sulfide= 0 % to 2 %	56-49	2,450	1,880	2,263	
	56-50	6,940	7,430	-	
	56-51	3,170	3,650	3,840	
	56-52	662	-	-	
	56-53	227	-	-	
	56-54	232	-	-	
	56-55	191	-	-	
	56-56	181	-	-	
	56-57	150	-	-	
	56-58	71	-	-	
	56-59	115	-	-	
60	water	56-60	127	104	-

KRC-57

Coordination: N500 E200
 Elevation: 331m
 Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au				
			ppb	ppb	ppb		
5	0.0-5.0m: Alluvial sediments brown colored soil, includig gravels	57-1	44	-	-		
		57-2	174	-	-		
		57-3	56	-	-		
		57-4	35	-	-		
		57-5	415	-	-		
6.0	5.0-6.0m: Carapace reddish brown colored carapace including lateritic nodules and quartz fragments	57-6	19,948	19,817	16,971		
10	6.0-10.0m: Mottled clay brownish yellowish colored clay with no lateritic nodules	57-7	284	-	-		
		57-8	95	-	-		
		57-9	151	-	-		
15	10.0-30.0m: Saprolite A greenish grey to greenish yellow colored fine grained saprolite	57-10	83	85	-		
		57-11	349	-	-		
		57-12	1,130	1,012	1,474		
		57-13	1,099	996	1,371		
		57-14	814	-	-		
		57-15	1,540	2,244	2,194		
		57-16	372	-	-		
		57-17	388	-	-		
		57-18	166	-	-		
		57-19	206	-	-		
		57-20	372	214	-		
		57-21	1,132	1,602	1,166		
		57-22	395	-	-		
25		57-23	87	-	-		
		57-24	219	-	-		
		57-25	42	-	-		
		57-26	78	-	-		
		57-27	1,823	3,980	2,606		
		57-28	403	-	-		
		57-29	882	-	-		
		57-30	384	814	1,008		
		30	30.0-36.0m: Saprolite B greenish grey colored fine grained powder, including chips of meta-sandstone that is stained by Fe-oxides	57-31	426	-	-
				57-32	104	-	-
57-33	233			-	-		
57-34	100			-	-		
57-35	97			-	-		
57-36	282			-	-		
35	36.0-60.0m: Meta-sandstone dark grey colored meta-sandstone, with fine grained pyrite dissemination, with pyrite veinlets, total amount of sulfide = 1 % to 2 %, with traces of quartz fragments and carbonate fragments, with traces of rock chips of green schist	57-37	166	-	-		
		57-38	60	-	-		
		57-39	196	-	-		
		57-40	362	225	-		
		57-41	128	-	-		
		57-42	100	-	-		
		57-43	29	-	-		
		57-44	102	-	-		
		57-45	41	-	-		
		57-46	34	-	-		
		57-47	29	-	-		
		57-48	35	-	-		
		57-49	23	-	-		
		57-50	64	37	-		
40		57-51	40	-	-		
		57-52	24	-	-		
		57-53	78	-	-		
		57-54	42	-	-		
		57-55	46	-	-		
		57-56	63	-	-		
		57-57	74	-	-		
		57-58	53	-	-		
		57-59	88	-	-		
		57-60	52	39	-		
45							
50							
55							
60							

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
0.0-3.0m	Carapace reddish brown colored, with iron nodules	42-1	40	-	-
		42-2	36	-	-
		42-3	31	-	-
3.0-13.0m	Saprolite A greenish grey colored powder with no rock fragments	42-4	49	-	-
		42-5	111	-	-
		42-6	99	-	-
		42-7	16	-	-
		42-8	16	-	-
		42-9	16	-	-
		42-10	15	3	-
		42-11	30	-	-
		42-12	39	-	-
		42-13	15	-	-
13.0-27.0m	Saprolite B greenish grey colored saprolite, including a lot of chips of peritic schist	42-14	23	-	-
		42-15	14	-	-
		42-16	21	-	-
		42-17	25	-	-
		42-18	12	-	-
		42-19	9	-	-
		42-20	1	9	-
		42-21	16	-	-
		42-22	3	-	-
		42-23	27	-	-
		42-24	20	-	-
		42-25	36	-	-
		42-26	37	-	-
		42-27	16	-	-
27.0-34.0m	Peritic schist dark grey colored peritic schist, with joint filling pyrite and disseminated pyrite, total amount of pyrite is 0 % to 2 %	42-28	13	-	-
		42-29	13	-	-
		42-30	15	21	-
		42-31	12	-	-
		42-32	5	-	-
		42-33	8	-	-
		42-34	25	-	-
		42-35	0	-	-
		42-36	4	-	-
		42-37	33	-	-
34.0-36.0m	Quarz diorite ? dark grey colored, medium grained rock	42-38	18	-	-
		42-39	10	-	-
		42-40	10	6	-
		42-41	26	-	-
		42-42	54	-	-
		42-43	11	-	-
		42-44	35	-	-
		42-45	0	-	-
		42-46	50	-	-
		42-47	47	-	-
36.0-54.0m	Meta-sandstone dark grey colored meta-graywacke, with disseminated pyrite, with fracture filling pyrite, total amount of pyrite is 1 % to 3 %	42-48	4	-	-
		42-49	9	-	-
		42-50	16	27	-
		42-51	0	-	-
		42-52	10	-	-
		42-53	3	-	-
		42-54	13	-	-
		42-55	4	-	-
		42-56	52	-	-
		42-57	34	-	-
54.0-55.0m	Alternation beds dark grey colored alternation beds of meta-graywacke and peritic schist, with disseminated pyrite, with fracture filling pyrite, total amount of pyrite is 1 % to 3 %	42-58	28	-	-
		42-59	53	-	-
		42-60	47	32	-
		42-60	47	32	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
0.0-3.0	0.0-3.0m: Carapace reddish brown colored, with laterite nodules	43-1	2	-	-
		43-2	16	-	-
		43-3	3,093	8,154	829
3.0-18.0	3.0-18.0m: Saprolite A greenish grey colored, soft and fine grained powder	43-4	110	-	-
		43-5	25	-	-
		43-6	73	-	-
		43-7	81	-	-
		43-8	9	-	-
		43-9	15	-	-
		43-10	184	236	-
		43-11	38	-	-
		43-12	46	-	-
		43-13	27	-	-
		43-14	56	-	-
		43-15	24	-	-
		43-16	39	-	-
		43-17	21	-	-
		43-18	22	-	-
18.0-28.0	18.0-28.0m: Saprolite B greenish grey colored saprolite, including chips of sandstone schist of greenish grey colored	43-19	34	-	-
		43-20	23	34	-
		43-21	18	-	-
		43-22	59	-	-
		43-23	42	-	-
		43-24	50	-	-
		43-25	68	-	-
		43-26	162	-	-
		43-27	42	-	-
		43-28	56	-	-
		43-29	41	-	-
28.0-60.0	28.0-60.0m: Sandstone schist grey to greenish grey colored schistose meta-sandstone, with schistosity filling pyrite and disseminated pyrite, total amount of pyrite = 1 % to 3 %, including meta-volcanics ? rock chip of 43-30 is siliceous tuff with strong dissemination of pyrite	43-30	27	11	-
		43-31	22	-	-
		43-32	21	-	-
		43-33	75	-	-
		43-34	108	-	-
		43-35	78	-	-
		43-36	9	-	-
		43-37	57	-	-
		43-38	68	-	-
		43-39	59	-	-
		43-40	51	50	-
		43-41	22	-	-
		43-42	10	-	-
		43-43	17	-	-
		43-44	17	-	-
		43-45	19	-	-
		43-46	39	-	-
		43-47	26	-	-
		43-48	11	-	-
		43-49	17	-	-
		43-50	12	9	-
43-51	61	-	-		
43-52	18	-	-		
43-53	6	-	-		
43-54	8	-	-		
43-55	38	-	-		
43-56	41	-	-		
43-57	49	-	-		
43-58	30	-	-		
43-59	14	-	-		
43-60	17	22	-		

Column (m)	Description	Sample No.	Au				
			ppb	ppb	ppb		
5	0.0-3.0m: Carapace reddish brown colored soil with laterite nodules	44-1	1	-	-		
		44-2	0	-	-		
		44-3	287	-	-		
	10	3.0-23.0m: Saprolite A greenish brown colored soft powder, including some fragments of quartz veinlets	44-4	53	-	-	
			44-5	179	-	-	
			44-6	46	-	-	
			44-7	17	-	-	
			44-8	28	-	-	
			44-9	122	-	-	
			44-10	18	10	-	
			44-11	24	-	-	
			44-12	47	-	-	
			44-13	26	-	-	
			44-14	103	-	-	
			44-15	145	-	-	
			44-16	291	-	-	
			44-17	81	-	-	
			44-18	64	-	-	
			44-19	781	-	-	
			20		44-20	58	36
	44-21	2			-	-	
	44-22	5			-	-	
	44-23	13			-	-	
	44-24	136			-	-	
	44-25	10			-	-	
	44-26	0			-	-	
	44-27	0			-	-	
	44-28	33			-	-	
	44-29	20			-	-	
	25	23.0-30.0m: Saprolite B greenish grey colored saprolite, including chips of meta-sandstone	44-30	26	7	-	
44-31			11	-	-		
44-32			60	-	-		
44-33			25	-	-		
44-34			23	-	-		
44-35			12	-	-		
44-36			66	-	-		
44-37			116	-	-		
44-38			86	-	-		
44-39			40	-	-		
44-40			11	2	-		
30		44-41	10	-	-		
		44-42	18	-	-		
		44-43	57	-	-		
		44-44	573	-	-		
		44-45	46	-	-		
		44-46	62	-	-		
		44-47	13	-	-		
		44-48	11	-	-		
		44-49	32	-	-		
		44-50	15	114	29		
35	30.0-60.0m: Meta-sandstone dark grey to greenish dark grey colored meta-greywacke, including trace of pyrite, 0 % to 1 %, including meta-volcanics ? 34m to 36m: fracture zone, rock chips are brown colored and oxidized 43m to 46m: fracture zone, rock chips are brown colored and oxidized	44-51	65	-	-		
		44-52	22	-	-		
		44-53	71	-	-		
		44-54	34	-	-		
		44-55	35	-	-		
		44-56	233	-	-		
		44-57	431	-	-		
		44-58	126	-	-		
		44-59	63	-	-		
		44-60	131	27	5		
40		44-56	233	-	-		
		44-57	431	-	-		
		44-58	126	-	-		
		44-59	63	-	-		
		44-60	131	27	5		
		45		44-56	233	-	-
				44-57	431	-	-
				44-58	126	-	-
				44-59	63	-	-
				44-60	131	27	5
50				44-56	233	-	-
				44-57	431	-	-
				44-58	126	-	-
				44-59	63	-	-
				44-60	131	27	5
		55		44-56	233	-	-
				44-57	431	-	-
				44-58	126	-	-
				44-59	63	-	-
				44-60	131	27	5
60				44-56	233	-	-
				44-57	431	-	-
				44-58	126	-	-
				44-59	63	-	-
				44-60	131	27	5
		60.0		44-56	233	-	-
				44-57	431	-	-
				44-58	126	-	-
				44-59	63	-	-
				44-60	131	27	5

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0	0.0-1.0m: Carapace	45-1	163	-	-
2.0	reddish brown colored soil with laterite nodules	45-2	50	-	-
		45-3	27	-	-
	1.0-2.0m: Mottled clay	45-4	12	-	-
	brown yellow colored clay	45-5	9	-	-
		45-6	24	-	-
	2.0-11.0m: Saprolite A	45-7	493	-	-
	greenish grey colored, fine grained powder	45-8	88	-	-
		45-9	32	-	-
		45-10	26	26	-
		45-11	16	-	-
		45-12	80	-	-
		45-13	65	-	-
		45-14	28	-	-
		45-15	19	-	-
		45-16	0	-	-
		45-17	8	-	-
		45-18	2	-	-
		45-19	22	-	-
		45-20	34	28	-
		45-21	20	-	-
		45-22	17	-	-
		45-23	14	-	-
		45-24	17	-	-
		45-25	54	-	-
		45-26	45	-	-
		45-27	18	-	-
		45-28	37	-	-
		45-29	34	-	-
		45-30	34	42	-
		45-31	38	-	-
		45-32	61	-	-
		45-33	91	-	-
		45-34	105	-	-
		45-35	73	-	-
		45-36	56	-	-
		45-37	39	-	-
		45-38	168	-	-
		45-39	71	-	-
		45-40	183	136	-
	40.0-60.0m: Meta-sandstone ?	45-41	29	-	-
	dark grey to greenish dark grey colored rock with disseminated pyrite	45-42	33	-	-
		45-43	31	-	-
	meta-greywacke ?	45-44	28	-	-
	meta-diorite ??, including meta-volcanics ?	45-45	19	-	-
		45-46	19	-	-
		45-47	49	-	-
		45-48	66	-	-
		45-49	55	-	-
		45-50	46	44	-
		45-51	66	-	-
		45-52	91	-	-
		45-53	48	-	-
		45-54	42	-	-
		45-55	41	-	-
		45-56	177	-	-
		45-57	90	-	-
		45-58	30	-	-
	rock chips of 45-59 are meta-graywacke or peritic schist	45-59	15	-	-
		45-60	22	9	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-7.0m: Carapace reddish brown colored hard soil with iron nodules	48-1	22	-	-
		48-2	13	-	-
		48-3	25	-	-
		48-4	38	-	-
		48-5	92	-	-
		48-6	38	-	-
		48-7	149	-	-
10	7.0-32.0m: Saprolite A reddish brown to brown colored, fine grained powder, gradually change to green saprolite oxide zone	48-8	125	-	-
		48-9	131	-	-
		48-10	199	172	-
		48-11	214	-	-
		48-12	325	-	-
		48-13	119	-	-
		48-14	110	-	-
		48-15	165	-	-
		48-16	212	-	-
		48-17	218	-	-
		48-18	183	-	-
		48-19	304	-	-
		48-20	626	239	-
		48-21	144	-	-
		48-22	326	-	-
		48-23	399	-	-
		48-24	184	-	-
		48-25	569	-	-
		48-26	892	583	1,029
		48-27	999	480	891
		48-28	1,098	4,114	857
		48-29	824	-	-
		48-30	583	589	-
		48-31	1,754	480	1,303
48-32	1,162	960	686		
35	32.0-45.0m: Saprolite A greenish grey colored, fine grained saprolite reduced zone	48-33	781	-	-
		48-34	618	-	-
		48-35	228	-	-
		48-36	294	-	-
		48-37	159	-	-
		48-38	134	-	-
		48-39	275	-	-
		48-40	393	836	-
		48-41	2,004	3,600	1,200
		48-42	1,656	3,497	2,331
		48-43	2,966	6,926	3,840
45	45.0-58.0m: Saprolite B greenish grey colored, medium grained powder including chips of meta-sandstone ("black tuff")	48-44	6,221	1,303	5,897
		48-45	1,346	1,646	1,851
		48-46	1,242	-	651
50		48-47	393	-	-
		48-48	2,060	-	1,714
		48-49	2,184	-	2,057
		48-50	677	442	-
		48-51	520	-	-
		48-52	661	-	-
		48-53	410	-	-
		48-54	986	-	-
		48-55	926	-	-
		48-56	249	-	-
55	58.0-60.0m: Meta-sandstone ("black tuff") dark grey colored, fine grained, massive rock including anhedral plagioclase grains, with dissemination of pyrite, amount of pyrite is 3% to 4%	48-57	280	-	-
		48-58	261	-	-
		48-59	436	-	-
60	60.0	48-60	257	329	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 14.0	0.0-14.0m: Carapace reddish brown colored hard soil with iron nodules not indurated, including white clay	49-1	5	-	-
		49-2	0	-	-
		49-3	80	-	-
		49-4	13	-	-
		49-5	73	-	-
		49-6	82	-	-
		49-7	45	-	-
		49-8	45	-	-
		49-9	108	-	-
		49-10	145	0	-
		49-11	44	-	-
		49-12	38	-	-
		49-13	48	-	-
		49-14	5,384	746	0
15 20 25 30 35 36.0	14.0-36.0m: Saprolite A brownish grey to greenish grey colored, fine grained powder	49-15	110	-	-
		49-16	81	-	-
		49-17	85	-	-
		49-18	234	-	-
		49-19	267	-	-
		49-20	86	90	-
		49-21	78	-	-
		49-22	73	-	-
		49-23	70	-	-
		49-24	70	-	-
		49-25	167	-	-
		49-26	53	-	-
		49-27	83	-	-
		49-28	49	-	-
		49-29	688	-	-
		49-30	9	81	-
		49-31	227	-	-
		49-32	163	-	-
		49-33	90	-	-
		49-34	102	-	-
		49-35	57	-	-
		49-36	18	-	-
36.0 40 44.0	36.0-44.0m: Saprolite B greenish grey colored, medium grained powder including diorite chips	49-37	104	-	-
		49-38	937	1,066	617
		49-39	435	-	-
		49-40	157	68	-
		49-41	225	-	-
		49-42	147	-	-
		49-43	121	-	-
		49-44	86	-	-
		49-45	135	-	-
		49-46	103	-	-
45 50 55 60	44.0-60.0m: Diorite (or Granodiorite) dark grey colored, medium grained diorite or granodiorite with disseminated pyrite, amount of pyrite is 0 % to 1 %	49-47	190	-	-
		49-48	139	-	-
		49-49	75	-	-
		49-50	71	-	-
		49-51	100	-	-
		49-52	70	-	-
		49-53	75	-	-
		49-54	67	-	-
		49-55	37	-	-
		49-56	22	-	-
		49-57	20	-	-
		49-58	59	-	-
		49-59	37	-	-
		49-60	29	54	-

Column (m)	Description	Sample No.	Au			
			ppb	ppb	ppb	
5 10 13.0	0.0-13.0m: Carapace reddish brown colored, coarse grained carapace, with a lot of iron nodules (diameter: 2-10mm)	84-1	30	-	-	
		84-2	19	-	-	
		84-3	15	-	-	
		84-4	24	-	-	
		84-5	33	-	-	
		84-6	33	-	-	
		84-7	54	-	-	
		84-8	82	-	-	
		84-9	132	-	-	
		84-10	57	73	-	
		84-11	109	-	-	
		84-12	162	-	-	
		84-13	185	-	-	
15 20 25 30 30.0	13.0-30.0m: Saprolite A reddish brown to yellowish brown colored, very fine grained saprolite, with no rock chips	84-14	391	-	-	
		84-15	502	-	-	
		84-16	971	-	-	
		84-17	471	-	-	
		84-18	600	-	-	
		84-19	482	-	-	
		84-20	863	245	-	
		84-21	84	-	-	
		84-22	99	-	-	
		84-23	204	-	-	
		84-24	203	-	-	
		84-25	666	-	-	
		84-26	9,460	9,531	10,491	
		84-27	10,290	8,777	9,189	
		84-28	886	-	-	
		84-29	144	-	-	
		84-30	110	125	-	
35 40 39.0	30.0-39.0m: Saprolite B yellowish brown colored, medium grained saprolite, with a lot of chips of strongly weathered sandstone (diameter: 5-20mm)	84-31	1,440	1,063	1,200	
		84-32	309	-	-	
		84-33	424	-	-	
		84-34	8,120	7,577	9,771	
		84-35	22,100	17,109	21,360	
		84-36	11,470	8,811	9,360	
		84-37	2,699	2,366	2,880	
		84-38	416	-	-	
		84-39	182	-	-	
		84-40	104	144	-	
45 50 55 60 60.0	39.0-41.0m: Meta-sandstone brown to dark gray colored, strongly weathered meta-sandstone, with oxidized sandstone and strongly weathered peritic schist	84-41	899	-	-	
		84-42	182	-	-	
		84-43	444	-	-	
		84-44	654	-	-	
		84-45	147	-	-	
		41.0-60.0m: Granodiorite fine grained granodiorite (plagioclase>>biotite >hornblende), with weak dissemination of pyrite, contents of pyrite is less than 1%, fresh, chloritization is not seen	84-46	53	-	-
			84-47	1,104	25	0
			84-48	728	-	-
			84-49	67	-	-
			84-50	174	99	-
			84-51	151	-	-
84-52	61		-	-		
84-53	58		-	-		
84-54	81		-	-		
84-55	57		-	-		
84-56	45		-	-		
84-57	94	-	-			
84-58	48	-	-			
84-59	130	-	-			
84-60	32	117	-			

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 13.0	0.0-13.0m: Carapace reddish brown colored carapace, including a lot of iron nodules	85-1	19	-	-
		85-2	60	-	-
		85-3	27	-	-
		85-4	26	-	-
		85-5	10	-	-
		85-6	11	-	-
		85-7	45	-	-
		85-8	21	-	-
		85-9	169	-	-
		85-10	3,591	2,518	-
		85-11	307	-	-
		85-12	242	-	-
		15 15.0	13.0-15.0m: Mottled clay brown colored clay, including some lateritic nodules	85-13	207
85-14	122	-		-	
20 25 30 35 40 45 46.0	15.0-46.0m: Saprolite A 15.0-38.0m: brown to pinkish brown colored, fine grained saprolite 38.0-46.0m: brownish yellow colored, soft saprolite	85-15	181	-	-
		85-16	180	-	-
		85-17	303	-	-
		85-18	193	-	-
		85-19	42	-	-
		85-20	123	1,649	446
		85-21	24	-	-
		85-22	64	-	-
		85-23	43	-	-
		85-24	32	-	-
		85-25	171	-	-
		85-26	31	-	-
		85-27	36	-	-
		85-28	38	-	-
		85-29	73	-	-
		85-30	112	107	-
		85-31	83	-	-
85-32	43	-	-		
85-33	123	-	-		
85-34	73	-	-		
85-35	53	-	-		
85-36	103	-	-		
85-37	134	-	-		
85-38	97	-	-		
85-39	147	-	-		
85-40	119	62	-		
85-41	10,710	9,806	10,971		
85-42	704	-	-		
85-43	575	-	-		
85-44	78	-	-		
85-45	211	-	-		
85-46	69	-	-		
50 50.0	46.0-50.0m: Saprolite B greenish gray colored saprolite, including chips of weathered intrusive	85-47	49	-	-
85-48		38	-	-	
85-49		30	-	-	
85-50		39	37	-	
55 58.0	50.0-58.0m: Granodiorite greenish dark gray colored, medium grained granodiorite, chloritized, with pyrite dissemination, contents of pyrite is 1% in volume	85-51	68	-	-
		85-52	55	-	-
		85-53	29	-	-
		85-54	162	-	-
		85-55	501	-	-
		85-56	1,533	1,783	2,366
		85-57	103	-	-
60 60.0	58.0-60.0m: Meta-andesite ? greenish gray colored meta-andesite, shistosed, with 1% pyrite dissemination	85-58	91	-	-
85-59		79	-	-	
85-60		60	72	-	

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
4.0	0.0-4.0m: Alluvial sediments dark gray colored sediments, mostly clay material, including a lot of organic material	50-1	78	-	-
		50-2	57	-	-
		50-3	38	-	-
		50-4	0	-	-
6.0	4.0-6.0m: Mottled clay brown colored clay, including some lateritic nodules	50-5	23	-	-
		50-6	86	-	-
		50-7	21	-	-
		50-8	22	-	-
27.0	6.0-27.0m: Saprolite A yellowish colored, fine grained saprolite	50-9	25	-	-
		50-10	40	17	-
		50-11	54	-	-
		50-12	28	-	-
		50-13	27	-	-
		50-14	35	-	-
		50-15	52	-	-
		50-16	57	-	-
		50-17	44	-	-
		50-18	31	-	-
		50-19	35	-	-
		50-20	17	32	-
		50-21	51	-	-
		50-22	51	-	-
		50-23	40	-	-
		50-24	30	-	-
		50-25	27	-	-
		50-26	59	-	-
		50-27	63	-	-
37.0	27.0-37.0m: Saprolite B dark green to greenish light gray colored, chloritized saprolite, with strong pyrite dissemination (2-3% in volume)	50-28	20	-	-
		50-29	15	-	-
		50-30	26	569	60
		50-31	40	-	-
		50-32	28	-	-
		50-33	94	-	-
		50-34	250	-	-
		50-35	145	-	-
		50-36	223	-	-
		50-37	369	-	-
45.0	37.0-45.0m: Meta-sandstone greenish to gray colored, fine grained meta-sandstone, chloritized, with sulfide (pyrite-arsenopyrite) dissemination (1-2% in volume), including quartz-calcite veinlets, contents of sulfide increase in silicified zone	50-38	57	-	-
		50-39	28	-	-
		50-40	20	27	-
		50-41	8	-	-
		50-42	13	-	-
		50-43	22	-	-
		50-44	6	-	-
		50-45	129	-	-
50					
55					
60					

Column (m)	Description	Sample No.	Au			
			ppb	ppb	ppb	
2.0 5 10 15	0.0-2.0m: Carapace reddish brown colored, with laterite nodules	34-1	99	-	-	
		34-2	58	-	-	
	2.0-15.0m: Saprolite A greenish grey colored, soft and fine grained powder	34-3	124	-	-	
		34-4	253	-	-	
		34-5	92	-	-	
		34-6	355	-	-	
		34-7	155	-	-	
		34-8	279	-	-	
		34-9	247	-	-	
		34-10	157	141	-	
		34-11	815	-	-	
		34-12	741	-	-	
	15.0 20 25 27.0	15.0-27.0m: Saprolite B greenish grey colored saprolite, including chips of weathered rock	34-13	209	-	-
			34-14	294	-	-
34-15		88	-	-		
34-16		208	-	-		
34-17		86	-	-		
34-18		46	-	-		
34-19		209	-	-		
34-20		116	119	-		
34-21		147	-	-		
34-22		174	-	-		
34-23		1,073	980	926		
27.0 30 35 40 45 50 55 60		27.0-60.0m: Meta-sandstone dark grey to black colored meta-greywacke with disseminated pyrite, total amount of pyrite = 0 % to 1 % rock chip of 34-29 is meta-andesite with strong dissemination of pyrite	34-24	186	-	-
	34-25		328	-	-	
	34-26		833	-	-	
	34-27		536	-	-	
	34-28		60	-	-	
	34-29		41	-	-	
	34-30		25	13	-	
	34-31		13	-	-	
	34-32		98	-	-	
	34-33		20	-	-	
	34-34		4	-	-	
	34-35		22	-	-	
	34-36		12	-	-	
	34-37		6	-	-	
34-38	27	-	-			
34-39	111	-	-			
34-40	38	35	-			
34-41	11	-	-			
34-42	74	-	-			
34-43	65	-	-			
34-44	20	-	-			
34-45	9	-	-			
34-46	15	-	-			
34-47	15	-	-			
34-48	75	-	-			
34-49	22	-	-			
34-50	24	26	-			
34-51	21	-	-			
34-52	34	-	-			
34-53	35	-	-			
34-54	228	-	-			
34-55	190	-	-			
34-56	95	-	-			
34-57	30	-	-			
34-58	28	-	-			
34-59	25	-	-			
34-60	146	136	-			

Column (m)	Description	Sample No.	Au				
			ppb	ppb	ppb		
2.0	0.0-2.0m: Carapace reddish brown colored soil with laterite nodules	35-1	734	-	-		
		35-2	3,785	2,707	2,777		
5	2.0-12.0m: Saprolite A greenish brown to greenish grey colored powder including no rock chips	35-3	146	-	-		
		35-4	75	-	-		
		35-5	82	-	-		
		35-6	44	-	-		
		35-7	61	-	-		
		35-8	90	-	-		
		35-9	59	-	-		
		35-10	247	291	-		
		35-11	29	-	-		
		35-12	191	-	-		
12.0	12.0-21.0m: Saprolite B greenish grey colored saprolite, including chips of weathered rock	35-13	255	-	-		
		35-14	86	-	-		
		35-15	57	-	-		
		35-16	17	-	-		
		35-17	37	-	-		
		35-18	42	-	-		
		35-19	22	-	-		
		35-20	1,390	75	686		
		35-21	33	-	-		
		35-22	26	-	-		
21.0	21.0-60.0m: Granodiorite light grey colored, fine grained granodiorite or tonalite, including trace of pyrite dissemination, with no alteration	35-23	16	-	-		
		35-24	20	-	-		
		35-25	32	-	-		
		35-26	33	-	-		
		35-27	13	-	-		
		35-28	23	-	-		
		35-29	23	-	-		
		35-30	25	29	-		
		35-31	8	-	-		
		35-32	0	-	-		
41.0	41m to 42m: fracture zone, shear zone, with epidotic alteration, with pyrite dissemination, 0 % to 1 % chip sample of 35-42 is peritic rock disseminated by pyrite	35-33	32	-	-		
		35-34	33	-	-		
		35-35	36	-	-		
		35-36	17	-	-		
		35-37	23	-	-		
		35-38	32	-	-		
		35-39	43	-	-		
		35-40	44	18	-		
		35-41	37	-	-		
		35-42	20	-	-		
42.0		35-43	53	-	-		
		35-44	31	-	-		
		35-45	31	-	-		
		35-46	34	-	-		
		35-47	31	-	-		
		35-48	27	-	-		
		35-49	63	-	-		
		35-50	32	53	-		
		35-51	73	-	-		
		35-52	2,659	2,910	2,811		
52.0	52m to 55m: epidotic alteration, with pyrite dissemination, 0 % to 1 %	35-53	205	-	-		
		35-54	77	-	-		
		35-55	197	-	-		
		35-56	38	-	-		
		35-57	30	-	-		
		35-58	69	-	-		
		35-59	20	-	-		
		35-60	78	92	-		
		55.0		35-60	78	92	-
				35-60	78	92	-

KRC-40

Coordination: N1000 E320

Elevation: 340m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
4.0	0.0-4.0m: Surface soil greenish brown colored silty soil, including iron nodules	40-1	57	-	-
		40-2	125	-	-
		40-3	14	-	-
		40-4	118	-	-
6.0	4.0-6.0m: Carapace reddish brown colored hard, indurated soil with iron nodules	40-5	33	-	-
		40-6	47	-	-
		40-7	1,408	1,466	-
		40-8	151	-	-
8.0	6.0-8.0m: Mottled clay reddish brown colored hard, indurated soil with iron nodules	40-9	115	-	-
		40-10	35	2,201	130
		40-11	0	-	-
		40-12	336	-	-
22.0	14.0-36.0m: Saprolite A brownish grey to greenish grey colored, fine grained powder	40-13	449	-	-
		40-14	699	-	-
		40-15	477	-	-
		40-16	469	-	-
		40-17	1,004	-	686
		40-18	161	-	-
		40-19	123	-	-
		40-20	102	107	-
		40-21	314	-	-
		40-22	604	-	-
40.0	36.0-44.0m: Saprolite B greenish grey colored, medium grained powder including diorite chips	40-23	418	-	-
		40-24	494	-	-
		40-25	164	-	-
		40-26	176	-	-
		40-27	2,290	-	651
		40-28	1,192	-	1,406
		40-29	172	-	-
		40-30	898	1,014	-
		40-31	686	-	-
		40-32	890	-	-
60.0	44.0-60.0m: Diorite (or Granodiorite) dark grey colored, medium grained diorite or granodiorite with disseminated pyrite, amount of pyrite is 0 % to 1 %	40-33	922	850	0
		40-34	1,376	-	-
		40-35	230	-	-
		40-36	210	-	-
		40-37	224	-	-
		40-38	326	-	-
		40-39	174	-	-
		40-40	134	136	-
		40-41	87	-	-
		40-42	1,073	262	377
60.0	44.0-60.0m: Diorite (or Granodiorite) dark grey colored, medium grained diorite or granodiorite with disseminated pyrite, amount of pyrite is 0 % to 1 %	40-43	74	-	-
		40-44	996	1,016	660
		40-45	193	-	-
		40-46	147	-	-
		40-47	531	-	-
		40-48	343	-	-
		40-49	258	-	-
		40-50	112	99	-
		40-51	78	-	-
		40-52	90	-	-
60.0	44.0-60.0m: Diorite (or Granodiorite) dark grey colored, medium grained diorite or granodiorite with disseminated pyrite, amount of pyrite is 0 % to 1 %	40-53	160	-	-
		40-54	209	-	-
		40-55	241	-	-
		40-56	73	-	-
		40-57	166	-	-
		40-58	206	-	-
		40-59	177	-	-
		40-60	401	318	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0 5 10	0.0-2.0m: Carapace brown to reddish brown colored carapace, including a lot of lateritic gravels 2.0-10.0m: Saprolite A greenish gray colored saprolite, fine grained, very soft, including quartz chips	78-1	3	-	-
		78-2	23	-	-
		78-3	7	-	-
		78-4	6	-	-
		78-5	4	0	-
		78-6	4	-	-
		78-7	153	-	-
		78-8	3	-	-
		78-9	2	-	-
		78-10	514	-	-
10.0 14.0	10.0-14.0m: Saprolite B greenish gray colored saprolite, medium grained, including chips of weathered meta-graywacke	78-11	27	-	-
		78-12	26	-	-
		78-13	82	-	-
		78-14	20	-	-
14.0 20 25	14.0-20.0m: Meta-sandstone dark gray colored meta-graywacke, including rounded plagioclase fragment (diameter: 0.5-1 mm), with silicification with arsenopyrite inside quartz veinlet (18-19m) 20.0-30.0m: Peritic schist dark gray colored peritic schist, with pyrite disseminaiton (1-2%)	78-15	3	0	-
		78-16	7	-	-
		78-17	12	-	-
		78-18	4	-	-
		78-19	30	-	-
		78-20	6	-	-
		78-21	12	-	-
		78-22	9	-	-
		78-23	9	-	-
		78-24	20	-	-
30.0 35 40 45 50 55 60	30.0-60.0m: Alternation beds alternation beds of peritic schist and meta- graywacke, with more sulfide in peritic schist (0-1%), with silicified fracture (41-46m) water	78-25	16	11	-
		78-26	8	-	-
		78-27	14	-	-
		78-28	10	-	-
		78-29	4	-	-
		78-30	36	-	-
		78-31	11	-	-
		78-32	9	-	-
		78-33	10	-	-
		78-34	12	-	-
		78-35	10	10	-
		78-36	8	-	-
		78-37	9	-	-
		78-38	5	-	-
		78-39	8	-	-
		78-40	15	-	-
78-41	14	-	-		
78-42	19	-	-		
78-43	161	-	-		
78-44	16	-	-		
78-45	16	8	-		
78-46	8	-	-		
78-47	20	-	-		
78-48	12	-	-		
78-49	23	-	-		
78-50	20	-	-		
78-51	22	-	-		
78-52	33	-	-		
78-53	20	-	-		
78-54	13	-	-		
78-55	12	12	-		
78-56	28	-	-		
78-57	115	-	-		
78-58	46	-	-		
78-59	22	-	-		
78-60	19	-	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 15 20 25 30 35 40 45	0.0-3.0m: Carapace brownish red colored carapace, including a lot of lateritic nodules, this material seems to be a little bit indurated	79-1	5	-	-
	3.0-7.0m: Saprolite A yellow colored saprolite, fine grained, very soft	79-2	5	-	-
		79-3	1	-	-
	7.0-16.0m: Saprolite B greenish gray to greenish yellow colored saprolite, including chips of weathered meta-sediment	79-4	<1	-	-
		79-5	15	-	-
		79-6	29	-	-
		79-7	43	-	-
		79-8	10	-	-
		79-9	4	-	-
		79-10	5	6	-
		79-11	7	-	-
		79-12	25	-	-
		79-13	18	-	-
	79-14	22	-	-	
	16.0-18.0m: Meta-sandstone green colored schistosed rock, chloritized, meta-volcanics?	79-15	7	-	-
		79-16	5	-	-
79-17		26	-	-	
79-18		8	-	-	
79-19		15	-	-	
18.0-22.0m: Meta-sandstone black colored meta-sandstone, coarse grained, massive		79-20	7	13	-
		79-21	20	-	-
22.0-24.0m: Peritic schist dark gray colored peritic schist, with no sulfide, schistosed		79-22	41	-	-
		79-23	11	-	-
24.0-44.0m: Meta-sandstone gray colored meta-graywacke, coarse grained, massive, not schistosed		79-24	18	-	-
	79-25	75	-	-	
	79-26	60	-	-	
	79-27	24	-	-	
	79-28	23	-	-	
	79-29	9	-	-	
	79-30	8	16	-	
	79-31	17	-	-	
	79-32	21	-	-	
	79-33	20	-	-	
	79-34	29	-	-	
	79-35	9	-	-	
	79-36	10	-	-	
	79-37	6	-	-	
79-38	6	-	-		
79-39	21	-	-		
79-40	29	23	-		
79-41	28	-	-		
79-42	33	-	-		
79-43	24	-	-		
79-44	29	-	-		
	24.0-44.0m: Fe (OH) nodules? fall from upper part?				

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 13.0	0.0-4.0m: Carapace brown colored carapace, silty, clay material, including some lateritic nodules	81-1	78	-	-
		81-2	268	-	-
		81-3	69	-	-
		81-4	144	-	-
	4.0-13.0m: Saprolite B greenish yellow to greensih gray colored saprolite, including chips of weathered meta-sediment	81-5	31	-	-
		81-6	23	22	-
		81-7	29	-	-
		81-8	44	-	-
		81-9	16	-	-
		81-10	11	-	-
		81-11	19	-	-
		81-12	3	-	-
		81-13	14	-	-
15 20 25 30 35 40 45 50 52.0	13.0-52.0m: Muddy sandstone black colored meta-graywacke, fine grained, massive, pyrite dissemination is less than 1% 40.0-60.0: contents of sulfide increase (2-5%) water	81-14	21	-	-
		81-15	30	-	-
		81-16	27	30	-
		81-17	115	-	-
		81-18	69	-	-
		81-19	59	-	-
		81-20	86	-	-
		81-21	26	-	-
		81-22	31	-	-
		81-23	24	-	-
		81-24	22	-	-
		81-25	50	-	-
		81-26	31	30	-
		81-27	33	-	-
		81-28	23	-	-
		81-29	21	-	-
		81-30	34	-	-
		81-31	24	-	-
		81-32	26	-	-
		81-33	34	-	-
		81-34	34	-	-
		81-35	24	-	-
		81-36	24	22	-
		81-37	10	-	-
		81-38	10	-	-
		81-39	10	-	-
		81-40	13	-	-
81-41	12	-	-		
81-42	52	-	-		
81-43	21	-	-		
81-44	15	-	-		
81-45	18	-	-		
81-46	36	61	-		
81-47	23	-	-		
81-48	29	-	-		
81-49	13	-	-		
81-50	15	-	-		
81-51	11	-	-		
81-52	13	-	-		
55 60	52.0-60.0m: Peritic schist dark gray to black colored peritic schist, with very fine grained pyrite dissemination, contents of pyrite is 3%	81-53	14	-	-
		81-54	51	-	-
		81-55	14	-	-
		81-56	257	19	-
		81-57	16	-	-
		81-58	31	-	-
		81-59	17	-	-
		81-60	14	-	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 9.0	0.0-9.0m: Crust, carapace reddish brown colored crust and carapace, including a lot of lateritic nodules	82-1	23	113	44
		82-2	8	32	-
		82-3	1	26	-
		82-4	10	85	54
		82-5	92	130	-
		82-6	146	179	-
		82-7	90	125	-
		82-8	0	163	114
		82-9	72	120	-
10 12.0	9.0-12.0m: Mottled clay brown colored clay, with some lateritic nodules	82-10	121	210	-
		82-11	140	188	-
		82-12	162	170	-
15 20 25 30 35 39.0	12.0-39.0m: Saprolite A pinkish brown to pinkish yellow colored, fine grained saprolite, very soft, with no rock chips	82-13	194	179	-
		82-14	182	215	-
		82-15	253	291	-
		82-16	311	290	-
		82-17	494	571	-
		82-18	466	552	-
		82-19	770	820	-
		82-20	698	787	-
		82-21	173	-	-
		82-22	127	-	-
		82-23	34	-	-
		82-24	179	-	-
		82-25	324	-	-
		82-26	726	-	-
		82-27	120	-	-
		82-28	479	-	-
		82-29	593	-	-
		82-30	352	389	-
82-31	74	-	-		
82-32	123	-	-		
82-33	121	-	-		
82-34	173	-	-		
82-35	133	-	-		
82-36	63	-	-		
82-37	170	-	-		
82-38	83	-	-		
82-39	178	-	-		
40 45 50 55 60	52.0-60.0m: Granodiorite gray colored, hornblende-biotite granodiorite, medium grained, fresh, with pyrite dissemination, contents of sulfide is 0-1% in volume	82-40	41	111	-
		82-41	124	-	-
		82-42	142	-	-
		82-43	242	-	-
		82-44	249	-	-
		82-45	3	-	-
		82-46	146	-	-
		82-47	47	-	-
		82-48	71	-	-
		82-49	559	-	-
		82-50	77	54	-
		82-51	51	-	-
82-52	137	-	-		
82-53	54	-	-		
82-54	54	-	-		
82-55	79	-	-		
82-56	260	-	-		
82-57	84	-	-		
82-58	101	-	-		
82-59	80	-	-		
82-60	29	30	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-9.0m: Crust, carapace reddish brown colored crust and carapace, including a lot of lateritic nodules	83-1	29	-	-
		83-2	20	-	-
		83-3	187	-	-
		83-4	25	-	-
		83-5	33	-	-
		83-6	424	-	-
		83-7	117	-	-
		83-8	134	-	-
		83-9	116	-	-
9.0	9.0-12.0m: Mottled clay brown colored clay, including iron oxide, with some lateritic nodules	83-10	129	113	-
83-11		112	-	-	
83-12		167	-	-	
12.0	12.0-39.0m: Saprolite A brown to pinkish brown colored fine grained saprolite, very soft, with no rock chips	83-13	206	-	-
83-14		118	-	-	
83-15		31	-	-	
83-16		23	-	-	
83-17		79	-	-	
83-18		44	-	-	
83-19		176	-	-	
83-20		341	46	-	
83-21		39	-	-	
83-22		65	-	-	
83-23		20	-	-	
83-24		324	-	-	
83-25		265	-	-	
83-26		57	-	-	
83-27		310	-	-	
83-28		141	-	-	
83-29		214	-	-	
83-30		68	62	-	
83-31		2,253	20	0	
83-32	45	-	-		
83-33	33	-	-		
83-34	19	-	-		
83-35	33	-	-		
83-36	69	-	-		
83-37	39	-	-		
83-38	52	-	-		
83-39	468	-	-		
39.0	52.0-60.0m: Saprolite B greensih gray colored saprolite, including chips of weathered granodiorite, with minor chlorite	83-40	111	56	-
83-41		122	-	-	
83-42		39	-	-	
83-43		25	-	-	
83-44		28	-	-	
83-45		10	-	-	
83-46		19	-	-	
83-47		229	-	-	
83-48		31	-	-	
83-49		22	-	-	
83-50		51	34	-	
83-51		46	-	-	
83-52	59	-	-		
83-53	63	-	-		
83-54	52	-	-		
83-55	41	-	-		
83-56	65	-	-		
56.0	56.0-60.0m: Silicified zone light gray colored silicified rock, with strongly disseminated of pyrite>arsenopyrite >pyrrhotite	83-57	91	-	-
83-58		176	-	-	
83-59		307	-	-	
83-60		188	162	-	
60.0					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 5.0	0.0-9.0m: Alluvial sediments brown colored sediments, including lateritic nodules 0.0-2.0m: dark brown colored sediments, with organic material	41-1	162	-	-
		41-2	91	-	-
		41-3	63	-	-
		41-4	29	-	-
		41-5	74	-	-
10 14.0	5.0-14.0m: Mottled clay brown colored, fine grained clay, soft, transition to saprolite is gradual	41-6	43	-	-
		41-7	137	-	-
		41-8	86	-	-
		41-9	22	-	-
		41-10	20	181	-
		41-11	48	-	-
		41-12	20	-	-
		41-13	29	-	-
		41-14	298	-	-
		41-15	194	-	-
15 20 25 30 35	14.0-35.0m: Saprolite A yellowish gray to greenish gray colored, fine grained saprolite, soft, including no rock chips	41-16	55	-	-
		41-17	36	-	-
		41-18	57	-	-
		41-19	13	-	-
		41-20	85	38	-
		41-21	26	-	-
		41-22	30	-	-
		41-23	7	-	-
		41-24	0	-	-
		41-25	8	-	-
		41-26	0	-	-
		41-27	0	-	-
		41-28	19	-	-
		41-29	4	-	-
		41-30	73	26	-
35 40 40.0	35.0-40.0m: Saprolite B greensih gray colored saprolite, including some chips of deeply weathered rock	41-31	0	-	-
		41-32	706	-	-
		41-33	0	-	-
		41-34	6	-	-
		41-35	14	-	-
		41-36	0	-	-
		41-37	60	-	-
		41-38	23	-	-
		41-39	27	-	-
		41-40	0	555	-
45					
50					
55					
60					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
0.0-2.0m 2.0-20.0m 5 10 15 20	0.0-2.0m: Carapace brown colored, with iron nodules	25-1	233	-	-
		25-2	82	-	-
		25-3	72	-	-
	2.0-20.0m: Saprolite B greenish grey colored saprolite, including a lot of chips of weathered rock	25-4	24	-	-
		25-5	14	-	-
		25-6	29	-	-
		25-7	38	-	-
		25-8	28	-	-
		25-9	34	-	-
		25-10	18	12	-
		25-11	14	-	-
		25-12	30	-	-
		25-13	33	-	-
		25-14	21	-	-
		25-15	38	-	-
		25-16	107	-	-
		25-17	30	-	-
		25-18	21	-	-
		25-19	12	-	-
		16.0-60.0m 25 30 35 40 45 50 55 60	16.0-60.0m: Meta-sandstone (greywacke) greenish dark grey colored, meta-greywacke with disseminated pyrite, total amount of pyrite = 0 % to 2 % epidotic alteration is observed, meta-volcanics	25-20	44
25-21	12			-	-
25-22	9			-	-
25-23	12			-	-
25-24	22			-	-
25-25	16			-	-
25-26	25			-	-
25-27	7			-	-
25-28	2			-	-
25-29	3			-	-
25-30	39			18	-
25-31	9			-	-
25-32	47			-	-
25-33	15			-	-
25-34	16			-	-
25-35	8			-	-
25-36	30			-	-
25-37	4			-	-
25-38	1			-	-
25-39	35			-	-
25-40	107			22	-
25-41	54			-	-
25-42	91			-	-
25-43	33			-	-
25-44	59			-	-
25-45	19			-	-
25-46	43			-	-
25-47	34			-	-
25-48	45			-	-
25-49	92			-	-
25-50	52			93	-
25-51	72			-	-
25-52	91			-	-
25-53	41			-	-
25-54	49			-	-
25-55	53			-	-
25-56	54			-	-
25-57	64			-	-
25-58	32			-	-
25-59	40			-	-
25-60	33	48	-		

Column (m)	Description	Sample No.	Au			
			ppb	ppb	ppb	
1.0 5 10 15 20 22.0	0.0-1.0m: Carapace brown colored carapace with laterite nodules	26-1	50	-	-	
		26-2	109	-	-	
		26-3	32	-	-	
	1.0-22.0m: Saprolite B greenish grey colored saprolite including chips weathered rock	26-4	688	-	-	
		26-5	56	-	-	
		26-6	45	-	-	
		26-7	18	-	-	
		26-8	19	-	-	
		26-9	47	-	-	
		26-10	54	59	-	
		26-11	30	-	-	
		26-12	11	-	-	
		26-13	41	-	-	
	22.0 25 30 35 40 45 50 55 60 60.0	22.0-60.0m: Meta-sandstone dark grey to greenish grey colored meta- sandstone, with disseminated pyrite, total amount of pyrite = 0 % to 1 % green color shows chloritic and / or epidotic alteration, meta-volcanics ? with traces of calcite films	26-14	39	-	-
			26-15	44	-	-
			26-16	66	-	-
			26-17	26	-	-
			26-18	64	-	-
			26-19	62	-	-
			26-20	218	235	-
			26-21	17	-	-
			26-22	142	-	-
26-23			134	-	-	
26-24	12	-	-			
26-25	8	-	-			
26-26	21	-	-			
26-27	31	-	-			
26-28	84	-	-			
26-29	13	-	-			
26-30	87	81	-			
26-31	31	-	-			
26-32	14	-	-			
26-33	7	-	-			
26-34	33	-	-			
26-35	38	-	-			
26-36	155	-	-			
26-37	123	-	-			
26-38	831	-	-			
26-39	107	-	-			
26-40	48	22	-			
26-41	1,189	102	686			
26-42	68	-	-			
26-43	145	-	-			
26-44	3,734	3,780	3,463			
26-45	23	-	-			
26-46	1,776	142	34			
26-47	171	-	-			
26-48	67	-	-			
26-49	1,514	1,110	1,234			
26-50	276	235	-			
26-51	834	-	-			
26-52	1,022	1,150	789			
26-53	277	-	-			
26-54	59	-	-			
26-55	495	-	-			
26-56	78	-	-			
26-57	47	29	22			
26-58	9,670	100	150			
26-59	308	400	410			
26-60	115	108	-			

KRC-27

Coordination: N1250 W400

Elevation: 345m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0	0.0-1.0m: Carapace brown colored soil with laterite nodules	27-1	49	-	-
3.0		27-2	364	-	-
5	2.0-3.0m: Mottled clay brown to greenish grey colored clay	27-3	362	-	-
		27-4	81	-	-
		27-5	21	-	-
		27-6	34	-	-
		27-7	22	-	-
		27-8	2	-	-
		27-9	28	-	-
		27-10	131	132	-
		27-11	24	-	-
		27-12	49	-	-
		27-13	179	-	-
		27-14	58	-	-
		27-15	22	-	-
		27-16	136	-	-
		27-17	56	-	-
27-18	56	-	-		
27-19	120	-	-		
27-20	1,403	1,718	-		
27-21	244	-	-		
27-22	146	-	-		
27-23	30	-	-		
27-24	30	-	-		
27-25	115	-	-		
27-26	42	-	-		
27-27	58	-	-		
27-28	508	-	-		
27-29	93	-	-		
27-30	131	206	-		
27-31	44	-	-		
27-32	102	-	-		
27-33	931	-	-		
27-34	92	-	-		
27-35	153	-	-		
27-36	482	-	-		
27-37	39	-	-		
27-38	220	-	-		
27-39	553	-	-		
27-40	77	40	-		
27-41	58	-	-		
27-42	418	-	-		
27-43	154	-	-		
27-44	36	-	-		
27-45	0	-	-		
27-46	12	-	-		
27-47	7	-	-		
27-48	492	-	-		
27-49	289	-	-		
27-50	1,151	158	124		
27-51	237	-	-		
27-52	2,835	2,668	2,263		
27-53	91	-	-		
27-54	18	-	-		
27-55	67	-	-		
27-56	20	-	-		
27-57	12	-	-		
27-58	8	-	-		
27-59	2	-	-		
27-60	0	2	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Surface soil grey colored soil, including quartz fragments	28-1	49	-	-
		28-2	10	-	-
5	2.0-10.0m: Saprolite A brown to greenish grey colored, fine grained powder	28-3	81	55	-
		28-4	16	-	-
		28-5	107	-	-
		28-6	27	-	-
		28-7	7	-	-
		28-8	9	-	-
		28-9	8	-	-
		28-10	10	-	-
10	10.0-25.0m: Saprolite B greenish grey colored saprolite, including chips of weathered rock	28-11	11	-	-
		28-12	8	-	-
		28-13	8	5	-
		28-14	8	-	-
		28-15	8	-	-
		28-16	22	-	-
		28-17	8	-	-
		28-18	164	-	-
		28-19	28	-	-
		28-20	15	-	-
15		28-21	111	-	-
		28-22	18	-	-
		28-23	22	10	-
		28-24	54	-	-
		28-25	6	-	-
		28-26	19	-	-
		28-27	1	-	-
		28-28	11	-	-
		28-29	43	-	-
		28-30	84	-	-
20		28-31	9	-	-
		28-32	6	-	-
		28-33	27	32	-
		28-34	56	-	-
		28-35	105	-	-
		28-36	43	-	-
		28-37	5	-	-
		28-38	5	-	-
		28-39	6	-	-
		28-40	168	-	-
25		28-41	261	-	-
		28-42	88	-	-
		28-43	0	7	-
		28-44	59	-	-
		28-45	116	-	-
		28-46	14	-	-
		28-47	6	-	-
		28-48	586	-	-
		28-49	14	-	-
		28-50	6	-	-
30		28-51	8	-	-
		28-52	3	-	-
		28-53	15	9	-
		28-54	19	-	-
		28-55	6	-	-
		28-56	10	-	-
		28-57	8	-	-
		28-58	1	-	-
		28-59	7	-	-
		28-60	1,391	1,460	0

Column (m)	Description	Sample No.	Au			
			ppb	ppb	ppb	
5	0.0-2.0m: Surface soil grey to brown colored soil, including silty material	30-1	84	-	-	
		30-2	82	-	-	
	2.0-4.0m: Carapace brown colored, Fe-oxide rich zone	30-3	55	34	-	
		30-4	37	-	-	
	4.0-7.0m: Carapace brown colored, fine grained soil	30-5	30	-	-	
		30-6	35	-	-	
	7.0	7.0-10.0m: Mottled clay brown to greenish brown, some times reddish brown colored clay	30-7	162	-	-
			30-8	23	-	-
	10	10.0-22.0m: Saprolite A greenish grey to light grey colored fine grained powder	30-9	41	-	-
			30-10	43	-	-
15	22.0-37.0m: Saprolite B greenish grey to light grey colored fine grained powder, including chips of weathered granodiorite	30-11	65	-	-	
		30-12	118	-	-	
20	37.0-60.0m: Fine grained granodiorite grey to light grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1 % including pyrrhotite ?	30-13	383	281	-	
		30-14	173	-	-	
		30-15	2,103	2,173	2,023	
		30-16	71	-	-	
		30-17	101	-	-	
		30-18	30	-	-	
		30-19	84	-	-	
		30-20	196	-	-	
		30-21	75	-	-	
		30-22	36	-	-	
25		30-23	96	120	-	
		30-24	66	-	-	
30		30-25	77	-	-	
		30-26	587	-	-	
35		30-27	174	-	-	
		30-28	55	-	-	
40		30-29	287	-	-	
		30-30	74	-	-	
45		30-31	27	-	-	
		30-32	62	-	-	
50		30-33	117	174	-	
		30-34	52	-	-	
55		30-35	60	-	-	
		30-36	31	-	-	
60		30-37	43	-	-	
		30-38	15	-	-	
		30-39	25	-	-	
		30-40	14	-	-	
		30-41	7	-	-	
		30-42	101	-	-	
		30-43	53	58	-	
		30-44	108	-	-	
		30-45	22	-	-	
		30-46	91	-	-	
		30-47	50	-	-	
		30-48	56	-	-	
		30-49	63	-	-	
		30-50	22	-	-	
		30-51	16	-	-	
		30-52	45	-	-	
		30-53	327	349	-	
		30-54	15	-	-	
		30-55	28	-	-	
		30-56	18	-	-	
		30-57	16	-	-	
		30-58	45	-	-	
		30-59	81	-	-	
		30-60	29	-	-	

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Surface soil grey to brown colored soil, including silty material	31-1	137	-	-
		31-2	124	-	-
4.0	2.0-4.0m: Carapace brown colored, Fe-oxide rich zone	31-3	373	385	-
		31-4	112	-	-
7.0	4.0-7.0m: Carapace brown colored, fine grained soil	31-5	29	-	-
		31-6	27	-	-
10.0	7.0-10.0m: Mottled clay brown to greenish brown, some times reddish brown colored clay	31-7	23	-	-
		31-8	27	-	-
15.0	10.0-22.0m: Saprolite A greenish grey to light grey colored fine grained powder	31-9	31	-	-
		31-10	20	-	-
20.0	22.0-37.0m: Saprolite B greenish grey to light grey colored fine grained powder, including chips of weathered granodiorite	31-11	53	-	-
		31-12	15	-	-
25.0		31-13	25	25	-
		31-14	7	-	-
30.0		31-15	20	-	-
		31-16	1,905	1,035	1,186
35.0		31-17	2,324	947	2,369
		31-18	48	-	-
40.0		31-19	713	-	-
		31-20	377	-	-
45.0		31-21	54	-	-
		31-22	100	-	-
50.0		31-23	46	40	-
		31-24	33	-	-
55.0		31-25	309	-	-
		31-26	64	-	-
60.0		31-27	23	-	-
		31-28	481	-	-
		31-29	60	-	-
		31-30	41	-	-
		31-31	15	-	-
		31-32	312	-	-
		31-33	85	50	-
		31-34	88	-	-
		31-35	19	-	-
		31-36	4	-	-
		31-37	7	-	-
		31-38	37	-	-
		31-39	7	-	-
		31-40	260	-	-
		31-41	175	-	-
		31-42	118	-	-
		31-43	9	927	24
		31-44	748	-	-
		31-45	116	-	-
		31-46	86	-	-
		31-47	41	-	-
		31-48	89	-	-
		31-49	134	-	-
		31-50	33	-	-
		31-51	188	-	-
		31-52	49	-	-
		31-53	124	125	-
		31-54	77	-	-
		31-55	59	-	-
		31-56	53	-	-
		31-57	118	-	-
		31-58	94	-	-
		31-59	25	-	-
		31-60	34	-	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Carapace reddish brown colored soil, including iron nodules	32-1	18	-	-
		32-2	31	-	-
5	2.0-8.0m: Mottled clay brown to yellowish brown colored clay, including limonite ?	32-3	27	33	-
		32-4	25	-	-
		32-5	12	-	-
		32-6	10	-	-
		32-7	13	-	-
		32-8	19	-	-
		32-9	11	-	-
10	8.0-22.0m: Saprolite A greenish grey colored fine grained saprolite	32-10	430	-	-
		32-11	693	-	-
		32-12	31	-	-
		32-13	21	27	-
		32-14	29	-	-
		32-15	15	-	-
		32-16	22	-	-
		32-17	11	-	-
		32-18	14	-	-
		32-19	15	-	-
		32-20	9	-	-
		32-21	10	-	-
		32-22	26	-	-
15	22.0-31.0m: Saprolite B greenish grey colored, fine grained saprolite including chips of weathered granodiorite	32-23	41	10	-
		32-24	19	-	-
		32-25	15	-	-
		32-26	110	-	-
		32-27	47	-	-
		32-28	8	-	-
		32-29	4	-	-
		32-30	4	-	-
		32-31	11	-	-
		32-32	4	-	-
20	31.0-39.0m: Fine grained granodiorite chloritized greenish grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1%	32-33	66	31	-
		32-34	11	-	-
		32-35	7	-	-
		32-36	42	-	-
		32-37	24	-	-
		32-38	21	-	-
		32-39	13	-	-
		32-40	23	-	-
		32-41	50	-	-
		32-42	25	-	-
25	39.0-42.0m: Fault ? small amount of rock chips of sericitized granodiorite	32-43	40	39	-
		32-44	9	-	-
		32-45	91	-	-
		32-46	107	-	-
		32-47	59	-	-
		32-48	14	-	-
		32-49	7	-	-
		32-50	5	-	-
		32-51	5	-	-
		32-52	5	-	-
30	42.0-60.0m: Fine grained granodiorite chloritized greenish grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1%	32-53	5	4	-
		32-54	3	-	-
		32-55	23	-	-
		32-56	28	-	-
		32-57	14	-	-
		32-58	7	-	-
		32-59	3	-	-
		32-60	3	-	-
		32-60	3	-	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 13.0	0.0-13.0m: Carapace reddish brown colored carapace, with a lot of iron nodules (diameter: 3-5mm), coarse grained sand	76-1	4	-	-
		76-2	6	-	-
		76-3	4	-	-
		76-4	18	-	-
		76-5	42	-	-
		76-6	32	-	-
		76-7	48	-	-
		76-8	74	-	-
		76-9	66	-	-
		76-10	68	90	-
		76-11	84	-	-
		76-12	124	-	-
		76-13	168	-	-
15 20 25 30 34.0	13.0-34.0m: Saprolite A reddish brown to yellowish brown colored, very fine grained saprolite, without rock chips	76-14	876	1,054	-
		76-15	1,840	1,440	1,782
		76-16	402	-	-
		76-17	454	-	-
		76-18	64	-	-
		76-19	254	-	-
		76-20	78	57	-
		76-21	111	-	-
		76-22	97	-	-
		76-23	15	-	-
		76-24	57	-	-
		76-25	44	-	-
		76-26	18	-	-
		76-27	32	-	-
		76-28	31	-	-
		76-29	19	-	-
		76-30	26	21	-
76-31	50	-	-		
76-32	29	-	-		
76-33	11	-	-		
76-34	14	-	-		
35 40 45 49.0	34.0-49.0m: Saprolite A light gray to greenish light gray colored, saprolite, medium to fine grained, without rock chips	76-35	11	-	-
		76-36	12	-	-
		76-37	13	-	-
		76-38	13	-	-
		76-39	6	-	-
		76-40	5	13	-
		76-41	13	-	-
		76-42	5	-	-
		76-43	2	-	-
		76-44	0	-	-
		76-45	38	-	-
		76-46	35	-	-
		76-47	77	-	-
		76-48	237	-	-
		76-49	59	-	-
		76-50	14	125	-
		76-51	9	-	-
76-52	5	-	-		
76-53	46	-	-		
55 60.0	52.0-60.0m: Volcanic sandstone black colored volcanic sandstone, weakly schistosed, including a lot of plagioclase grains (subhedral, 1-2mm, rounded), Fe-oxides stain along fractures	76-54	165	-	-
		76-55	42	-	-
		76-56	21	-	-
		76-57	243	-	-
		76-58	19	-	-
		76-59	407	-	-
		76-60	11	7	-

KRC-77

Coordination: N1250 E350

Elevation: 370m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au				
			ppb	ppb	ppb		
5 10 15 16.0	0.0-16.0m: Carapace reddish brown colored carapace, coarse grained, including a lot of iron nodules (diameter: 0.3-0.8cm)	77-1	20	-	-		
		77-2	352	-	-		
		77-3	56	-	-		
		77-4	36	-	-		
		77-5	42	-	-		
		77-6	36	-	-		
		77-7	20	-	-		
		77-8	34	-	-		
		77-9	48	-	-		
		77-10	28	21	-		
		77-11	58	-	-		
		77-12	76	-	-		
		77-13	52	-	-		
		77-14	56	-	-		
		77-15	84	-	-		
		16.0 20 25 30 35 38.0	16.0-38.0m: Saprolite A reddish brown to brown colored saprolite, very fine grained saprolite (silty), without rock chips	77-16	76	-	-
77-17	44			-	-		
77-18	54			-	-		
77-19	32			-	-		
77-20	144			143	-		
77-21	25			-	-		
77-22	36			-	-		
77-23	34			-	-		
77-24	35			-	-		
77-25	26			-	-		
77-26	19			-	-		
77-27	31			-	-		
77-28	21			-	-		
77-29	21			-	-		
77-30	26			25	-		
77-31	20			-	-		
77-32	20			-	-		
77-33	12			-	-		
77-34	15			-	-		
77-35	20			-	-		
77-36	17			-	-		
77-37	10			-	-		
38.0 40 45 50 55 60	38.0-60.0m: Saprolite A brown to yellowish light gray colored saprolite, fine grained, without rock chips			77-38	15	-	-
				77-39	13	-	-
		77-40	16	17	-		
		77-41	7	-	-		
		77-42	18	-	-		
		77-43	23	-	-		
		77-44	18	-	-		
		77-45	20	-	-		
		77-46	32	-	-		
		77-47	13	-	-		
		77-48	15	-	-		
		77-49	16	-	-		
		77-50	16	26	-		
77-51	16	-	-				
77-52	13	-	-				
77-53	22	-	-				
77-54	10	-	-				
77-55	13	-	-				
77-56	13	-	-				
77-57	10	-	-				
77-58	20	-	-				
77-59	13	-	-				
77-60	5	10	-				

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-5.0m: Carapace reddish brown colored carapace, with lateritic nodules	67-1	173	-	-
		67-2	29	-	-
		67-3	55	-	-
		67-4	30	-	-
		67-5	14	-	-
10	5.0-18.0m: Saprolite A yellowish brown colored saprolite, fine grained	67-6	6	-	-
		67-7	0	-	-
		67-8	2,970	49	0
		67-9	32	-	-
		67-10	0	21	-
		67-11	62	-	-
		67-12	33	-	-
		67-13	29	-	-
		67-14	43	-	-
		67-15	0	-	-
		67-16	14	-	-
		67-17	35	-	-
		15	18.0-37.0m: Saprolite A yellowish gray to brownish gray colored saprolite, fine grained, including no rock chips	67-18	21
67-19	29			-	-
67-20	28			0	-
67-21	166			-	-
67-22	52			-	-
67-23	32			-	-
67-24	32			-	-
67-25	52			-	-
67-26	22			-	-
67-27	130			-	-
67-28	20			-	-
67-29	22			-	-
67-30	28			28	-
67-31	18			-	-
67-32	20			-	-
67-33	32			-	-
20	37.0-47.0m: Saprolite A gray to yellowish gray colored saprolite, fine grained, including no rock chips			67-34	26
		67-35	10	-	-
		67-36	14	-	-
		67-37	20	-	-
		67-38	24	-	-
		67-39	8	-	-
		67-40	20	14	-
		67-41	30	-	-
		67-42	23	-	-
		67-43	0	-	-
25	47.0-51.0m: Saprolite B gray colored saprolite, including small amount of rock fragments	67-44	7	-	-
		67-45	130	-	-
		67-46	19	-	-
		67-47	0	-	-
		67-48	20	-	-
		67-49	37	-	-
		67-50	33	26	-
30	51.0-60.0m: Peritic schist dark gray colored peritic schist, with schistosity filling iron oxide and pyrite, fracture rich, including rock fragment (diameter: 2-5mm)	67-51	0	-	-
		67-52	36	-	-
		67-53	15	-	-
		67-54	10	-	-
		67-55	20	-	-
		67-56	16	-	-
		67-57	52	-	-
		67-58	13	-	-
		67-59	22	-	-
		67-60	9	38	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-9.0m: Carapace reddish brown colored carapace, including lateritic gravels, with iron oxides	68-1	8	-	-
		68-2	7	-	-
		68-3	11	-	-
		68-4	9	-	-
		68-5	16	-	-
		68-6	174	-	-
		68-7	10	-	-
		68-8	9	-	-
		68-9	24	-	-
9.0	9.0-29.0m: Saprolite A greenish gray colored saprolite, fine grained and very soft saprolite	68-10	9	11	-
68-11		8	-	-	
68-12		7	-	-	
68-13		13	-	-	
68-14		18	-	-	
68-15		5	-	-	
68-16		5	-	-	
68-17		4	-	-	
68-18		6	-	-	
68-19		8	-	-	
68-20		17	17	-	
68-21		26	-	-	
68-22		11	-	-	
68-23		6	-	-	
68-24		19	-	-	
68-25		32	-	-	
68-26		63	-	-	
68-27		25	-	-	
68-28		0	-	-	
68-29	27	-	-		
29.0	29.0-39.0m: Saprolite B greenish gray colored saprolite, including chips of deeply weathered rock, with many quartz vein chips	68-30	56	189	85
water		68-31	66	-	-
68-32		53	-	-	
68-33		40	-	-	
68-34		36	-	-	
68-35		19	-	-	
68-36		21	-	-	
68-37		37	-	-	
68-38		20	-	-	
68-39		10	-	-	
39.0		39.0-60.0m: Peritic schist gray to dark gray colored peritic schist, fine grained and schistosed, rock chips clearly show schistosity, sulfides are mainly included in schistosity and fracture planes	68-40	13	11
68-41	3		-	-	
68-42	3		-	-	
68-43	3		-	-	
68-44	15		-	-	
68-45	5		-	-	
68-46	6		-	-	
68-47	5		-	-	
68-48	4		-	-	
68-49	5		-	-	
68-50	7		1	-	
68-51	2	-	-		
68-52	5	-	-		
68-53	5	-	-		
68-54	3	-	-		
68-55	miss	miss	miss		
68-56	2	-	-		
68-57	5	-	-		
68-58	45	-	-		
68-59	0	-	-		
68-60	2	-	-		

Column (m)	Description	Sample No.	Au				
			ppb	ppb	ppb		
3.0	0.0-3.0m: Surface soil dark brown colored soil, including lateritic gravels	70-1	87	-	-		
		70-2	336	-	-		
		70-3	69	-	-		
5.0	3.0-5.0m: Carapace red brown colored carapace, including lateritic gravels	70-4	58	-	-		
		70-5	40	-	-		
		70-6	57	-	-		
12.0	5.0-12.0m: Mottled clay brown to yellowish brown colored mottled clay, soft	70-7	4,262	1,798	3,531		
		70-8	1,495	1,156	1,646		
		70-9	1,732	1,374	1,646		
		70-10	849	423	560		
		70-11	217	-	-		
		70-12	69	-	-		
		16.0	12.0-16.0m: Saprolite A greenish gray colored saprolite, fine grained, soft	70-13	161	-	-
				70-14	108	-	-
				70-15	59	-	-
		21.0	16.0-21.0m: Saprolite B greenish gray to gray colored saprolite, including chips of weathered meta-sandstone	70-16	44	-	-
70-17	81			-	-		
70-18	269			-	-		
70-19	84			-	-		
70-20	41			42	-		
70-21	41			-	-		
21.0-60.0m	21.0-60.0m: Meta-sandstone gray to dark gray colored meta-sandstone, fine grained and hard, with pyrite-arsenopyrite dissemination (30-31, 40-41m), including quartz vein chips, total amount of sulfide=0-2%, and also sulfide in small veinlets	70-22	54	-	-		
		70-23	30	-	-		
		70-24	97	-	-		
		70-25	390	-	-		
		70-26	282	-	-		
		70-27	104	-	-		
		70-28	181	-	-		
		70-29	102	-	-		
		70-30	193	190	-		
		70-31	765	-	-		
		70-32	1,214	1,151	1,234		
		70-33	1,217	1,328	1,749		
		70-34	5,865	7,050	6,271		
		70-35	4,245	5,425	5,486		
		70-36	2,835	2,900	2,983		
		70-37	4,513	5,694	5,040		
		70-38	4,865	5,244	5,863		
		70-39	1,323	-	1,200		
		70-40	129	123	-		
70-41	94	-	-				
70-42	116	-	-				
70-43	86	-	-				
70-44	165	-	-				
70-45	329	-	-				
70-46	146	-	-				
70-47	60	-	-				
70-48	143	-	-				
70-49	34	-	-				
70-50	34	32	-				
70-51	44	-	-				
70-52	80	-	-				
70-53	76	-	-				
70-54	40	-	-				
70-55	71	-	-				
70-56	44	-	-				
70-57	29	-	-				
70-58	20	-	-				
70-59	10	-	-				
70-60	5	7	-				

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Carapace brown colored soil, including iron nodules	18-1	218	-	-
		18-2	169	-	-
5	2.0-17.0m: Saprolite B yellowish grey colored, coarse grained powder including chips of weathered granodiorite	18-3	73	-	-
		18-4	76	-	-
		18-5	47	-	-
		18-6	36	-	-
		18-7	20	-	-
		18-8	29	-	-
		18-9	329	-	-
		18-10	38	47	-
		18-11	14	-	-
		18-12	46	-	-
15		18-13	18	-	-
		18-14	12	-	-
		18-15	29	-	-
		18-16	42	-	-
		18-17	19	-	-
		18-18	14	-	-
		18-19	7	-	-
		18-20	7	6	-
		18-21	8	-	-
		18-22	12	-	-
25		18-23	25	-	-
		18-24	12	-	-
		18-25	5	-	-
		18-26	6	-	-
		18-27	5	-	-
		18-28	5	-	-
		18-29	16	-	-
		18-30	21	34	-
		18-31	7	-	-
		18-32	16	-	-
35		18-33	12	-	-
		18-34	10	-	-
		18-35	54	-	-
		18-36	78	-	-
		18-37	25	-	-
		18-38	15	-	-
		18-39	23	-	-
		18-40	41	57	-
		18-41	58	-	-
		18-42	1	-	-
45	40.0-46.0m: porphyry or porphyritic dacite dark grey colored porphyry, with plagioclase phenocrysts in a dark grey matrix	18-43	0	-	-
		18-44	3	-	-
		18-45	0	-	-
		18-46	49	-	-
46.0	46.0-48.0m: Fault ?	18-47	19	-	-
		18-48	104	-	-
48.0	dark grey colored sand	18-49	90	-	-
		18-50	17	14	-
55	40.0-46.0m: porphyry or porphyritic dacite dark grey colored porphyry, with plagioclase phenocrysts in a dark grey matrix, rock chip of 18-56 shows sedimentary texture with strong dissemination of pyrite and arsenopyrite	18-51	28	-	-
		18-52	8	-	-
		18-53	23	-	-
		18-54	16	-	-
		18-55	6	-	-
		18-56	68	-	-
		18-57	58	-	-
		18-58	37	-	-
		18-59	51	-	-
		18-60	45	44	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0 5 10 15 20	0.0-1.0m: Carapace reddish brown colored soil with nodules	19-1	503	-	-
		19-2	88	-	-
	1.0-20.0m: Saprolite B yellowish grey to greenish grey colored, coarse grained sand coming from weathered granodiorite	19-3	48	-	-
		19-4	15	-	-
		19-5	9	-	-
		19-6	19	-	-
		19-7	50	-	-
		19-8	18	-	-
		19-9	65	-	-
		19-10	35	24	-
		19-11	11	-	-
		19-12	17	-	-
		19-13	6	-	-
		19-14	11	-	-
		19-15	9	-	-
		19-16	35	-	-
		19-17	22	-	-
		19-18	45	-	-
		19-19	45	-	-
		20.0	17.0-40.0m: Fine grained granodiorite light grey colored, fine to medium grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1%	19-20	56
25 30 35 40	19-21	57		-	-
	19-22	9		-	-
	19-23	12		-	-
	19-24	17		-	-
	19-25	34		-	-
	19-26	104		-	-
	19-27	38		-	-
	19-28	42		-	-
	19-29	26		-	-
	19-30	19		22	-
	19-31	31		-	-
	19-32	27		-	-
	19-33	16		-	-
	19-34	13		-	-
	19-35	22		-	-
	19-36	18		-	-
	19-37	15		-	-
	19-38	17		-	-
	19-39	38		-	-
43.0	43.0-53.0m: meta-sandstone dark grey colored, fine grained meta -greywacke with pyrrhotite and pyrite dissemination, with small amount of epidote	19-40	23	19	-
45 50		19-41	16	-	-
		19-42	16	-	-
		19-43	40	-	-
		19-44	37	-	-
		19-45	18	-	-
		19-46	29	-	-
		19-47	9	-	-
		19-48	61	-	-
		19-49	17	-	-
19-50	5	11	-		
53.0	53.0-60.0m: Fault ? dark grey colored sand with no rock chips	19-51	19	-	-
55 60		19-52	63	-	-
		19-53	988	1,097	1,063
		19-54	1,051	1,131	1,166
		19-55	37	-	-
		19-56	36	-	-
		19-57	9	-	-
		19-58	28	-	-
		19-59	7	-	-
		19-60	9	20	-

KRC-20

Coordination: N1500 W300

Elevation: 352m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Carapace brown colored soil with nodules	20-1	176	-	-
		20-2	164	-	-
		20-3	222	-	-
5	2.0-9.0m: Saprolite A greenish yellow to greenish grey colored, fine grained powder	20-4	21	-	-
		20-5	22	-	-
		20-6	24	-	-
		20-7	13	-	-
		20-8	10	-	-
		20-9	12	-	-
9.0	9.0-28.0m: Saprolite B pinkish grey to greenish grey colored saprolite including small amount of weathered rock chips	20-10	6	10	-
		20-11	8	-	-
		20-12	29	-	-
		20-13	16	-	-
		20-14	12	-	-
		20-15	98	-	-
		20-16	73	-	-
		20-17	536	-	-
		20-18	86	-	-
		20-19	58	-	-
		20-20	392	428	-
23.0	23.0-24.0m: hard rock of granodiorite	20-21	22	-	-
		20-22	38	-	-
24.0	23.0-24.0m: hard rock of granodiorite	20-23	41	-	-
		20-24	15	-	-
		20-25	35	-	-
		20-26	56	-	-
		20-27	85	-	-
		20-28	116	-	-
		20-29	41	-	-
		20-30	28	36	-
		20-31	63	-	-
		20-32	44	-	-
		20-33	102	-	-
		20-34	130	-	-
		20-35	146	-	-
		20-36	38	-	-
		20-37	64	-	-
		20-38	130	-	-
		28.0	28.0-60.0m: granodiorite light grey colored, fine to medium grained granodiorite (or quartz diorite), not altered total amount of pyrite = 0 % to 1%	20-39	106
20-40	170			194	-
20-41	101			-	-
20-42	263			-	-
20-43	222			-	-
20-44	62			-	-
20-45	59			-	-
20-46	12			-	-
20-47	92			-	-
20-48	54			-	-
20-49	43			-	-
20-50	48			46	-
20-51	30			-	-
55	28.0-60.0m: granodiorite	20-52	34	-	-
		20-53	44	-	-
		20-54	57	-	-
		20-55	120	-	-
		20-56	62	-	-
		20-57	39	-	-
		20-58	41	-	-
		20-59	33	-	-
		20-60	60	44	-
		60.0			

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-3.0m: Carapace brown to brownish grey colored soil with iron nodules	21-1	129	-	-
		21-2	64	-	-
		21-3	56	-	-
		21-4	22	-	-
	3.0-14.0m: Saprolite A greenish yellow to greenish grey colored, saprolite	21-5	25	-	-
		21-6	27	-	-
		21-7	39	-	-
		21-8	28	-	-
		21-9	25	-	-
		21-10	34	39	-
10	11.0-12.0m: chips of quartz veinlets	21-11	26	-	-
		21-12	52	-	-
15	14.0-26.0m: Saprolite B greenish grey to light grey colored saprolite including chips of weathered granodiorite	21-13	58	-	-
		21-14	30	-	-
		21-15	14	-	-
		21-16	29	-	-
		21-17	38	-	-
		21-18	31	-	-
		21-19	17	-	-
		21-20	16	19	-
		21-21	15	-	-
		21-22	218	-	-
		21-23	53	-	-
		21-24	51	-	-
		21-25	92	-	-
		21-26	162	-	-
20	26.0-31.0m: Fine grained granodiorite light grey colored, fine to medium grained granodiorite, or quartz diorite, including small amount of pyrite and / or pyrrhotite total amount of sulfides = 0 % to 1%	21-27	22	-	-
		21-28	25	-	-
		21-29	63	-	-
		21-30	72	54	-
		21-31	65	-	-
		21-32	66	-	-
		21-33	253	-	-
		21-34	100	-	-
		21-35	28	-	-
		21-36	110	-	-
25	31.0-33.0m: Fault ? oxide zone, dark grey colored, fine grained rock	21-37	1,154	12	0
		21-38	578	-	-
		21-39	24	-	-
		21-40	25	290	23
		21-41	7	-	-
		21-42	8	-	-
		21-43	15	-	-
		21-44	14	-	-
		21-45	7	-	-
		21-46	7	-	-
30	33.0-60.0m: meta-sandstone dark grey colored, fine grained meta-greywacke with weak dissemination of pyrite	21-47	7	-	-
		21-48	73	-	-
		21-49	155	-	-
		21-50	13	7	-
		21-51	12	-	-
		21-52	55	-	-
		21-53	17	-	-
		21-54	7	-	-
		21-55	5	-	-
		21-56	6	-	-
35		21-57	18	-	-
		21-58	5	-	-
		21-59	5	-	-
		21-60	10	8	-
40					
45					
50					
55					
60					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
4.0	0.0-4.0m: Carapace reddish brown colored soil with nodules	22-1	264	-	-
		22-2	1,066	274	0
		22-3	53	-	-
		22-4	393	-	-
24.0	4.0-24.0m: Saprolite A greenish grey colored, fine to medium grained saprolite, with no rock chips	22-5	157	-	-
		22-6	36	-	-
		22-7	25	-	-
		22-8	23	-	-
		22-9	84	-	-
		22-10	348	386	-
		22-11	260	-	-
		22-12	198	-	-
		22-13	41	-	-
		22-14	20	-	-
		22-15	20	-	-
		22-16	24	-	-
		22-17	15	-	-
		22-18	15	-	-
		22-19	16	-	-
		22-20	15	19	-
		22-21	12	-	-
		22-22	17	-	-
		22-23	12	-	-
		22-24	3	-	-
60.0	28.0-60.0m: granodiorite light grey colored, fine to medium grained granodiorite (or quartz diorite), not altered, sometimes it shows porphyritic texture with big plagioclase phenocrysts total amount of pyrite and pyrrhotite = 0 % to 1%	22-25	100	-	-
		22-26	117	-	-
		22-27	56	-	-
		22-28	36	-	-
		22-29	25	-	-
		22-30	21	25	-
		22-31	8	-	-
		22-32	202	-	-
		22-33	52	-	-
		22-34	5	-	-
		22-35	57	-	-
		22-36	8	-	-
		22-37	10	-	-
		22-38	10	-	-
		22-39	186	-	-
		22-40	11	8	-
		22-41	85	-	-
		22-42	98	-	-
		22-43	18	-	-
		22-44	13	-	-
		22-45	10	-	-
		22-46	7	-	-
		22-47	16	-	-
		22-48	47	-	-
		22-49	55	-	-
		22-50	150	424	-
22-51	25	-	-		
22-52	54	-	-		
22-53	196	-	-		
22-54	17	-	-		
22-55	213	-	-		
22-56	91	-	-		
22-57	47	-	-		
22-58	173	-	-		
22-59	29	-	-		
22-60	6	11	-		

Column (m)	Description	Sample No.	Au			
			ppb	ppb	ppb	
2.0 5 8.0 10 15 20 25	0.0-2.0m: Carapace brown to reddish brown colored soil with iron nodules	23-1	185	-	-	
		23-2	583	-	-	
	2.0-8.0m: Saprolite A yellow to brownish yellow colored saprolite	23-3	52	-	-	
		23-4	24	-	-	
		23-5	60	-	-	
		23-6	40	-	-	
		23-7	69	-	-	
		23-8	48	-	-	
		23-9	54	-	-	
	8.0-25.0m: Saprolite B greenish grey to yellowish grey colored saprolite including chips of weathered sandstone	23-10	42	46	-	
		23-11	35	-	-	
		23-12	20	-	-	
		23-13	20	-	-	
		23-14	32	-	-	
		23-15	57	-	-	
		23-16	66	-	-	
		23-17	56	-	-	
		23-18	65	-	-	
		23-19	41	-	-	
	23-20	50	50	-		
	25.0	25.0-37.0m: Meta-sandstone dark grey to grey colored meta-greywacke including quartz grains (1mm of diameter) with pyrite and arsenopyrite dissemination total amount of sulfides = 1 % to 3%	23-21	8	-	-
			23-22	30	-	-
			23-23	26	-	-
			23-24	12	-	-
			23-25	15	-	-
23-26			9	-	-	
23-27			28	-	-	
23-28			35	-	-	
23-29			23	-	-	
23-30			17	22	-	
37.0	37.0-48.0m: Silicified zone ? slightly silicified meta-sandstone, with pyrite and arsenopyrite dissemination quartzite ??	23-31	24	-	-	
		23-32	25	-	-	
		23-33	41	-	-	
		23-34	17	-	-	
		23-35	26	-	-	
		23-36	87	-	-	
		23-37	219	-	-	
		23-38	558	-	-	
		23-39	311	-	-	
		23-40	400	300	-	
48.0	48.0-57.0m: Meta-sandstone dark grey to grey colored meta-greywacke including quartz grains (1mm of diameter) with pyrite dissemination	23-41	41	-	-	
		23-42	57	-	-	
		23-43	285	-	-	
		23-44	485	-	-	
		23-45	52	-	-	
		23-46	178	-	-	
		23-47	720	-	-	
57.0		23-48	431	-	-	
		23-49	69	-	-	
		23-50	33	35	-	
		23-51	7	-	-	
		23-52	8	-	-	
		23-53	4	-	-	
		23-54	13	-	-	
		23-55	14	-	-	
		23-56	4	-	-	
		23-57	31	-	-	

Column (m)	Description	Sample No.	Au			
			ppb	ppb	ppb	
5	0.0-1.0m: Carapace brown colored soil with iron nodules	24-1	32	-	-	
		24-2	44	-	-	
		24-3	42	-	-	
	2.0-8.0m: Saprolite A greenish grey to yellow colored, fine grained powder	24-4	34	-	-	
		24-5	35	-	-	
		24-6	20	-	-	
		24-7	13	-	-	
		24-8	24	-	-	
		24-9	33	-	-	
	10	8.0-16.0m: Saprolite B greenish grey colored, medium grained powder including chips of meta-sandstone	24-10	23	9	-
			24-11	24	-	-
			24-12	33	-	-
			24-13	22	-	-
			24-14	16	-	-
			24-15	163	-	-
24-16			72	-	-	
24-17			351	-	-	
24-18			267	-	-	
24-19			79	-	-	
20	16.0-38.0m: Meta-sandstone dark grey colored, fine to medium grained sandstone (greywacke), including quartz grains, with sulfide dissemination, sulfide amount is 1% to 3 %, sometimes higher than 3 %	24-20	44	36	-	
		24-21	195	-	-	
		24-22	148	-	-	
		24-23	24	-	-	
		24-24	22	-	-	
		24-25	66	-	-	
		24-26	26	-	-	
		24-27	25	-	-	
		24-28	12	-	-	
		24-29	10	-	-	
		24-30	8	18	-	
		24-31	238	-	-	
		24-32	6	-	-	
		24-33	2	-	-	
		24-34	0	-	-	
35	38.0-60.0m: Alternation beds of meta-sandstone and peritic schist dark grey colored, alternation beds of meta-sandstone (greywacke) and peritic schist, with sulfide dissemination, sulfide amount is 1% to 3 %, sometimes higher than 3 % rock chip of "24-48" contains quartz veinlets	24-35	4	-	-	
		24-36	7	-	-	
		24-37	12	-	-	
		24-38	12	-	-	
		24-39	5	-	-	
		24-40	15	13	-	
		24-41	9	-	-	
		24-42	11	-	-	
		24-43	16	-	-	
		24-44	12	-	-	
		24-45	25	-	-	
		24-46	4	-	-	
		24-47	24	-	-	
		24-48	6	-	-	
		24-49	15	-	-	
50	38.0-60.0m: Alternation beds of meta-sandstone and peritic schist dark grey colored, alternation beds of meta-sandstone (greywacke) and peritic schist, with sulfide dissemination, sulfide amount is 1% to 3 %, sometimes higher than 3 % rock chip of "24-48" contains quartz veinlets	24-50	25	22	-	
		24-51	21	-	-	
		24-52	13	-	-	
		24-53	11	-	-	
		24-54	25	-	-	
		24-55	30	-	-	
		24-56	34	-	-	
		24-57	12	-	-	
		24-58	12	-	-	
		24-59	8	-	-	
		24-60	15	20	-	

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 11.0	0.0-11.0m: Carapace reddish brown colored carapace, medium grained, with a small amount of iron nodules	74-1	11	-	-
		74-2	2	-	-
		74-3	3	-	-
		74-4	8	-	-
		74-5	9	8	-
		74-6	17	-	-
		74-7	13	-	-
		74-8	14	-	-
		74-9	22	-	-
		74-10	28	-	-
		74-11	27	-	-
15 20 25 30 31.0	11.0-31.0m: Saprolite A reddish brown colored saprolite, medium-coarse grained, homogeneous, without rock chips	74-12	104	-	-
		74-13	33	-	-
		74-14	20	-	-
		74-15	226	27	16
		74-16	13	-	-
		74-17	33	-	-
		74-18	17	-	-
		74-19	17	-	-
		74-20	34	-	-
		74-21	51	-	-
		74-22	18	-	-
		74-23	27	-	-
		74-24	30	-	-
		74-25	34	23	-
		74-26	105	-	-
		74-27	462	-	-
		74-28	54	-	-
		74-29	400	-	-
		74-30	60	-	-
		35 40 45 50 50.0	31.0-50.0m: Saprolite A greenish light gray to yellowish gray colored saprolite, very fine grained powder, without rock chips	74-31	43
74-32	83			-	-
74-33	46			-	-
74-34	31			-	-
74-35	4			18	-
74-36	24			-	-
74-37	33			-	-
74-38	36			-	-
74-39	26			-	-
74-40	9			-	-
74-41	12			-	-
74-42	6			-	-
74-43	8			-	-
74-44	8			-	-
74-45	194			130	-
74-46	4			-	-
74-47	3			-	-
74-48	1			-	-
74-49	3			-	-
55 60 60.0	50.0-60.0m: Saprolite B yellowish light gray colored saprolite, coarse grained, including a lot of chips of weathered sandstone, (diameter: 3-4mm, sometimes black colored, meta-graywacke and peritic schist can be observed, iron-oxides along fracture) 59.0-60.0m: Oxide zone	74-50	130	-	-
		74-51	140	-	-
		74-52	19	-	-
		74-53	16	-	-
		74-54	2	-	-
		74-55	4	7	-
		74-56	9	-	-
		74-57	0	-	-
		74-58	24	-	-
		74-59	3	-	-
		74-60	2	-	-

KRC-75

Coordination: N1500 E400

Elevation: 382m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-8.0m: Carapace red to brown colored hard carapace, including a lot of lateritic gravels	75-1	32	-	-
		75-2	6	-	-
		75-3	6	-	-
		75-4	4	-	-
		75-5	6	15	-
		75-6	121	-	-
		75-7	28	-	-
		75-8	25	-	-
10	8.0-15.0m: Mottled clay brown to reddish brown colored clay, including some lateritic gravels	75-9	51	-	-
		75-10	51	-	-
		75-11	34	-	-
		75-12	24	-	-
		75-13	21	-	-
		75-14	15	-	-
		75-15	205	99	-
15	15.0-54.0m: Saprolite A brownish to yellowish brown to greenish gray colored saprolite, fine grained, soft	75-16	239	-	-
		75-17	15	-	-
		75-18	459	-	-
		75-19	21	-	-
		75-20	58	-	-
		75-21	39	-	-
		75-22	8	-	-
		75-23	11	-	-
		75-24	10	-	-
		75-25	10	5	-
		75-26	7	-	-
		75-27	4	-	-
		75-28	19	-	-
		75-29	12	-	-
		75-30	45	-	-
		75-31	9	-	-
		75-32	49	-	-
		75-33	9	-	-
		75-34	7	-	-
		75-35	6	6	-
		75-36	16	-	-
		75-37	14	-	-
		75-38	11	-	-
		75-39	3	-	-
		75-40	6	-	-
		75-41	6	-	-
		75-42	5	-	-
		75-43	4	-	-
		75-44	9	-	-
		75-45	8	7	-
		75-46	10	-	-
		75-47	6	-	-
75-48	1	-	-		
75-49	6	-	-		
75-50	0	-	-		
75-51	6	-	-		
75-52	15	-	-		
75-53	11	-	-		
75-54	12	-	-		
55	54.0-60.0m: Saprolite B green colored saprolite, including chips of deeply weathered schistosed meta-sandstone	75-55	10	6	-
		75-56	0	-	-
		75-57	0	-	-
		75-58	0	-	-
		75-59	0	-	-
		75-60	54	-	-
60	60.0				