

Apc.22 Diagraphie géologique des trous de forages à circulation

inverse (RC) dans le Secteur de Kékoro

+

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-5.0m: Surface soil yellowish brown on the top, brown on the bottom, this layer contains white clay and some quartz fragments	51-1	46	-	-
		51-2	42	-	-
		51-3	40	-	-
		51-4	31	-	-
		51-5	42	-	-
6.0	5.0-6.0m: Crust and Carapace reddish brown colored, with iron nodules	51-6	204	-	-
		51-7	221	-	-
		51-8	404	-	-
		51-9	33	-	-
10	6.0-12.0m: Saprolite A greenish grey colored saprolite	51-10	11	6	-
		51-11	7	-	-
		51-12	7	-	-
15	12.0-36.0m: Saprolite B greenish grey colored saprolite, including a lot of chips of weathered meta-greywacke	51-13	7	-	-
		51-14	6	-	-
		51-15	5	-	-
		51-16	3	-	-
		51-17	6	-	-
		51-18	5	-	-
		51-19	5	-	-
		51-20	7	7	-
		51-21	29	-	-
		51-22	5	-	-
		51-23	5	-	-
		51-24	3	-	-
		51-25	4	-	-
		51-26	21	-	-
		51-27	6	-	-
20		51-28	3	-	-
		51-29	8	-	-
		51-30	5	6	-
		51-31	8	-	-
		51-32	3	-	-
		51-33	2	-	-
		51-34	7	-	-
		51-35	11	-	-
		51-36	5	-	-
		51-37	4	-	-
25	36.0-47.0m: Meta-sandstone dark grey colored meta-greywacke, with disseminated pyrite, with joint filling pyrite, total amount of pyrite is 1 % to 2 %	51-38	4	-	-
		51-39	2	-	-
		51-40	4	2	-
		51-41	9	-	-
		51-42	10	-	-
		51-43	3	-	-
		51-44	7	-	-
		51-45	3	-	-
		51-46	7	-	-
		51-47	13	-	-
30	47.0-51.0m: Peritic schist dark grey colored peritic schist, with disseminated pyrite, with schistosity filling pyrite total amount of pyrite is 2 %	51-48	12	-	-
		51-49	16	-	-
		51-50	11	10	-
		51-51	7	-	-
35	51.0-57.0m: Meta-sandstone dark grey colored meta-greywacke, with disseminated pyrite, total amount of pyrite is 0 % to 1 %	51-52	5	-	-
		51-53	7	-	-
		51-54	5	-	-
		51-55	6	-	-
		51-56	8	-	-
40	57.0-60.0m: Peritic schist dark grey colored peritic schist, with disseminated pyrite, with schistosity filling pyrite total amount of pyrite is 1 % to 2 %	51-57	10	-	-
		51-58	12	-	-
		51-59	10	-	-
		51-60	14	7	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0	0.0-1.0m: Surface soil brown colored, with laterite nodules	52-1	26	-	-
3.0		52-2	96	-	-
		52-3	31	-	-
5	2.0-3.0m: Carapace reddish brown colored, with a lot of lateritic nodules	52-4	28	-	-
		52-5	11	-	-
		52-6	385	-	-
		52-7	19	-	-
10	3.0-14.0m: Saprolite A greenish grey colored saprolite	52-8	32	-	-
		52-9	19	-	-
		52-10	10	7	-
		52-11	302	-	-
		52-12	23	-	-
		52-13	27	-	-
15	14.0-22.0m: Saprolite B greenish grey colored saprolite, including chips of weathered meta-greywacke	52-14	22	-	-
		52-15	14	-	-
		52-16	11	-	-
		52-17	9	-	-
		52-18	10	-	-
		52-19	6	-	-
		52-20	16	8	-
		52-21	5	-	-
20	22.0-26.0m: Saprolite B greenish grey colored saprolite, including chips of silicified? meta-greywacke (quartz schist?)	52-22	19	-	-
		52-23	8	-	-
		52-24	5	-	-
		52-25	24	-	-
		52-26	11	-	-
		52-27	17	-	-
25	26.0-34.0m: Saprolite B greenish grey colored saprolite, including chips of weathered meta-greywacke	52-28	8	-	-
		52-29	6	-	-
		52-30	7	12	-
		52-31	23	-	-
		52-32	50	-	-
		52-33	30	-	-
30	34.0-38.0m: Alternation beds of meta-sandstone and peritic schist dark grey colored rock, including 1 % to 4 % of disseminated pyrite	52-34	55	-	-
		52-35	23	-	-
		52-36	15	-	-
		52-37	8	-	-
		52-38	18	-	-
		52-39	10	-	-
35	38.0-45.0m: Meta-sandstone meta-sandstone with 1 % to 4 % of disseminated pyrite	52-40	7	21	-
		52-41	37	-	-
		52-42	218	-	-
		52-43	0	-	-
		52-44	42	-	-
		52-45	1	-	-
40	45.0-52.0m: Alternation beds of meta-sandstone and peritic schist dark grey colored rock, with schistosity filling and disseminated pyrite, 1% to 4% in peritic schist, 1% in meta-sandstone	52-46	5	-	-
		52-47	31	-	-
		52-48	5	-	-
		52-49	4	-	-
		52-50	8	1	-
		52-51	1	-	-
45	52.0-56.0m: Meta-sandstone meta-sandstone with great amount of disseminated pyrite, more than 4 % ?	52-52	0	-	-
		52-53	0	-	-
		52-54	0	-	-
		52-55	0	-	-
		52-56	0	-	-
		52-57	0	-	-
50	56.0-60.0m: Peritic schist peritic schist with schistosity filling pyrite and disseminated pyrite, more than 4 % ?	52-58	2	-	-
		52-59	0	-	-
		52-60	1	2	-
55					
60					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
0.0-3.0	0.0-3.0m: Carapace reddish brown colored soil with laterite nodules	53-1	17	-	-
		53-2	14	-	-
		53-3	17	-	-
3.0-20.0	3.0-20.0m: Saprolite A greenish brown colored saprolite	53-4	23	-	-
		53-5	14	-	-
		53-6	19	-	-
		53-7	13	-	-
		53-8	15	-	-
		53-9	61	-	-
		53-10	37	28	-
		53-11	32	-	-
		53-12	6	-	-
		53-13	5	-	-
		53-14	7	-	-
		53-15	10	-	-
		53-16	4	-	-
		53-17	40	-	-
		53-18	36	-	-
		53-19	71	-	-
		53-20	16	-	-
20.0-39.0	20.0-39.0m: Saprolite B greenish grey colored saprolite, including chips of meta-greywacke and of peritic schist	53-21	3	-	-
		53-22	4	-	-
		53-23	8	-	-
		53-24	14	-	-
		53-25	19	-	-
		53-26	13	-	-
		53-27	12	-	-
		53-28	8	-	-
		53-29	14	-	-
		53-30	30	4	-
		53-31	1	-	-
		53-32	6	-	-
		53-33	2	-	-
		53-34	6	-	-
		53-35	4	-	-
		53-36	5	-	-
		53-37	8	-	-
		53-38	7	-	-
		53-39	2	-	-
		39.0-57.0	39.0-57.0m: Peritic schist dark grey colored schist, with schistosity filling and disseminated fine grained pyrite, 1% to 3% ?	53-40	4
53-41	20			-	-
53-42	19			-	-
53-43	14			-	-
53-44	21			-	-
53-45	17			-	-
53-46	20			-	-
53-47	17			-	-
53-48	19			-	-
53-49	41			-	-
53-50	17			18	-
57.0-58.0	57.0-58.0m: Meta-sandstone grey colored rock, with 1 % to 3 % ? pyrite	53-51	25	-	-
		53-52	30	-	-
		53-53	66	-	-
		53-54	78	-	-
		53-55	37	-	-
		53-56	47	-	-
58.0-60.0	58.0-60.0m: Peritic schist dark grey colored schist, with schistosity filling and disseminated fine grained pyrite, 1% to 3% ?	53-57	36	-	-
		53-58	8	-	-
		53-59	11	-	-
		53-60	13	14	-

Column (m)	Description	Sample No.	Au				
			ppb	ppb	ppb		
5	0.0-5.0m: Carapace reddish brown colored soil with laterite nodules at the depth of 1m to 2m	55-1	0	-	-		
		55-2	16	-	-		
		55-3	17	-	-		
		55-4	18	-	-		
		55-5	19	-	-		
10	5.0-32.0m: Saprolite A 5.0m to 11.0m: brown colored silty soil	55-6	32	-	-		
		55-7	83	-	-		
		55-8	180	-	-		
		55-9	329	-	-		
		55-10	229	493	-		
		55-11	241	-	-		
		15	11.0m to 19.0m: yellowish light grey colored silty soil	55-12	40	-	-
				55-13	26	-	-
				55-14	39	-	-
				55-15	58	-	-
55-16	55			-	-		
55-17	134			-	-		
55-18	33			-	-		
55-19	21			-	-		
20	19.0m to 27.0m: grey to brown colored silty soil	55-20	7	39	-		
		55-21	32	-	-		
		55-22	25	-	-		
		55-23	25	-	-		
		55-24	53	-	-		
		55-25	27	-	-		
		55-26	42	-	-		
		55-27	36	-	-		
		55-28	30	-	-		
		55-29	27	-	-		
30	27.0m to 32.0m: greenish grey to light grey to brown colored silty soil	55-30	62	50	-		
		55-31	95	-	-		
		55-32	33	-	-		
		55-33	36	-	-		
		55-34	37	-	-		
		55-35	46	-	-		
		55-36	21	-	-		
		55-37	27	-	-		
		55-38	251	-	-		
		55-39	539	-	-		
40	32.0-47.0m: Saprolite B greenish light grey colored fine grained soil, with chips of rock fragments (diameter: 5mm)	55-40	210	193	-		
		55-41	69	-	-		
		55-42	44	-	-		
		55-43	29	-	-		
		55-44	27	-	-		
		55-45	11	-	-		
		55-46	71	-	-		
		55-47	10	-	-		
		55-48	112	-	-		
		55-49	478	-	-		
50	47.0-54.0m: Peritic schist dark grey colored schist, with schistosity filling and disseminated fine grained pyrite, 1% to 3% with small amount of Fe-oxides	55-50	525	714	-		
		55-51	1,607	71	69		
		55-52	494	-	-		
		55-53	69	-	-		
		55-54	144	-	-		
55	54.0-60.0m: Alternation beds dark grey colored to greenish light grey colored alternation beds of meta-sandstone and peritic schist with schistosity filling and disseminated fine grained pyrite, 2% to 3% or more, including arsenopyrite with quartz veinlets	55-55	95	-	-		
		55-56	1,538	78	103		
		55-57	33	-	-		
		55-58	41	-	-		
		55-59	315	-	-		
		55-60	166	158	-		
60							

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-4.0m: Crust and carapace reddish brown colored, hard material with lateritic nodules	56-1	55	-	-
		56-2	35	-	-
		56-3	512	-	-
		56-4	83	-	-
5	4.0-8.0m: Carapace reddish brown colored carapace including white clay and lateritic nodules	56-5	158	-	-
		56-6	889	-	-
		56-7	117	-	-
		56-8	120	-	-
10	8.0-32.0m: Mottled clay brown to yellowish brown colored soft material, with no lateritic nodules	56-9	137	-	-
		56-10	50	91	-
		56-11	63	-	-
		56-12	68	-	-
		56-13	103	-	-
		56-14	177	-	-
		56-15	190	-	-
		56-16	330	-	-
		56-17	95	-	-
		56-18	109	-	-
20		56-19	76	-	-
		56-20	253	614	-
		56-21	166	-	-
		56-22	405	-	-
		56-23	2,550	2,854	3,189
		56-24	117	-	-
		56-25	61	-	-
		56-26	16,470	11,451	8,709
		56-27	53	-	-
		56-28	172	-	-
30		56-29	1,505	2,306	1,303
		56-30	1,052	1,016	-
		56-31	123	-	-
		56-32	613	-	-
		56-33	135	-	-
		56-34	359	-	-
		56-35	172	-	-
		56-36	304	-	-
		56-37	210	-	-
		56-38	130	-	-
35	32.0-38.0m: Saprolite A greenish grey colored fine to midium grained saprolite	56-39	259	-	-
		56-40	122	-	-
		56-41	433	-	-
		56-42	230	-	-
		56-43	328	-	-
		56-44	48	-	-
		56-45	137	-	-
		56-46	138	-	-
		56-47	174	-	-
		56-48	333	-	-
40	38.0-48.0m: Saprolite B greenish grey to bluish grey colored midium grained material including rock chips	56-49	2,450	1,880	2,263
		56-50	6,940	7,430	-
		56-51	3,170	3,650	3,840
		56-52	662	-	-
		56-53	227	-	-
		56-54	232	-	-
		56-55	191	-	-
		56-56	181	-	-
		56-57	150	-	-
		56-58	71	-	-
45		56-59	115	-	-
		56-60	127	104	-
50	48.0-60.0m: Granodiorite light grey to greenish grey colored granodiorite, with medium grained facies and fine grained facies, partly chloritized, with dissemination of pyrite and traces of arsenopyrite, sulfide= 0 % to 2 %				
55	water				
60	60.0				

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-5.0m: Alluvial sediments brown colored soil, includig gravels	57-1	44	-	-
		57-2	174	-	-
		57-3	56	-	-
		57-4	35	-	-
		57-5	415	-	-
6.0	5.0-6.0m: Carapace	57-6	19,948	19,817	16,971
10	reddish brown colored carapace including lateritic nodules and quartz fragments	57-7	284	-	-
		57-8	95	-	-
		57-9	151	-	-
10.0	6.0-10.0m: Mottled clay	57-10	83	85	-
15	brownish yellowish colored clay with no lateritic nodules	57-11	349	-	-
		57-12	1,130	1,012	1,474
		57-13	1,099	996	1,371
		57-14	814	-	-
		57-15	1,540	2,244	2,194
		57-16	372	-	-
		57-17	388	-	-
		57-18	166	-	-
		57-19	206	-	-
		57-20	372	214	-
		57-21	1,132	1,602	1,166
		57-22	395	-	-
		57-23	87	-	-
57-24	219	-	-		
57-25	42	-	-		
57-26	78	-	-		
57-27	1,823	3,980	2,606		
57-28	403	-	-		
57-29	882	-	-		
57-30	384	814	1,008		
30.0	30.0-36.0m: Saprolite B	57-31	426	-	-
35	greenish grey colored fine grained powder, including chips of meta-sandstone that is stained by Fe-oxides	57-32	104	-	-
		57-33	233	-	-
		57-34	100	-	-
		57-35	97	-	-
		57-36	282	-	-
		57-37	166	-	-
36.0	36.0-60.0m: Meta-sandstone	57-38	60	-	-
40	dark grey colored meta-sandstone, with fine grained pyrite dissemination, with pyrite veinlets, total amount of sulfide = 1 % to 2 %, with traces of quartz fragments and carbonate fragments, with traces of rock chips of green schist	57-39	196	-	-
		57-40	362	225	-
		57-41	128	-	-
		57-42	100	-	-
		57-43	29	-	-
		57-44	102	-	-
		57-45	41	-	-
		57-46	34	-	-
		57-47	29	-	-
		57-48	35	-	-
		57-49	23	-	-
		57-50	64	37	-
57-51	40	-	-		
57-52	24	-	-		
57-53	78	-	-		
57-54	42	-	-		
57-55	46	-	-		
57-56	63	-	-		
57-57	74	-	-		
57-58	53	-	-		
57-59	88	-	-		
60	60.0	57-60	52	39	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
0.0-3.0m	0.0-3.0m: Carapace reddish brown colored, with iron nodules	42-1	40	-	-
		42-2	36	-	-
		42-3	31	-	-
3.0-13.0m	3.0-13.0m: Saprolite A greenish grey colored powder with no rock fragments	42-4	49	-	-
		42-5	111	-	-
		42-6	99	-	-
		42-7	16	-	-
		42-8	16	-	-
		42-9	16	-	-
		42-10	15	3	-
		42-11	30	-	-
		42-12	39	-	-
		42-13	15	-	-
13.0-27.0m	13.0-27.0m: Saprolite B greenish grey colored saprolite, including a lot of chips of peritic schist	42-14	23	-	-
		42-15	14	-	-
		42-16	21	-	-
		42-17	25	-	-
		42-18	12	-	-
		42-19	9	-	-
		42-20	1	9	-
		42-21	16	-	-
		42-22	3	-	-
		42-23	27	-	-
		42-24	20	-	-
		42-25	36	-	-
		42-26	37	-	-
		42-27	16	-	-
27.0-34.0m	27.0-34.0m: Peritic schist dark grey colored peritic schist, with joint filling pyrite and disseminated pyrite, total amount of pyrite is 0 % to 2 %	42-28	13	-	-
		42-29	13	-	-
		42-30	15	21	-
		42-31	12	-	-
		42-32	5	-	-
		42-33	8	-	-
		42-34	25	-	-
		42-35	0	-	-
		42-36	4	-	-
		42-37	33	-	-
34.0-36.0m	34.0-36.0m: Quarz diorite ? dark grey colored, medium grained rock	42-38	18	-	-
		42-39	10	-	-
		42-40	10	6	-
		42-41	26	-	-
		42-42	54	-	-
		42-43	11	-	-
		42-44	35	-	-
		42-45	0	-	-
		42-46	50	-	-
		42-47	47	-	-
36.0-54.0m	36.0-54.0m: Meta-sandstone dark grey colored meta-graywacke, with disseminated pyrite, with fracture filling pyrite, total amount of pyrite is 1 % to 3 %	42-48	4	-	-
		42-49	9	-	-
		42-50	16	27	-
		42-51	0	-	-
		42-52	10	-	-
		42-53	3	-	-
		42-54	13	-	-
		42-55	4	-	-
		42-56	52	-	-
		42-57	34	-	-
54.0-60.0m	36.0-54.0m: Alternation beds dark grey colored alternation beds of meta-graywacke and peritic schist, with disseminated pyrite, with fracture filling pyrite, total amount of pyrite is 1 % to 3 %	42-58	28	-	-
		42-59	53	-	-
		42-60	47	32	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
3.0	0.0-3.0m: Carapace reddish brown colored, with laterite nodules	43-1	2	-	-
		43-2	16	-	-
		43-3	3,093	8,154	829
5	3.0-18.0m: Saprolite A greenish grey colored, soft and fine grained powder	43-4	110	-	-
		43-5	25	-	-
		43-6	73	-	-
		43-7	81	-	-
		43-8	9	-	-
		43-9	15	-	-
		43-10	184	236	-
		43-11	38	-	-
		43-12	46	-	-
		43-13	27	-	-
10		43-14	56	-	-
		43-15	24	-	-
		43-16	39	-	-
		43-17	21	-	-
		43-18	22	-	-
		43-19	34	-	-
		43-20	23	34	-
		43-21	18	-	-
		43-22	59	-	-
		43-23	42	-	-
15		43-24	50	-	-
		43-25	68	-	-
		43-26	162	-	-
		43-27	42	-	-
		43-28	56	-	-
		43-29	41	-	-
		43-30	27	11	-
		43-31	22	-	-
		43-32	21	-	-
		43-33	75	-	-
20	18.0-28.0m: Saprolite B greenish grey colored saprolite, including chips of sandstone schist of greenish grey colored	43-34	108	-	-
		43-35	78	-	-
		43-36	9	-	-
		43-37	57	-	-
		43-38	68	-	-
		43-39	59	-	-
		43-40	51	50	-
		43-41	22	-	-
		43-42	10	-	-
		43-43	17	-	-
25		43-44	17	-	-
		43-45	19	-	-
		43-46	39	-	-
		43-47	26	-	-
		43-48	11	-	-
		43-49	17	-	-
		43-50	12	9	-
		43-51	61	-	-
		43-52	18	-	-
		43-53	6	-	-
30	28.0-60.0m: Sandstone schist grey to greenish grey colored schistose meta-sandstone, with schistosity filling pyrite and disseminated pyrite, total amount of pyrite = 1 % to 3 %, including meta-volcanics ? rock chip of 43-30 is siliceous tuff with strong dissemination of pyrite	43-54	8	-	-
		43-55	38	-	-
		43-56	41	-	-
		43-57	49	-	-
		43-58	30	-	-
		43-59	14	-	-
		43-60	17	22	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
3.0	0.0-3.0m: Carapace reddish brown colored soil with laterite nodules	44-1	1	-	-
		44-2	0	-	-
		44-3	287	-	-
5 10 15 20 23.0	3.0-23.0m: Saprolite A greenish brown colored soft powder, including some fragments of quartz veinlets	44-4	53	-	-
		44-5	179	-	-
		44-6	46	-	-
		44-7	17	-	-
		44-8	28	-	-
		44-9	122	-	-
		44-10	18	10	-
		44-11	24	-	-
		44-12	47	-	-
		44-13	26	-	-
		44-14	103	-	-
		44-15	145	-	-
		44-16	291	-	-
		44-17	81	-	-
		44-18	64	-	-
		44-19	781	-	-
		44-20	58	36	-
44-21	2	-	-		
44-22	5	-	-		
44-23	13	-	-		
25 30 30.0	23.0-30.0m: Saprolite B greenish grey colored saprolite, including chips of meta-sandstone	44-24	136	-	-
		44-25	10	-	-
		44-26	0	-	-
		44-27	0	-	-
		44-28	33	-	-
		44-29	20	-	-
		44-30	26	7	-
		44-31	11	-	-
		44-32	60	-	-
		44-33	25	-	-
35 40 45 50 55 60 60.0	30.0-60.0m: Meta-sandstone dark grey to greenish dark grey colored meta-greywacke, including trace of pyrite, 0 % to 1 %, including meta-volcanics ? 34m to 36m: fracture zone, rock chips are brown colored and oxidized 43m to 46m: fracture zone, rock chips are brown colored and oxidized rock chips of 44-56 are meta-graywacke with strong dissemination of pyrite 56m to 60m: very fine grained glassy rock	44-34	23	-	-
		44-35	12	-	-
		44-36	66	-	-
		44-37	116	-	-
		44-38	86	-	-
		44-39	40	-	-
		44-40	11	2	-
		44-41	10	-	-
		44-42	18	-	-
		44-43	57	-	-
		44-44	573	-	-
		44-45	46	-	-
		44-46	62	-	-
		44-47	13	-	-
		44-48	11	-	-
		44-49	32	-	-
		44-50	15	114	29
44-51	65	-	-		
44-52	22	-	-		
44-53	71	-	-		
44-54	34	-	-		
44-55	35	-	-		
44-56	233	-	-		
44-57	431	-	-		
44-58	126	-	-		
44-59	63	-	-		
44-60	131	27	5		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0	0.0-1.0m: Carapace	45-1	163	-	-
2.0	reddish brown colored soil with laterite nodules	45-2	50	-	-
		45-3	27	-	-
5	1.0-2.0m: Mottled clay	45-4	12	-	-
	brown yellow colored clay	45-5	9	-	-
		45-6	24	-	-
10	2.0-11.0m: Saprolite A	45-7	493	-	-
	greenish grey colored, fine grained powder	45-8	88	-	-
		45-9	32	-	-
		45-10	26	26	-
		45-11	16	-	-
11.0	11.0-40.0m: Saprolite B	45-12	80	-	-
15	greenish grey colored saprolite, including chips of greenish grey colored meta-sandstone ?	45-13	65	-	-
		45-14	28	-	-
		45-15	19	-	-
		45-16	0	-	-
		45-17	8	-	-
		45-18	2	-	-
		45-19	22	-	-
		45-20	34	28	-
		45-21	20	-	-
		45-22	17	-	-
		45-23	14	-	-
		45-24	17	-	-
		45-25	54	-	-
		45-26	45	-	-
		45-27	18	-	-
		45-28	37	-	-
		45-29	34	-	-
		45-30	34	42	-
		45-31	38	-	-
		45-32	61	-	-
45-33	91	-	-		
45-34	105	-	-		
45-35	73	-	-		
45-36	56	-	-		
45-37	39	-	-		
45-38	168	-	-		
45-39	71	-	-		
40	40.0	45-40	183	136	-
45	40.0-60.0m: Meta-sandstone ?	45-41	29	-	-
	dark grey to greenish dark grey colored rock with disseminated pyrite meta-greywacke ? meta-diorite ??, including meta-volcanics ?	45-42	33	-	-
		45-43	31	-	-
		45-44	28	-	-
		45-45	19	-	-
		45-46	19	-	-
		45-47	49	-	-
		45-48	66	-	-
		45-49	55	-	-
		45-50	46	44	-
45-51	66	-	-		
45-52	91	-	-		
45-53	48	-	-		
45-54	42	-	-		
45-55	41	-	-		
45-56	177	-	-		
45-57	90	-	-		
45-58	30	-	-		
45-59	15	-	-		
60	60.0	45-60	22	9	-
	rock chips of 45-59 are meta-graywacke or peritic schist				

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-7.0m: Carapace reddish brown colored hard soil with iron nodules	48-1	22	-	-
		48-2	13	-	-
		48-3	25	-	-
		48-4	38	-	-
		48-5	92	-	-
		48-6	38	-	-
		48-7	149	-	-
10	7.0-32.0m: Saprolite A reddish brown to brown colored, fine grained powder, gradually change to green saprolite oxide zone	48-8	125	-	-
		48-9	131	-	-
		48-10	199	172	-
		48-11	214	-	-
		48-12	325	-	-
		48-13	119	-	-
		48-14	110	-	-
		48-15	165	-	-
		48-16	212	-	-
		48-17	218	-	-
		48-18	183	-	-
		48-19	304	-	-
		48-20	626	239	-
		48-21	144	-	-
		48-22	326	-	-
		48-23	399	-	-
		48-24	184	-	-
		48-25	569	-	-
		48-26	892	583	1,029
		48-27	999	480	891
		48-28	1,098	4,114	857
		48-29	824	-	-
		48-30	583	589	-
48-31	1,754	480	1,303		
48-32	1,162	960	686		
35	32.0-45.0m: Saprolite A greenish grey colored, fine grained saprolite reduced zone	48-33	781	-	-
		48-34	618	-	-
		48-35	228	-	-
		48-36	294	-	-
		48-37	159	-	-
		48-38	134	-	-
		48-39	275	-	-
		48-40	393	836	-
		48-41	2,004	3,600	1,200
		48-42	1,656	3,497	2,331
		48-43	2,966	6,926	3,840
		48-44	6,221	1,303	5,897
45	45.0-58.0m: Saprolite B greenish grey colored, medium grained powder including chips of meta-sandstone ("black tuff")	48-45	1,346	1,646	1,851
		48-46	1,242	-	651
50		48-47	393	-	-
		48-48	2,060	-	1,714
		48-49	2,184	-	2,057
		48-50	677	442	-
		48-51	520	-	-
		48-52	661	-	-
		48-53	410	-	-
		48-54	986	-	-
		48-55	926	-	-
		48-56	249	-	-
55	58.0-60.0m: Meta-sandstone ("black tuff") dark grey colored, fine grained, massive rock including anhedral plagioclase grains, with dissemination of pyrite, amount of pyrite is 3% to 4 %	48-57	280	-	-
		48-58	261	-	-
		48-59	436	-	-
60	water	48-60	257	329	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 14.0	0.0-14.0m: Carapace reddish brown colored hard soil with iron nodules not indurated, including white clay	49-1	5	-	-
		49-2	0	-	-
		49-3	80	-	-
		49-4	13	-	-
		49-5	73	-	-
		49-6	82	-	-
		49-7	45	-	-
		49-8	45	-	-
		49-9	108	-	-
		49-10	145	0	-
		49-11	44	-	-
		49-12	38	-	-
		49-13	48	-	-
		49-14	5,384	746	0
15 20 25 30 35 36.0	14.0-36.0m: Saprolite A brownish grey to greenish grey colored, fine grained powder	49-15	110	-	-
		49-16	81	-	-
		49-17	85	-	-
		49-18	234	-	-
		49-19	267	-	-
		49-20	86	90	-
		49-21	78	-	-
		49-22	73	-	-
		49-23	70	-	-
		49-24	70	-	-
		49-25	167	-	-
		49-26	53	-	-
		49-27	83	-	-
		49-28	49	-	-
		49-29	688	-	-
		49-30	9	81	-
		49-31	227	-	-
		49-32	163	-	-
		49-33	90	-	-
		49-34	102	-	-
		49-35	57	-	-
		49-36	18	-	-
36.0 40 44.0	36.0-44.0m: Saprolite B greenish grey colored, medium grained powder including diorite chips	49-37	104	-	-
		49-38	937	1,066	617
		49-39	435	-	-
		49-40	157	68	-
		49-41	225	-	-
		49-42	147	-	-
		49-43	121	-	-
		49-44	86	-	-
45 50 55 60	44.0-60.0m: Diorite (or Granodiorite) dark grey colored, medium grained diorite or granodiorite with disseminated pyrite, amount of pyrite is 0 % to 1 %	49-45	135	-	-
		49-46	103	-	-
		49-47	190	-	-
		49-48	139	-	-
		49-49	75	-	-
		49-50	71	-	-
		49-51	100	-	-
		49-52	70	-	-
		49-53	75	-	-
		49-54	67	-	-
		49-55	37	-	-
		49-56	22	-	-
		49-57	20	-	-
		49-58	59	-	-
		49-59	37	-	-
		49-60	29	54	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 13.0	0.0-13.0m: Carapace reddish brown colored, coarse grained carapace, with a lot of iron nodules (diameter: 2-10mm)	84-1	30	-	-
		84-2	19	-	-
		84-3	15	-	-
		84-4	24	-	-
		84-5	33	-	-
		84-6	33	-	-
		84-7	54	-	-
		84-8	82	-	-
		84-9	132	-	-
		84-10	57	73	-
		84-11	109	-	-
		84-12	162	-	-
		84-13	185	-	-
15 20 25 30 30.0	13.0-30.0m: Saprolite A reddish brown to yellowish brown colored, very fine grained saprolite, with no rock chips	84-14	391	-	-
		84-15	502	-	-
		84-16	971	-	-
		84-17	471	-	-
		84-18	600	-	-
		84-19	482	-	-
		84-20	863	245	-
		84-21	84	-	-
		84-22	99	-	-
		84-23	204	-	-
		84-24	203	-	-
		84-25	666	-	-
		84-26	9,460	9,531	10,491
		84-27	10,290	8,777	9,189
		84-28	886	-	-
		84-29	144	-	-
		84-30	110	125	-
35 39.0	30.0-39.0m: Saprolite B yellowish brown colored, medium grained saprolite, with a lot of chips of strongly weathered sandstone (diameter: 5-20mm)	84-31	1,440	1,063	1,200
		84-32	309	-	-
		84-33	424	-	-
		84-34	8,120	7,577	9,771
		84-35	22,100	17,109	21,360
		84-36	11,470	8,811	9,360
		84-37	2,699	2,366	2,880
		84-38	416	-	-
40 41.0	39.0-41.0m: Meta-sandstone brown to dark gray colored, strongly weathered meta-sandstone, with oxidized sandstone and strongly weathered peritic schist	84-39	182	-	-
		84-40	104	144	-
45 50 55 60 60.0	41.0-60.0m: Granodiorite fine grained granodiorite (plagioclase>>biotite >hornblende), with weak dissemination of pyrite, contents of pyrite is less than 1%, fresh, chloritization is not seen	84-41	899	-	-
		84-42	182	-	-
		84-43	444	-	-
		84-44	654	-	-
		84-45	147	-	-
		84-46	53	-	-
		84-47	1,104	25	0
		84-48	728	-	-
		84-49	67	-	-
		84-50	174	99	-
		84-51	151	-	-
		84-52	61	-	-
84-53	58	-	-		
84-54	81	-	-		
84-55	57	-	-		
84-56	45	-	-		
84-57	94	-	-		
84-58	48	-	-		
84-59	130	-	-		
84-60	32	117	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-13.0m: Carapace reddish brown colored carapace, including a lot of iron nodules	85-1	19	-	-
		85-2	60	-	-
		85-3	27	-	-
		85-4	26	-	-
		85-5	10	-	-
		85-6	11	-	-
		85-7	45	-	-
		85-8	21	-	-
		85-9	169	-	-
		85-10	3,591	2,518	-
10		85-11	307	-	-
		85-12	242	-	-
13.0		85-13	207	-	-
15	13.0-15.0m: Mottled clay brown colored clay, including some lateritic nodules	85-14	122	-	-
		85-15	181	-	-
20	15.0-46.0m: Saprolite A 15.0-38.0m: brown to pinkish brown colored, fine grained saprolite	85-16	180	-	-
		85-17	303	-	-
25		85-18	193	-	-
		85-19	42	-	-
30		85-20	123	1,649	446
		85-21	24	-	-
35		85-22	64	-	-
		85-23	43	-	-
40		85-24	32	-	-
		85-25	171	-	-
45		85-26	31	-	-
		85-27	36	-	-
50		85-28	38	-	-
		85-29	73	-	-
55		85-30	112	107	-
		85-31	83	-	-
60		85-32	43	-	-
		85-33	123	-	-
60		85-34	73	-	-
		85-35	53	-	-
60		85-36	103	-	-
		85-37	134	-	-
60		85-38	97	-	-
		85-39	147	-	-
60		85-40	119	62	-
		85-41	10,710	9,806	10,971
60		85-42	704	-	-
		85-43	575	-	-
60		85-44	78	-	-
		85-45	211	-	-
60		85-46	69	-	-
		85-47	49	-	-
60	46.0-50.0m: Saprolite B greenish gray colored saprolite, including chips of weathered intrusive	85-48	38	-	-
		85-49	30	-	-
60		85-50	39	37	-
		85-51	68	-	-
60	50.0-58.0m: Granodiorite greenish dark gray colored, medium grained granodiorite, chloritized, with pyrite dissemination, contents of pyrite is 1% in volume	85-52	55	-	-
		85-53	29	-	-
60		85-54	162	-	-
		85-55	501	-	-
60		85-56	1,533	1,783	2,366
		85-57	103	-	-
60	58.0-60.0m: Meta-andesite ? greenish gray colored meta-andesite, shistosed, with 1% pyrite dissemination	85-58	91	-	-
		85-59	79	-	-
60		85-60	60	72	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
4.0	0.0-4.0m: Alluvial sediments dark gray colored sediments, mostly clay material, including a lot of organic material	50-1	78	-	-
		50-2	57	-	-
		50-3	38	-	-
		50-4	0	-	-
6.0	4.0-6.0m: Mottled clay brown colored clay, including some lateritic nodules	50-5	23	-	-
		50-6	86	-	-
		50-7	21	-	-
		50-8	22	-	-
27.0	6.0-27.0m: Saprolite A yellowish colored, fine grained saprolite	50-9	25	-	-
		50-10	40	17	-
		50-11	54	-	-
		50-12	28	-	-
		50-13	27	-	-
		50-14	35	-	-
		50-15	52	-	-
		50-16	57	-	-
		50-17	44	-	-
		50-18	31	-	-
		50-19	35	-	-
		50-20	17	32	-
		50-21	51	-	-
		50-22	51	-	-
		50-23	40	-	-
		50-24	30	-	-
		50-25	27	-	-
		50-26	59	-	-
50-27	63	-	-		
37.0	27.0-37.0m: Saprolite B dark green to greenish light gray colored, chloritized saprolite, with strong pyrite dissemination (2-3% in volume)	50-28	20	-	-
		50-29	15	-	-
		50-30	26	569	60
		50-31	40	-	-
		50-32	28	-	-
		50-33	94	-	-
		50-34	250	-	-
		50-35	145	-	-
		50-36	223	-	-
		50-37	369	-	-
45.0	37.0-45.0m: Meta-sandstone greenish to gray colored, fine grained meta-sandstone, chloritized, with sulfide (pyrite-arsenopyrite) dissemination (1-2% in volume), including quartz-calcite veinlets, contents of sulfide increase in silicified zone	50-38	57	-	-
		50-39	28	-	-
		50-40	20	27	-
		50-41	8	-	-
		50-42	13	-	-
		50-43	22	-	-
		50-44	6	-	-
		50-45	129	-	-
50					
55					
60					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Carapace brown colored, with iron nodules	33-1	70	-	-
		33-2	252	-	-
5	2.0-16.0m: Saprolite B greenish grey colored saprolite, including a lot of chips of weathered peritic rock	33-3	57	-	-
		33-4	38	-	-
		33-5	88	-	-
		33-6	55	-	-
		33-7	59	-	-
		33-8	79	-	-
		33-9	46	-	-
		33-10	83	34	-
		33-11	43	-	-
		33-12	44	-	-
		33-13	48	-	-
		33-14	56	-	-
		33-15	38	-	-
		33-16	88	-	-
		16.0	16.0-60.0m: Meta-sandstone (muddy) dark grey colored, schistose meta-greywacke with disseminated pyrite, total amount of pyrite = 0 % to 2 % peritic schist ?	33-17	61
20	33-18	51		-	-
	33-19	27		-	-
25	33-20	37		23	-
	33-21	25		-	-
30	33-22	26		-	-
	33-23	26		-	-
35	33-24	26		-	-
	33-25	22		-	-
40	33-26	30		-	-
	33-27	46		-	-
45	33-28	22		-	-
	33-29	58		-	-
50	33-30	33		26	-
	33-31	40		-	-
55	33-32	38		-	-
	33-33	61		-	-
60	33-34	72		-	-
	33-35	25		-	-
60.0	33-36	37		-	-
	33-37	60		-	-
60.0	33-38	41		-	-
	33-39	25		-	-
60.0	33-40	24		37	-
	33-41	8		-	-
60.0	33-42	13		-	-
	33-43	10		-	-
60.0	33-44	25		-	-
	33-45	2		-	-
60.0	33-46	2		-	-
	33-47	0	-	-	
60.0	33-48	26	-	-	
	33-49	0	-	-	
60.0	33-50	35	58	-	
	33-51	34	-	-	
60.0	33-52	16	-	-	
	33-53	19	-	-	
60.0	33-54	24	-	-	
	33-55	23	-	-	
60.0	33-56	23	-	-	
	33-57	23	-	-	
60.0	33-58	32	-	-	
	33-59	18	-	-	
60.0	33-60	135	136	-	

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0 5 10 15.0	0.0-2.0m: Carapace reddish brown colored, with laterite nodules	34-1	99	-	-
		34-2	58	-	-
		34-3	124	-	-
	2.0-15.0m: Saprolite A greenish grey colored, soft and fine grained powder	34-4	253	-	-
		34-5	92	-	-
		34-6	355	-	-
		34-7	155	-	-
		34-8	279	-	-
		34-9	247	-	-
		34-10	157	141	-
		34-11	815	-	-
		34-12	741	-	-
		34-13	209	-	-
		34-14	294	-	-
		34-15	88	-	-
15.0 20 25 27.0	15.0-27.0m: Saprolite B greenish grey colored saprolite, including chips of weathered rock	34-16	208	-	-
		34-17	86	-	-
		34-18	46	-	-
		34-19	209	-	-
		34-20	116	119	-
		34-21	147	-	-
		34-22	174	-	-
		34-23	1,073	980	926
		34-24	186	-	-
		34-25	328	-	-
		34-26	833	-	-
		34-27	536	-	-
27.0 30 35 40 45 50 55 60.0	27.0-60.0m: Meta-sandstone dark grey to black colored meta-greywacke with disseminated pyrite, total amount of pyrite = 0 % to 1 % rock chip of 34-29 is meta-andesite with strong dissemination of pyrite	34-28	60	-	-
		34-29	41	-	-
		34-30	25	13	-
		34-31	13	-	-
		34-32	98	-	-
		34-33	20	-	-
		34-34	4	-	-
		34-35	22	-	-
		34-36	12	-	-
		34-37	6	-	-
		34-38	27	-	-
		34-39	111	-	-
		34-40	38	35	-
water 45 50 55 60	34-41	11	-	-	
	34-42	74	-	-	
	34-43	65	-	-	
	34-44	20	-	-	
	34-45	9	-	-	
	34-46	15	-	-	
	34-47	15	-	-	
	34-48	75	-	-	
	34-49	22	-	-	
	34-50	24	26	-	
	34-51	21	-	-	
	34-52	34	-	-	
34-53	35	-	-		
34-54	228	-	-		
34-55	190	-	-		
34-56	95	-	-		
34-57	30	-	-		
34-58	28	-	-		
34-59	25	-	-		
34-60	146	136	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 12.0	0.0-12.0m: Mottled clay brown colored clay, including quartz fragments	36-1	99	-	-
		36-2	100	-	-
		36-3	86	-	-
		36-4	38	-	-
		36-5	504	-	-
		36-6	172	-	-
		36-7	78	-	-
		36-8	67	-	-
		36-9	45	-	-
		36-10	36	42	-
		36-11	20	-	-
15 20 20.0	12.0-20.0m: Saprolite A greenish grey colored, fine grained powder	36-12	19	-	-
		36-13	44	-	-
		36-14	31	-	-
		36-15	27	-	-
		36-16	33	-	-
		36-17	70	-	-
		36-18	77	-	-
		36-19	10	-	-
		36-20	32	36	-
		36-21	53	-	-
25 30 31.0	20.0-31.0m: Saprolite B greenish grey colored saprolite, including chips of weathered granodiorite	36-22	30	-	-
		36-23	20	-	-
		36-24	20	-	-
		36-25	15	-	-
		36-26	54	-	-
		36-27	20	-	-
		36-28	33	-	-
		36-29	30	-	-
		36-30	13	40	-
		36-31	186	-	-
35 40 45 50 55 60	31.0-60.0m: Granodiorite light grey colored, medium to fine grained granodiorite or quartz diorite, weakly chloritized with pyrite (+ arsenopyrite ?) dissemination, 0% to 1% or more	36-32	29	-	-
		36-33	2,421	2,218	1,851
		36-34	37	-	-
		36-35	37	-	-
		36-36	42	-	-
		36-37	14	-	-
		36-38	32	-	-
		36-39	38	-	-
		36-40	54	58	-
		36-41	63	-	-
36-42	58	-	-		
36-43	54	-	-		
36-44	64	-	-		
36-45	74	-	-		
36-46	55	-	-		
36-47	52	-	-		
36-48	48	-	-		
36-49	40	-	-		
36-50	108	125	-		
36-51	59	-	-		
36-52	159	-	-		
36-53	152	-	-		
36-54	104	-	-		
36-55	49	-	-		
36-56	76	-	-		
36-57	117	-	-		
36-58	101	-	-		
36-59	81	-	-		
36-60	73	82	-		

KRC-40

Coordination: N1000 E320

Elevation: 340m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
4.0	0.0-4.0m: Surface soil greenish brown colored silty soil, including iron nodules	40-1	57	-	-
		40-2	125	-	-
		40-3	14	-	-
		40-4	118	-	-
6.0	4.0-6.0m: Carapace reddish brown colored hard, indurated soil with iron nodules	40-5	33	-	-
		40-6	47	-	-
		40-7	1,408	1,466	-
		40-8	151	-	-
8.0	6.0-8.0m: Mottled clay reddish brown colored hard, indurated soil with iron nodules	40-9	115	-	-
		40-10	35	2,201	130
		40-11	0	-	-
		40-12	336	-	-
10.0		40-13	449	-	-
		40-14	699	-	-
		40-15	477	-	-
		40-16	469	-	-
15.0	14.0-36.0m: Saprolite A brownish grey to greenish grey colored, fine grained powder	40-17	1,004	-	686
		40-18	161	-	-
		40-19	123	-	-
		40-20	102	107	-
20.0		40-21	314	-	-
		40-22	604	-	-
		40-23	418	-	-
		40-24	494	-	-
25.0		40-25	164	-	-
		40-26	176	-	-
		40-27	2,290	-	651
		40-28	1,192	-	1,406
30.0		40-29	172	-	-
		40-30	898	1,014	-
		40-31	686	-	-
		40-32	890	-	-
35.0		40-33	922	850	0
		40-34	1,376	-	-
		40-35	230	-	-
		40-36	210	-	-
40.0	36.0-44.0m: Saprolite B greenish grey colored, medium grained powder including diorite chips	40-37	224	-	-
		40-38	326	-	-
		40-39	174	-	-
		40-40	134	136	-
45.0		40-41	87	-	-
		40-42	1,073	262	377
		40-43	74	-	-
		40-44	996	1,016	660
50.0	44.0-60.0m: Diorite (or Granodiorite) dark grey colored, medium grained diorite or granodiorite with disseminated pyrite, amount of pyrite is 0 % to 1 %	40-45	193	-	-
		40-46	147	-	-
		40-47	531	-	-
		40-48	343	-	-
55.0		40-49	258	-	-
		40-50	112	99	-
		40-51	78	-	-
		40-52	90	-	-
60.0		40-53	160	-	-
		40-54	209	-	-
		40-55	241	-	-
		40-56	73	-	-
60.0		40-57	166	-	-
		40-58	206	-	-
		40-59	177	-	-
		40-60	401	318	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0 5 10 14.0	0.0-2.0m: Carapace brown to reddish brown colored carapace, including a lot of lateritic gravels 2.0-10.0m: Saprolite A greenish gray colored saprolite, fine grained, very soft, including quartz chips 10.0-14.0m: Saprolite B greenish gray colored saprolite, medium grained, including chips of weathered meta-graywacke	78-1	3	-	-
		78-2	23	-	-
		78-3	7	-	-
		78-4	6	-	-
		78-5	4	0	-
		78-6	4	-	-
		78-7	153	-	-
		78-8	3	-	-
		78-9	2	-	-
		78-10	514	-	-
15 20 25 30 35 40 45 50 55 60	14.0-20.0m: Meta-sandstone dark gray colored meta-graywacke, including rounded plagioclase fragment (diameter: 0.5-1 mm), with silicification with arsenopyrite inside quartz veinlet (18-19m) 20.0-30.0m: Peritic schist dark gray colored peritic schist, with pyrite dissemination (1-2%) 30.0-60.0m: Alternation beds alternation beds of peritic schist and meta- graywacke, with more sulfide in peritic schist (0-1%), with silicified fracture (41-46m)	78-11	27	-	-
		78-12	26	-	-
		78-13	82	-	-
		78-14	20	-	-
		78-15	3	0	-
		78-16	7	-	-
		78-17	12	-	-
		78-18	4	-	-
		78-19	30	-	-
		78-20	6	-	-
water		78-21	12	-	-
		78-22	9	-	-
		78-23	9	-	-
		78-24	20	-	-
		78-25	16	11	-
		78-26	8	-	-
		78-27	14	-	-
		78-28	10	-	-
		78-29	4	-	-
		78-30	36	-	-
		78-31	11	-	-
		78-32	9	-	-
		78-33	10	-	-
		78-34	12	-	-
		78-35	10	10	-
		78-36	8	-	-
		78-37	9	-	-
		78-38	5	-	-
		78-39	8	-	-
		78-40	15	-	-
		78-41	14	-	-
		78-42	19	-	-
		78-43	161	-	-
		78-44	16	-	-
		78-45	16	8	-
		78-46	8	-	-
		78-47	20	-	-
		78-48	12	-	-
		78-49	23	-	-
		78-50	20	-	-
		78-51	22	-	-
		78-52	33	-	-
		78-53	20	-	-
		78-54	13	-	-
		78-55	12	12	-
		78-56	28	-	-
		78-57	115	-	-
		78-58	46	-	-
		78-59	22	-	-
		78-60	19	-	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
0.0-3.0	0.0-3.0m: Carapace brownish red colored carapace, including a lot of lateritic nodules, this material seems to be a little bit indurated	79-1	5	-	-
		79-2	5	-	-
3.0-7.0	3.0-7.0m: Saprolite A yellow colored saprolite, fine grained, very soft	79-3	1	-	-
		79-4	<1	-	-
7.0-16.0	7.0-16.0m: Saprolite B greenish gray to greenish yellow colored saprolite, including chips of weathered meta-sediment	79-5	15	-	-
		79-6	29	-	-
16.0-18.0	16.0-18.0m: Meta-sandstone green colored schistosed rock, chloritized, meta-volcanics?	79-7	43	-	-
		79-8	10	-	-
18.0-22.0	18.0-22.0m: Meta-sandstone black colored meta-sandstone, coarse grained, massive	79-9	4	-	-
		79-10	5	6	-
22.0-24.0	22.0-24.0m: Peritic schist dark gray colored peritic schist, with no sulfide, schistosed	79-11	7	-	-
		79-12	25	-	-
24.0-44.0	24.0-44.0m: Meta-sandstone gray colored meta-graywacke, coarse grained, massive, not schistosed	79-13	18	-	-
		79-14	22	-	-
44.0-45.0	24.0-44.0m: Fe (OH) nodules? fall from upper part?	79-15	7	-	-
		79-16	5	-	-
45.0-46.0	water	79-17	26	-	-
		79-18	8	-	-
46.0-47.0		79-19	15	-	-
		79-20	7	13	-
47.0-48.0		79-21	20	-	-
		79-22	41	-	-
48.0-49.0		79-23	11	-	-
		79-24	18	-	-
49.0-50.0		79-25	75	-	-
		79-26	60	-	-
50.0-51.0		79-27	24	-	-
		79-28	23	-	-
51.0-52.0		79-29	9	-	-
		79-30	8	16	-
52.0-53.0		79-31	17	-	-
		79-32	21	-	-
53.0-54.0		79-33	20	-	-
		79-34	29	-	-
54.0-55.0		79-35	9	-	-
		79-36	10	-	-
55.0-56.0		79-37	6	-	-
		79-38	6	-	-
56.0-57.0		79-39	21	-	-
		79-40	29	23	-
57.0-58.0		79-41	28	-	-
		79-42	33	-	-
58.0-59.0		79-43	24	-	-
		79-44	29	-	-

KRC-80

Coordination: N1000 W600

Elevation: 340m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
4.0	0.0-4.0m: Carapace brown to gray colored carapace, silty, including a lot of lateritic nodules	80-1	72	-	-
		80-2	46	-	-
		80-3	136	-	-
		80-4	99	-	-
10.0	4.0-10.0m: Saprolite B greenish gray to brown colored saprolite, including chips of deeply weathered meta -sediment	80-5	29	-	-
		80-6	1,073	5	3
		80-7	16	-	-
		80-8	26	-	-
		80-9	28	-	-
		80-10	16	-	-
60.0	10.0-60.0m: Peritic schist black colored peritic schist, with pyrite dissemination along schistosity, estimated contents of pyrite is 2-5%, with minor quartz-calcite veinlets, including black material (graphite?)	80-11	56	-	-
		80-12	31	-	-
		80-13	13	-	-
		80-14	23	-	-
		80-15	14	-	-
		80-16	39	19	-
		80-17	18	-	-
		80-18	56	-	-
		80-19	14	-	-
		80-20	17	-	-
		80-21	14	-	-
		80-22	16	-	-
		80-23	111	-	-
		80-24	18	-	-
		80-25	13	-	-
		80-26	10	7	-
		80-27	10	-	-
		80-28	20	-	-
		80-29	12	-	-
		80-30	9	-	-
		80-31	18	-	-
		80-32	13	-	-
		80-33	16	-	-
		80-34	9	-	-
		80-35	5	-	-
		80-36	16	11	-
		80-37	16	-	-
		80-38	11	-	-
		80-39	14	-	-
		80-40	5	-	-
80-41	14	-	-		
80-42	42	-	-		
80-43	24	-	-		
80-44	18	-	-		
80-45	18	-	-		
80-46	15	15	-		
80-47	34	-	-		
80-48	40	-	-		
80-49	16	-	-		
80-50	13	-	-		
80-51	12	-	-		
80-52	11	-	-		
80-53	9	-	-		
80-54	11	-	-		
80-55	18	-	-		
80-56	17	17	-		
80-57	12	-	-		
80-58	25	-	-		
80-59	17	-	-		
80-60	14	-	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 13.0	0.0-4.0m: Carapace brown colored carapace, silty, clay material, including some lateritic nodules	81-1	78	-	-
		81-2	268	-	-
		81-3	69	-	-
		81-4	144	-	-
	4.0-13.0m: Saprolite B greenish yellow to greensih gray colored saprolite, including chips of weathered meta-sediment	81-5	31	-	-
		81-6	23	22	-
		81-7	29	-	-
		81-8	44	-	-
		81-9	16	-	-
		81-10	11	-	-
		81-11	19	-	-
		81-12	3	-	-
		81-13	14	-	-
15 20 25 30 35 40 45 50 52.0	13.0-52.0m: Muddy sandstone black colored meta-graywacke, fine grained, massive, pyrite dissemination is less than 1%	81-14	21	-	-
		81-15	30	-	-
		81-16	27	30	-
		81-17	115	-	-
		81-18	69	-	-
		81-19	59	-	-
		81-20	86	-	-
		81-21	26	-	-
		81-22	31	-	-
		81-23	24	-	-
		81-24	22	-	-
		81-25	50	-	-
		81-26	31	30	-
		81-27	33	-	-
		81-28	23	-	-
		81-29	21	-	-
		81-30	34	-	-
		81-31	24	-	-
		81-32	26	-	-
		81-33	34	-	-
		81-34	34	-	-
		81-35	24	-	-
		81-36	24	22	-
		81-37	10	-	-
		81-38	10	-	-
81-39	10	-	-		
81-40	13	-	-		
81-41	12	-	-		
81-42	52	-	-		
81-43	21	-	-		
81-44	15	-	-		
81-45	18	-	-		
81-46	36	61	-		
81-47	23	-	-		
81-48	29	-	-		
81-49	13	-	-		
81-50	15	-	-		
81-51	11	-	-		
81-52	13	-	-		
81-53	14	-	-		
81-54	51	-	-		
81-55	14	-	-		
81-56	257	19	-		
81-57	16	-	-		
81-58	31	-	-		
81-59	17	-	-		
81-60	14	-	-		

water

40.0-60.0: contents of sulfide increase (2-5%)

52.0-60.0m: Peritic schist
dark gray to black colored peritic schist,
with very fine grained pyrite dissemination,
contents of pyrite is 3%

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 9.0	0.0-9.0m: Crust, carapace reddish brown colored crust and carapace, including a lot of lateritic nodules	82-1	23	113	44
		82-2	8	32	-
		82-3	1	26	-
		82-4	10	85	54
		82-5	92	130	-
		82-6	146	179	-
		82-7	90	125	-
		82-8	0	163	114
		82-9	72	120	-
10 12.0	9.0-12.0m: Mottled clay brown colored clay, with some lateritic nodules	82-10	121	210	-
		82-11	140	188	-
		82-12	162	170	-
15 20 25 30 35 39.0	12.0-39.0m: Saprolite A pinkish brown to pinkish yellow colored, fine grained saprolite, very soft, with no rock chips	82-13	194	179	-
		82-14	182	215	-
		82-15	253	291	-
		82-16	311	290	-
		82-17	494	571	-
		82-18	466	552	-
		82-19	770	820	-
		82-20	698	787	-
		82-21	173	-	-
		82-22	127	-	-
		82-23	34	-	-
		82-24	179	-	-
		82-25	324	-	-
		82-26	726	-	-
		82-27	120	-	-
		82-28	479	-	-
		82-29	593	-	-
		82-30	352	389	-
82-31	74	-	-		
82-32	123	-	-		
82-33	121	-	-		
82-34	173	-	-		
82-35	133	-	-		
82-36	63	-	-		
82-37	170	-	-		
82-38	83	-	-		
82-39	178	-	-		
40 45 50 55 60.0	52.0-60.0m: Granodiorite gray colored, hornblende-biotite granodiorite, medium grained, fresh, with pyrite dissemination, contents of sulfide is 0-1% in volume	82-40	41	111	-
		82-41	124	-	-
		82-42	142	-	-
		82-43	242	-	-
		82-44	249	-	-
		82-45	3	-	-
		82-46	146	-	-
		82-47	47	-	-
		82-48	71	-	-
		82-49	559	-	-
		82-50	77	54	-
		82-51	51	-	-
82-52	137	-	-		
82-53	54	-	-		
82-54	54	-	-		
82-55	79	-	-		
82-56	260	-	-		
82-57	84	-	-		
82-58	101	-	-		
82-59	80	-	-		
82-60	29	30	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-9.0m: Crust, carapace reddish brown colored crust and carapace, including a lot of lateritic nodules	83-1	29	-	-
		83-2	20	-	-
		83-3	187	-	-
		83-4	25	-	-
		83-5	33	-	-
		83-6	424	-	-
		83-7	117	-	-
		83-8	134	-	-
		83-9	116	-	-
9.0	9.0-12.0m: Mottled clay brown colored clay, including iron oxide, with some lateritic nodules	83-10	129	113	-
83-11		112	-	-	
83-12		167	-	-	
12.0	12.0-39.0m: Saprolite A brown to pinkish brown colored fine grained saprolite, very soft, with no rock chips	83-13	206	-	-
83-14		118	-	-	
83-15		31	-	-	
83-16		23	-	-	
83-17		79	-	-	
83-18		44	-	-	
83-19		176	-	-	
83-20		341	46	-	
83-21		39	-	-	
83-22		65	-	-	
83-23		20	-	-	
83-24		324	-	-	
83-25		265	-	-	
83-26		57	-	-	
83-27		310	-	-	
83-28		141	-	-	
83-29		214	-	-	
83-30		68	62	-	
83-31		2,253	20	0	
83-32		45	-	-	
83-33		33	-	-	
83-34	19	-	-		
83-35	33	-	-		
83-36	69	-	-		
83-37	39	-	-		
83-38	52	-	-		
83-39	468	-	-		
39.0	52.0-60.0m: Saprolite B greensih gray colored saprolite, including chips of weathered granodiorite, with minor chlorite	83-40	111	56	-
83-41		122	-	-	
83-42		39	-	-	
83-43		25	-	-	
83-44		28	-	-	
83-45		10	-	-	
83-46		19	-	-	
83-47		229	-	-	
83-48		31	-	-	
83-49		22	-	-	
83-50		51	34	-	
83-51	46	-	-		
83-52	59	-	-		
83-53	63	-	-		
83-54	52	-	-		
83-55	41	-	-		
83-56	65	-	-		
56.0	56.0-60.0m: Silicified zone light gray colored silicified rock, with strongly disseminated of pyrite>arsenopyrite >pyrrhotite	83-57	91	-	-
83-58		176	-	-	
83-59		307	-	-	
83-60		188	162	-	
60.0					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 5.0	0.0-9.0m: Alluvial sediments brown colored sediments, including lateritic nodules 0.0-2.0m: dark brown colored sediments, with organic material	41-1	162	-	-
		41-2	91	-	-
		41-3	63	-	-
		41-4	29	-	-
		41-5	74	-	-
10 14.0	5.0-14.0m: Mottled clay brown colored, fine grained clay, soft, transition to saprolite is gradual	41-6	43	-	-
		41-7	137	-	-
		41-8	86	-	-
		41-9	22	-	-
		41-10	20	181	-
		41-11	48	-	-
		41-12	20	-	-
		41-13	29	-	-
		41-14	298	-	-
		41-15	194	-	-
15 20 25 30 35	14.0-35.0m: Saprolite A yellowish gray to greenish gray colored, fine grained saprolite, soft, including no rock chips	41-16	55	-	-
		41-17	36	-	-
		41-18	57	-	-
		41-19	13	-	-
		41-20	85	38	-
		41-21	26	-	-
		41-22	30	-	-
		41-23	7	-	-
		41-24	0	-	-
		41-25	8	-	-
		41-26	0	-	-
		41-27	0	-	-
		41-28	19	-	-
		41-29	4	-	-
		41-30	73	26	-
35 40 40.0	35.0-40.0m: Saprolite B greensih gray colored saprolite, including some chips of deeply weathered rock	41-31	0	-	-
		41-32	706	-	-
		41-33	0	-	-
		41-34	6	-	-
		41-35	14	-	-
		41-36	0	-	-
		41-37	60	-	-
		41-38	23	-	-
		41-39	27	-	-
		41-40	0	555	-
45					
50					
55					
60					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0 5 10 15 20 20.0	0.0-2.0m: Carapace brown colored, with iron nodules	25-1	233	-	-
		25-2	82	-	-
	2.0-20.0m: Saprolite B greenish grey colored saprolite, including a lot of chips of weathered rock	25-3	72	-	-
		25-4	24	-	-
		25-5	14	-	-
		25-6	29	-	-
		25-7	38	-	-
		25-8	28	-	-
		25-9	34	-	-
		25-10	18	12	-
		25-11	14	-	-
		25-12	30	-	-
		25-13	33	-	-
		25-14	21	-	-
		25-15	38	-	-
		25-16	107	-	-
		25-17	30	-	-
		25-18	21	-	-
		25-19	12	-	-
		25-20	44	26	-
25 30 35 40 45 50 55 60 60.0	16.0-60.0m: Meta-sandstone (greywacke) greenish dark grey colored, meta-greywacke with disseminated pyrite, total amount of pyrite = 0 % to 2 % epidotic alteration is observed, meta-volcanics	25-21	12	-	-
		25-22	9	-	-
		25-23	12	-	-
		25-24	22	-	-
		25-25	16	-	-
		25-26	25	-	-
		25-27	7	-	-
		25-28	2	-	-
		25-29	3	-	-
		25-30	39	18	-
		25-31	9	-	-
		25-32	47	-	-
		25-33	15	-	-
		25-34	16	-	-
		25-35	8	-	-
		25-36	30	-	-
		25-37	4	-	-
		25-38	1	-	-
		25-39	35	-	-
		25-40	107	22	-
		25-41	54	-	-
		25-42	91	-	-
		25-43	33	-	-
		25-44	59	-	-
		25-45	19	-	-
		25-46	43	-	-
		25-47	34	-	-
		25-48	45	-	-
		25-49	92	-	-
		25-50	52	93	-
		25-51	72	-	-
		25-52	91	-	-
		25-53	41	-	-
		25-54	49	-	-
		25-55	53	-	-
		25-56	54	-	-
		25-57	64	-	-
		25-58	32	-	-
		25-59	40	-	-
		25-60	33	48	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0 5 10 15 20 22.0	0.0-1.0m: Carapace brown colored carapace with laterite nodules	26-1	50	-	-
		26-2	109	-	-
	1.0-22.0m: Saprolite B greenish grey colored saprolite including chips weathered rock	26-3	32	-	-
		26-4	688	-	-
		26-5	56	-	-
		26-6	45	-	-
		26-7	18	-	-
		26-8	19	-	-
		26-9	47	-	-
		26-10	54	59	-
		26-11	30	-	-
		26-12	11	-	-
		26-13	41	-	-
		26-14	39	-	-
		26-15	44	-	-
		26-16	66	-	-
		26-17	26	-	-
		26-18	64	-	-
		26-19	62	-	-
		26-20	218	235	-
	26-21	17	-	-	
	26-22	142	-	-	
22.0 25 30 35 40 45 50 55 60 60.0	22.0-60.0m: Meta-sandstone dark grey to greenish grey colored meta-sandstone, with disseminated pyrite, total amount of pyrite = 0 % to 1 % green color shows chloritic and / or epidotic alteration, meta-volcanics ? with traces of calcite films	26-23	134	-	-
		26-24	12	-	-
		26-25	8	-	-
		26-26	21	-	-
		26-27	31	-	-
		26-28	84	-	-
		26-29	13	-	-
		26-30	87	81	-
		26-31	31	-	-
		26-32	14	-	-
		26-33	7	-	-
		26-34	33	-	-
		26-35	38	-	-
		26-36	155	-	-
		26-37	123	-	-
		26-38	831	-	-
		26-39	107	-	-
		26-40	48	22	-
		26-41	1,189	102	686
		26-42	68	-	-
		26-43	145	-	-
		26-44	3,734	3,780	3,463
		26-45	23	-	-
		26-46	1,776	142	34
		26-47	171	-	-
		26-48	67	-	-
		26-49	1,514	1,110	1,234
		26-50	276	235	-
		26-51	834	-	-
		26-52	1,022	1,150	789
		26-53	277	-	-
		26-54	59	-	-
		26-55	495	-	-
		26-56	78	-	-
		26-57	47	29	22
		26-58	9,670	100	150
		26-59	308	400	410
		26-60	115	108	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0	0.0-1.0m: Carapace	27-1	49	-	-
3.0	brown colored soil with laterite nodules	27-2	364	-	-
		27-3	362	-	-
	2.0-3.0m: Mottled clay	27-4	81	-	-
5	brown to greenish grey colored clay	27-5	21	-	-
		27-6	34	-	-
	3.0-22.0m: Saprolite B	27-7	22	-	-
	greenish grey colored saprolite, including chips of wethered rock	27-8	2	-	-
		27-9	28	-	-
10		27-10	131	132	-
		27-11	24	-	-
		27-12	49	-	-
		27-13	179	-	-
		27-14	58	-	-
15		27-15	22	-	-
		27-16	136	-	-
		27-17	56	-	-
		27-18	56	-	-
20		27-19	120	-	-
		27-20	1,403	1,718	-
		27-21	244	-	-
		27-22	146	-	-
22.0	22.0-60.0m: Meta-sandstone	27-23	30	-	-
	dark grey to greenish grey colored meta-sandstone, with disseminated pyrite, total amount of pyrite = 0 % to 1 %	27-24	30	-	-
25	green color shows chloritic and / or epidotic alteration, meta-volcanics ?	27-25	115	-	-
		27-26	42	-	-
		27-27	58	-	-
		27-28	508	-	-
		27-29	93	-	-
30		27-30	131	206	-
		27-31	44	-	-
		27-32	102	-	-
		27-33	931	-	-
		27-34	92	-	-
35		27-35	153	-	-
		27-36	482	-	-
		27-37	39	-	-
		27-38	220	-	-
		27-39	553	-	-
40		27-40	77	40	-
		27-41	58	-	-
		27-42	418	-	-
		27-43	154	-	-
		27-44	36	-	-
45		27-45	0	-	-
		27-46	12	-	-
		27-47	7	-	-
		27-48	492	-	-
		27-49	289	-	-
50		27-50	1,151	158	124
		27-51	237	-	-
		27-52	2,835	2,668	2,263
		27-53	91	-	-
		27-54	18	-	-
55		27-55	67	-	-
		27-56	20	-	-
		27-57	12	-	-
		27-58	8	-	-
		27-59	2	-	-
60		27-60	0	2	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Surface soil grey colored soil, including quartz fragments	28-1	49	-	-
		28-2	10	-	-
5	2.0-10.0m: Saprolite A brown to greenish grey colored, fine grained powder	28-3	81	55	-
		28-4	16	-	-
		28-5	107	-	-
		28-6	27	-	-
		28-7	7	-	-
		28-8	9	-	-
		28-9	8	-	-
		28-10	10	-	-
10	10.0-25.0m: Saprolite B greenish grey colored saprolite, including chips of weathered rock	28-11	11	-	-
		28-12	8	-	-
		28-13	8	5	-
		28-14	8	-	-
		28-15	8	-	-
		28-16	22	-	-
		28-17	8	-	-
		28-18	164	-	-
		28-19	28	-	-
		28-20	15	-	-
20		28-21	111	-	-
		28-22	18	-	-
		28-23	22	10	-
		28-24	54	-	-
		28-25	6	-	-
		28-26	19	-	-
		28-27	1	-	-
		28-28	11	-	-
		28-29	43	-	-
		28-30	84	-	-
30	dark grey to greenish grey colored meta-andesite ?, with disseminated pyrite, total amount of pyrite = 0 % to 1 % green color shows chloritic and / or epidotic alteration meta-greywacke ?	28-31	9	-	-
		28-32	6	-	-
		28-33	27	32	-
		28-34	56	-	-
		28-35	105	-	-
		28-36	43	-	-
		28-37	5	-	-
		28-38	5	-	-
		28-39	6	-	-
		28-40	168	-	-
40		28-41	261	-	-
		28-42	88	-	-
		28-43	0	7	-
		28-44	59	-	-
		28-45	116	-	-
		28-46	14	-	-
		28-47	6	-	-
		28-48	586	-	-
		28-49	14	-	-
		28-50	6	-	-
50		28-51	8	-	-
		28-52	3	-	-
		28-53	15	9	-
		28-54	19	-	-
		28-55	6	-	-
		28-56	10	-	-
		28-57	8	-	-
		28-58	1	-	-
		28-59	7	-	-
		28-60	1,391	1,460	0
60	60.0				

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Surface soil grey colored soil, including organic matter	29-1	91	-	-
		29-2	70	-	-
5	2.0-7.0m: Mottled clay brown colored clay and white clay, with oxide minerals	29-3	58	88	-
		29-4	102	-	-
		29-5	261	-	-
		29-6	65	-	-
		29-7	306	-	-
		29-8	33	-	-
		29-9	16	-	-
10	7.0-15.0m: Saprolite A greenish grey to brown colored, fine grained powder	29-10	11	-	-
		29-11	26	-	-
		29-12	20	-	-
		29-13	75	32	-
		29-14	46	-	-
		29-15	22	-	-
		29-16	14	-	-
		29-17	8	-	-
		29-18	7	-	-
		29-19	52	-	-
		29-20	10	-	-
		29-21	14	-	-
		29-22	17	-	-
15	15.0-22.0m: Saprolite B greenish grey to light grey colored saprolite including chips of granodiorite	29-23	15	13	-
		29-24	9	-	-
		29-25	3	-	-
		29-26	8	-	-
		29-27	13	-	-
		29-28	38	-	-
		29-29	7	-	-
		29-30	51	-	-
		29-31	8	-	-
		29-32	25	-	-
		29-33	4	8	-
		29-34	6	-	-
		29-35	8	-	-
		29-36	31	-	-
		29-37	9	-	-
		29-38	17	-	-
		29-39	7	-	-
		29-40	70	-	-
		29-41	123	-	-
		29-42	15	-	-
		29-43	541	591	-
		29-44	3	-	-
29-45	0	-	-		
20	22.0-45.0m: Fine grained granodiorite grey to light grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite	29-46	121	-	-
		29-47	81	-	-
		29-48	84	-	-
		29-49	286	-	-
		29-50	256	-	-
		29-51	161	-	-
		29-52	118	-	-
		29-53	31	27	-
		29-54	17	-	-
		29-55	1,241	1,490	1,474
		29-56	195	-	-
29-57	48	-	-		
29-58	46	-	-		
29-59	749	-	-		
29-60	231	-	-		
25	45.0-52.0m: Diorite dark grey colored, fine grained diorite or quartz diorite, including small amount of pyrite				
30	52.0-60.0m: Fine grained granodiorite grey to light grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite				
35					
40					
45					
50					
55					
60					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Surface soil grey to brown colored soil, including silty material	30-1	84	-	-
		30-2	82	-	-
4.0	2.0-4.0m: Carapace brown colored, Fe-oxide rich zone	30-3	55	34	-
		30-4	37	-	-
7.0	4.0-7.0m: Carapace brown colored, fine grained soil	30-5	30	-	-
		30-6	35	-	-
10.0	7.0-10.0m: Mottled clay brown to greenish brown, some times reddish brown colored clay	30-7	162	-	-
		30-8	23	-	-
15.0	10.0-22.0m: Saprolite A greenish grey to light grey colored fine grained powder	30-9	41	-	-
		30-10	43	-	-
20.0	22.0-37.0m: Saprolite B greenish grey to light grey colored fine grained powder, including chips of weathered granodiorite	30-11	65	-	-
		30-12	118	-	-
25.0	37.0-60.0m: Fine grained granodiorite grey to light grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1 % including pyrrhotite ?	30-13	383	281	-
		30-14	173	-	-
30.0		30-15	2,103	2,173	2,023
		30-16	71	-	-
35.0		30-17	101	-	-
		30-18	30	-	-
40.0		30-19	84	-	-
		30-20	196	-	-
45.0		30-21	75	-	-
		30-22	36	-	-
50.0		30-23	96	120	-
		30-24	66	-	-
55.0		30-25	77	-	-
		30-26	587	-	-
60.0		30-27	174	-	-
		30-28	55	-	-
		30-29	287	-	-
		30-30	74	-	-
		30-31	27	-	-
		30-32	62	-	-
		30-33	117	174	-
		30-34	52	-	-
		30-35	60	-	-
		30-36	31	-	-
		30-37	43	-	-
		30-38	15	-	-
		30-39	25	-	-
		30-40	14	-	-
		30-41	7	-	-
		30-42	101	-	-
		30-43	53	58	-
		30-44	108	-	-
		30-45	22	-	-
		30-46	91	-	-
		30-47	50	-	-
		30-48	56	-	-
		30-49	63	-	-
		30-50	22	-	-
		30-51	16	-	-
		30-52	45	-	-
		30-53	327	349	-
		30-54	15	-	-
		30-55	28	-	-
		30-56	18	-	-
		30-57	16	-	-
		30-58	45	-	-
		30-59	81	-	-
		30-60	29	-	-

Column (m)	Description	Sample No.	Au			
			ppb	ppb	ppb	
5	0.0-2.0m: Surface soil grey to brown colored soil, including silty material	31-1	137	-	-	
		31-2	124	-	-	
	2.0-4.0m: Carapace brown colored, Fe-oxide rich zone	31-3	373	385	-	
		31-4	112	-	-	
	4.0-7.0m: Carapace brown colored, fine grained soil	31-5	29	-	-	
		31-6	27	-	-	
	7.0	7.0-10.0m: Mottled clay brown to greenish brown, some times reddish brown colored clay	31-7	23	-	-
			31-8	27	-	-
	10	10.0-22.0m: Saprolite A greenish grey to light grey colored fine grained powder	31-9	31	-	-
			31-10	20	-	-
31-11			53	-	-	
31-12			15	-	-	
31-13			25	25	-	
31-14			7	-	-	
31-15			20	-	-	
31-16			1,905	1,035	1,186	
31-17			2,324	947	2,369	
31-18			48	-	-	
15	22.0-37.0m: Saprolite B greenish grey to light grey colored fine grained powder, including chips of weathered granodiorite	31-19	713	-	-	
		31-20	377	-	-	
		31-21	54	-	-	
		31-22	100	-	-	
		31-23	46	40	-	
		31-24	33	-	-	
		31-25	309	-	-	
		31-26	64	-	-	
		31-27	23	-	-	
		31-28	481	-	-	
20	37.0-51.0m: Fine grained granodiorite grey to light grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1 % including pyrrhotite ?	31-29	60	-	-	
		31-30	41	-	-	
		31-31	15	-	-	
		31-32	312	-	-	
		31-33	85	50	-	
		31-34	88	-	-	
		31-35	19	-	-	
		31-36	4	-	-	
		31-37	7	-	-	
		31-38	37	-	-	
25	37.0-51.0m: Fault ? brown colored soil, with no rock chips	31-39	7	-	-	
		31-40	260	-	-	
		31-41	175	-	-	
		31-42	118	-	-	
		31-43	9	927	24	
		31-44	748	-	-	
		31-45	116	-	-	
		31-46	86	-	-	
		31-47	41	-	-	
		31-48	89	-	-	
30		31-49	134	-	-	
		31-50	33	-	-	
		31-51	188	-	-	
		31-52	49	-	-	
		31-53	124	125	-	
		31-54	77	-	-	
		31-55	59	-	-	
		31-56	53	-	-	
		31-57	118	-	-	
		31-58	94	-	-	
35		31-59	25	-	-	
		31-60	34	-	-	
40						
45						
50						
55						
60						

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Carapace reddish brown colored soil, including iron nodules	32-1	18	-	-
		32-2	31	-	-
5	2.0-8.0m: Mottled clay brown to yellowish brown colored clay, including limonite ?	32-3	27	33	-
		32-4	25	-	-
		32-5	12	-	-
		32-6	10	-	-
		32-7	13	-	-
		32-8	19	-	-
		32-9	11	-	-
10	8.0-22.0m: Saprolite A greenish grey colored fine grained saprolite	32-10	430	-	-
		32-11	693	-	-
		32-12	31	-	-
		32-13	21	27	-
		32-14	29	-	-
		32-15	15	-	-
		32-16	22	-	-
		32-17	11	-	-
		32-18	14	-	-
		32-19	15	-	-
		32-20	9	-	-
		32-21	10	-	-
		32-22	26	-	-
15	22.0-31.0m: Saprolite B greenish grey colored, fine grained saprolite including chips of weathered granodiorite	32-23	41	10	-
		32-24	19	-	-
		32-25	15	-	-
		32-26	110	-	-
		32-27	47	-	-
		32-28	8	-	-
		32-29	4	-	-
		32-30	4	-	-
		32-31	11	-	-
		32-32	4	-	-
20	31.0-39.0m: Fine grained granodiorite chloritized greenish grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1%	32-33	66	31	-
		32-34	11	-	-
		32-35	7	-	-
		32-36	42	-	-
		32-37	24	-	-
		32-38	21	-	-
		32-39	13	-	-
		32-40	23	-	-
25	39.0-42.0m: Fault ? small amount of rock chips of sericitized granodiorite	32-41	50	-	-
		32-42	25	-	-
		32-43	40	39	-
30	42.0-60.0m: Fine grained granodiorite chloritized greenish grey colored, fine grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1%	32-44	9	-	-
		32-45	91	-	-
		32-46	107	-	-
		32-47	59	-	-
		32-48	14	-	-
		32-49	7	-	-
		32-50	5	-	-
		32-51	5	-	-
		32-52	5	-	-
		32-53	5	4	-
		32-54	3	-	-
		32-55	23	-	-
		32-56	28	-	-
35		32-57	14	-	-
		32-58	7	-	-
		32-59	3	-	-
		32-60	3	-	-
40					
45					
50					
55					
60	60.0				

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 13.0	0.0-13.0m: Carapace reddish brown colored carapace, with a lot of iron nodules (diameter: 3-5mm), coarse grained sand	76-1	4	-	-
		76-2	6	-	-
		76-3	4	-	-
		76-4	18	-	-
		76-5	42	-	-
		76-6	32	-	-
		76-7	48	-	-
		76-8	74	-	-
		76-9	66	-	-
		76-10	68	90	-
		76-11	84	-	-
		76-12	124	-	-
		15 20 25 30 34.0	13.0-34.0m: Saprolite A reddish brown to yellowish brown colored, very fine grained saprolite, without rock chips	76-13	168
76-14	876			1,054	-
76-15	1,840			1,440	1,782
76-16	402			-	-
76-17	454			-	-
76-18	64			-	-
76-19	254			-	-
76-20	78			57	-
76-21	111			-	-
76-22	97			-	-
76-23	15			-	-
76-24	57			-	-
76-25	44			-	-
76-26	18			-	-
76-27	32			-	-
76-28	31			-	-
76-29	19			-	-
35 40 45 49.0	34.0-49.0m: Saprolite A light gray to greenish light gray colored, saprolite, medium to fine grained, without rock chips	76-30	26	21	-
		76-31	50	-	-
		76-32	29	-	-
		76-33	11	-	-
		76-34	14	-	-
		76-35	11	-	-
		76-36	12	-	-
		76-37	13	-	-
		76-38	13	-	-
		76-39	6	-	-
		76-40	5	13	-
		76-41	13	-	-
		76-42	5	-	-
		76-43	2	-	-
50 52.0	49.0-52.0m: Saprolite B light gray colored, coarse grained saprolite, including a lot of rockchips (diameter: 5-7mm)	76-44	0	-	-
		76-45	38	-	-
		76-46	35	-	-
		76-47	77	-	-
		76-48	237	-	-
		76-49	59	-	-
		76-50	14	125	-
		76-51	9	-	-
		76-52	5	-	-
		76-53	46	-	-
55 60.0	52.0-60.0m: Volcanic sandstone black colored volcanic sandstone, weakly schistosed, including a lot of plagioclase grains (subhedral, 1-2mm, rounded), Fe-oxides stain along fractures	76-54	165	-	-
		76-55	42	-	-
		76-56	21	-	-
		76-57	243	-	-
		76-58	19	-	-
		76-59	407	-	-
		76-60	11	7	-

KRC-77

Coordination: N1250 E350

Elevation: 370m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 15 16.0	0.0-16.0m: Carapace reddish brown colored carapace, coarse grained, including a lot of iron nodules (diameter: 0.3-0.8cm)	77-1	20	-	-
		77-2	352	-	-
		77-3	56	-	-
		77-4	36	-	-
		77-5	42	-	-
		77-6	36	-	-
		77-7	20	-	-
		77-8	34	-	-
		77-9	48	-	-
		77-10	28	21	-
		77-11	58	-	-
		77-12	76	-	-
		77-13	52	-	-
		77-14	56	-	-
		77-15	84	-	-
		16.0 20 25 30 35 38.0	16.0-38.0m: Saprolite A reddish brown to brown colored saprolite, very fine grained saprolite (silty), without rock chips	77-16	76
77-17	44			-	-
77-18	54			-	-
77-19	32			-	-
77-20	144			143	-
77-21	25			-	-
77-22	36			-	-
77-23	34			-	-
77-24	35			-	-
77-25	26			-	-
77-26	19			-	-
77-27	31			-	-
77-28	21			-	-
77-29	21			-	-
77-30	26			25	-
77-31	20			-	-
77-32	20			-	-
77-33	12			-	-
77-34	15			-	-
77-35	20			-	-
77-36	17			-	-
77-37	10			-	-
77-38	15			-	-
38.0 40 45 50 55 60	38.0-60.0m: Saprolite A brown to yellowish light gray colored saprolite, fine grained, without rock chips			77-39	13
		77-40	16	17	-
		77-41	7	-	-
		77-42	18	-	-
		77-43	23	-	-
		77-44	18	-	-
		77-45	20	-	-
		77-46	32	-	-
		77-47	13	-	-
		77-48	15	-	-
		77-49	16	-	-
		77-50	16	26	-
		77-51	16	-	-
		77-52	13	-	-
		77-53	22	-	-
		77-54	10	-	-
		77-55	13	-	-
		77-56	13	-	-
		77-57	10	-	-
		77-58	20	-	-
		77-59	13	-	-
		77-60	5	10	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-5.0m: Carapace reddish brown colored carapace, with lateritic nodules	67-1	173	-	-
		67-2	29	-	-
		67-3	55	-	-
		67-4	30	-	-
		67-5	14	-	-
10	5.0-18.0m: Saprolite A yellowish brown colored saprolite, fine grained	67-6	6	-	-
		67-7	0	-	-
		67-8	2,970	49	0
		67-9	32	-	-
		67-10	0	21	-
		67-11	62	-	-
		67-12	33	-	-
		67-13	29	-	-
		67-14	43	-	-
		67-15	0	-	-
		67-16	14	-	-
		67-17	35	-	-
		20	18.0-37.0m: Saprolite A yellowish gray to brownish gray colored saprolite, fine grained, including no rock chips	67-18	21
67-19	29			-	-
67-20	28			0	-
67-21	166			-	-
67-22	52			-	-
67-23	32			-	-
67-24	32			-	-
67-25	52			-	-
67-26	22			-	-
67-27	130			-	-
67-28	20			-	-
67-29	22			-	-
67-30	28			28	-
67-31	18			-	-
67-32	20			-	-
67-33	32			-	-
67-34	26			-	-
67-35	10			-	-
67-36	14			-	-
30	37.0-47.0m: Saprolite A gray to yellowish gray colored saprolite, fine grained, including no rock chips			67-37	20
		67-38	24	-	-
		67-39	8	-	-
		67-40	20	14	-
		67-41	30	-	-
		67-42	23	-	-
		67-43	0	-	-
		67-44	7	-	-
		67-45	130	-	-
		67-46	19	-	-
		67-47	0	-	-
		40	47.0-51.0m: Saprolite B gray colored saprolite, including small amount of rock fragments	67-48	20
67-49	37			-	-
67-50	33			26	-
67-51	0			-	-
67-52	36			-	-
50	51.0-60.0m: Peritic schist dark gray colored peritic schist, with schistosity filling iron oxide and pyrite, fracture rich, including rock fragment (diameter: 2-5mm)	67-53	15	-	-
		67-54	10	-	-
		67-55	20	-	-
		67-56	16	-	-
		67-57	52	-	-
		67-58	13	-	-
		67-59	22	-	-
		67-60	9	38	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-9.0m: Carapace reddish brown colored carapace, including lateritic gravels, with iron oxides	68-1	8	-	-
		68-2	7	-	-
		68-3	11	-	-
		68-4	9	-	-
		68-5	16	-	-
		68-6	174	-	-
		68-7	10	-	-
		68-8	9	-	-
		68-9	24	-	-
		9.0			
10	9.0-29.0m: Saprolite A greenish gray colored saprolite, fine grained and very soft saprolite	68-10	9	11	-
		68-11	8	-	-
		68-12	7	-	-
		68-13	13	-	-
		68-14	18	-	-
		68-15	5	-	-
		68-16	5	-	-
		68-17	4	-	-
		68-18	6	-	-
		68-19	8	-	-
		68-20	17	17	-
		68-21	26	-	-
		68-22	11	-	-
		68-23	6	-	-
		68-24	19	-	-
		68-25	32	-	-
		68-26	63	-	-
		68-27	25	-	-
		68-28	0	-	-
29.0					
30	29.0-39.0m: Saprolite B greenish gray colored saprolite, including chips of deeply weathered rock, with many quartz vein chips	68-29	27	-	-
		68-30	56	189	85
		68-31	66	-	-
		68-32	53	-	-
		68-33	40	-	-
		68-34	36	-	-
		68-35	19	-	-
		68-36	21	-	-
		68-37	37	-	-
		68-38	20	-	-
		68-39	10	-	-
39.0					
40	39.0-60.0m: Peritic schist gray to dark gray colored peritic schist, fine grained and schistosed, rock chips clearly show schistosity, sulfides are mainly included in schistosity and fracture planes	68-40	13	11	-
		68-41	3	-	-
		68-42	3	-	-
		68-43	3	-	-
		68-44	15	-	-
		68-45	5	-	-
		68-46	6	-	-
		68-47	5	-	-
		68-48	4	-	-
		68-49	5	-	-
		68-50	7	1	-
68-51	2	-	-		
68-52	5	-	-		
68-53	5	-	-		
68-54	3	-	-		
68-55	miss	miss	miss		
68-56	2	-	-		
68-57	5	-	-		
68-58	45	-	-		
68-59	0	-	-		
68-60	2	-	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
3.0	0.0-3.0m: Surface soil dark brown colored soil, including lateritic gravels	70-1	87	-	-
		70-2	336	-	-
		70-3	69	-	-
5.0	3.0-5.0m: Carapace red brown colored carapace, including lateritic gravels	70-4	58	-	-
		70-5	40	-	-
		70-6	57	-	-
12.0	5.0-12.0m: Mottled clay brown to yellowish brown colored mottled clay, soft	70-7	4,262	1,798	3,531
		70-8	1,495	1,156	1,646
		70-9	1,732	1,374	1,646
		70-10	849	423	560
		70-11	217	-	-
		70-12	69	-	-
16.0	12.0-16.0m: Saprolite A greenish gray colored saprolite, fine grained, soft	70-13	161	-	-
		70-14	108	-	-
		70-15	59	-	-
21.0	16.0-21.0m: Saprolite B greenish gray to gray colored saprolite, including chips of weathered meta-sandstone	70-16	44	-	-
		70-17	81	-	-
		70-18	269	-	-
		70-19	84	-	-
		70-20	41	42	-
		70-21	41	-	-
21.0-60.0m	21.0-60.0m: Meta-sandstone gray to dark gray colored meta-sandstone, fine grained and hard, with pyrite-arsenopyrite dissemination (30-31, 40-41m), including quartz vein chips, total amount of sulfide=0-2%, and also sulfide in small veinlets	70-22	54	-	-
		70-23	30	-	-
		70-24	97	-	-
		70-25	390	-	-
		70-26	282	-	-
		70-27	104	-	-
		70-28	181	-	-
		70-29	102	-	-
		70-30	193	190	-
		70-31	765	-	-
		70-32	1,214	1,151	1,234
		70-33	1,217	1,328	1,749
		70-34	5,865	7,050	6,271
		70-35	4,245	5,425	5,486
		70-36	2,835	2,900	2,983
		70-37	4,513	5,694	5,040
		70-38	4,865	5,244	5,863
		70-39	1,323	-	1,200
70-40	129	123	-		
70-41	94	-	-		
70-42	116	-	-		
70-43	86	-	-		
70-44	165	-	-		
70-45	329	-	-		
70-46	146	-	-		
70-47	60	-	-		
70-48	143	-	-		
70-49	34	-	-		
70-50	34	32	-		
70-51	44	-	-		
70-52	80	-	-		
70-53	76	-	-		
70-54	40	-	-		
70-55	71	-	-		
70-56	44	-	-		
70-57	29	-	-		
70-58	20	-	-		
70-59	10	-	-		
70-60	5	7	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Carapace brown colored soil, including iron nodules	18-1	218	-	-
		18-2	169	-	-
5	2.0-17.0m: Saprolite B yellowish grey colored, coarse grained powder including chips of weathered granodiorite	18-3	73	-	-
		18-4	76	-	-
		18-5	47	-	-
		18-6	36	-	-
		18-7	20	-	-
		18-8	29	-	-
		18-9	329	-	-
		18-10	38	47	-
		18-11	14	-	-
		18-12	46	-	-
		18-13	18	-	-
		18-14	12	-	-
		18-15	29	-	-
		18-16	42	-	-
17.0	17.0-40.0m: Fine grained granodiorite light grey colored, fine to medium grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1%	18-17	19	-	-
		18-18	14	-	-
		18-19	7	-	-
		18-20	7	6	-
		18-21	8	-	-
		18-22	12	-	-
		18-23	25	-	-
		18-24	12	-	-
		18-25	5	-	-
		18-26	6	-	-
		18-27	5	-	-
		18-28	5	-	-
		18-29	16	-	-
		18-30	21	34	-
		18-31	7	-	-
		18-32	16	-	-
		18-33	12	-	-
		18-34	10	-	-
		18-35	54	-	-
		18-36	78	-	-
18-37	25	-	-		
18-38	15	-	-		
18-39	23	-	-		
18-40	41	57	-		
40.0	40.0-46.0m: porphyry or porphyritic dacite dark grey colored porphyry, with plagioclase phenocrysts in a dark grey matrix	18-41	58	-	-
		18-42	1	-	-
		18-43	0	-	-
		18-44	3	-	-
		18-45	0	-	-
		18-46	49	-	-
46.0	46.0-48.0m: Fault ? dark grey colored sand	18-47	19	-	-
		18-48	104	-	-
48.0	40.0-46.0m: porphyry or porphyritic dacite dark grey colored porphyry, with plagioclase phenocrysts in a dark grey matrix, rock chip of 18-56 shows sedimentary texture with strong dissemination of pyrite and arsenopyrite	18-49	90	-	-
		18-50	17	14	-
		18-51	28	-	-
		18-52	8	-	-
		18-53	23	-	-
		18-54	16	-	-
		18-55	6	-	-
		18-56	68	-	-
		18-57	58	-	-
		18-58	37	-	-
		18-59	51	-	-
		18-60	45	44	-
60.0					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0 5 10 15 20	0.0-1.0m: Carapace reddish brown colored soil with nodules	19-1	503	-	-
	1.0-20.0m: Saprolite B yellowish grey to greenish grey colored, coarse grained sand coming from weathered granodiorite	19-2	88	-	-
		19-3	48	-	-
		19-4	15	-	-
		19-5	9	-	-
		19-6	19	-	-
		19-7	50	-	-
		19-8	18	-	-
		19-9	65	-	-
		19-10	35	24	-
		19-11	11	-	-
		19-12	17	-	-
		19-13	6	-	-
		19-14	11	-	-
		19-15	9	-	-
		19-16	35	-	-
		19-17	22	-	-
		19-18	45	-	-
		19-19	45	-	-
		20.0	17.0-40.0m: Fine grained granodiorite light grey colored, fine to medium grained granodiorite, or quartz diorite, including small amount of pyrite, amount of pyrite = 0 % to 1%	19-20	56
25 30 35 40 43.0		19-21	57	-	-
		19-22	9	-	-
		19-23	12	-	-
		19-24	17	-	-
		19-25	34	-	-
		19-26	104	-	-
		19-27	38	-	-
		19-28	42	-	-
		19-29	26	-	-
		19-30	19	22	-
		19-31	31	-	-
		19-32	27	-	-
		19-33	16	-	-
		19-34	13	-	-
		19-35	22	-	-
		19-36	18	-	-
		19-37	15	-	-
		19-38	17	-	-
		19-39	38	-	-
		19-40	23	19	-
		19-41	16	-	-
19-42	16	-	-		
19-43	40	-	-		
45 50 53.0	43.0-53.0m: meta-sandstone dark grey colored, fine grained meta -greywacke with pyrrhotite and pyrite dissemination, with small amount of epidote	19-44	37	-	-
	19-45	18	-	-	
	19-46	29	-	-	
	19-47	9	-	-	
	19-48	61	-	-	
	19-49	17	-	-	
	19-50	5	11	-	
19-51	19	-	-		
19-52	63	-	-		
53.0	53.0-60.0m: Fault ? dark grey colored sand with no rock chips	19-53	988	1,097	1,063
55 60		19-54	1,051	1,131	1,166
		19-55	37	-	-
		19-56	36	-	-
		19-57	9	-	-
		19-58	28	-	-
		19-59	7	-	-
		19-60	9	20	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Carapace brown colored soil with nodules	20-1	176	-	-
		20-2	164	-	-
5	2.0-9.0m: Saprolite A greenish yellow to greenish grey colored, fine grained powder	20-3	222	-	-
		20-4	21	-	-
		20-5	22	-	-
		20-6	24	-	-
		20-7	13	-	-
		20-8	10	-	-
		20-9	12	-	-
9.0	9.0-28.0m: Saprolite B pinkish grey to greenish grey colored saprolite including small amount of weathered rock chips	20-10	6	10	-
		20-11	8	-	-
		20-12	29	-	-
		20-13	16	-	-
		20-14	12	-	-
		20-15	98	-	-
		20-16	73	-	-
		20-17	536	-	-
		20-18	86	-	-
		20-19	58	-	-
23.0	23.0-24.0m: hard rock of granodiorite	20-20	392	428	-
		20-21	22	-	-
24.0		20-22	38	-	-
		20-23	41	-	-
25		20-24	15	-	-
		20-25	35	-	-
		20-26	56	-	-
		20-27	85	-	-
		20-28	116	-	-
		20-29	41	-	-
		20-30	28	36	-
28.0	28.0-60.0m: granodiorite light grey colored, fine to medium grained granodiorite (or quartz diorite), not altered total amount of pyrite = 0 % to 1%	20-31	63	-	-
		20-32	44	-	-
		20-33	102	-	-
		20-34	130	-	-
		20-35	146	-	-
		20-36	38	-	-
		20-37	64	-	-
		20-38	130	-	-
		20-39	106	-	-
		20-40	170	194	-
30		20-41	101	-	-
		20-42	263	-	-
		20-43	222	-	-
		20-44	62	-	-
		20-45	59	-	-
		20-46	12	-	-
		20-47	92	-	-
		20-48	54	-	-
		20-49	43	-	-
		20-50	48	46	-
35		20-51	30	-	-
		20-52	34	-	-
		20-53	44	-	-
		20-54	57	-	-
		20-55	120	-	-
		20-56	62	-	-
		20-57	39	-	-
		20-58	41	-	-
		20-59	33	-	-
		20-60	60	44	-
60.0					

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
0.0-3.0m 3.0 5 10 11.0 12.0 14.0 15 20 25 26.0 30 31.0 33.0 35 40 45 50 55 60	0.0-3.0m: Carapace brown to brownish grey colored soil with iron nodules	21-1	129	-	-
		21-2	64	-	-
		21-3	56	-	-
		21-4	22	-	-
		21-5	25	-	-
		21-6	27	-	-
		21-7	39	-	-
		21-8	28	-	-
		21-9	25	-	-
		21-10	34	39	-
		21-11	26	-	-
		21-12	52	-	-
		21-13	58	-	-
		21-14	30	-	-
		21-15	14	-	-
		21-16	29	-	-
		21-17	38	-	-
		21-18	31	-	-
		21-19	17	-	-
		21-20	16	19	-
		21-21	15	-	-
		21-22	218	-	-
		21-23	53	-	-
		21-24	51	-	-
		21-25	92	-	-
		21-26	162	-	-
		21-27	22	-	-
		21-28	25	-	-
		21-29	63	-	-
		21-30	72	54	-
	21-31	65	-	-	
	21-32	66	-	-	
	21-33	253	-	-	
	21-34	100	-	-	
	21-35	28	-	-	
	21-36	110	-	-	
	21-37	1,154	12	0	
	21-38	578	-	-	
	21-39	24	-	-	
	21-40	25	290	23	
	21-41	7	-	-	
	21-42	8	-	-	
	21-43	15	-	-	
	21-44	14	-	-	
	21-45	7	-	-	
	21-46	7	-	-	
	21-47	7	-	-	
	21-48	73	-	-	
	21-49	155	-	-	
	21-50	13	7	-	
	21-51	12	-	-	
	21-52	55	-	-	
	21-53	17	-	-	
	21-54	7	-	-	
	21-55	5	-	-	
	21-56	6	-	-	
	21-57	18	-	-	
	21-58	5	-	-	
	21-59	5	-	-	
	21-60	10	8	-	

KRC-22

Coordination: N1500 W100

Elevation: 349m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
4.0	0.0-4.0m: Carapace reddish brown colored soil with nodules	22-1	264	-	-
		22-2	1,066	274	0
		22-3	53	-	-
		22-4	393	-	-
24.0	4.0-24.0m: Saprolite A greenish grey colored, fine to medium grained saprolite, with no rock chips	22-5	157	-	-
		22-6	36	-	-
		22-7	25	-	-
		22-8	23	-	-
		22-9	84	-	-
		22-10	348	386	-
		22-11	260	-	-
		22-12	198	-	-
		22-13	41	-	-
		22-14	20	-	-
		22-15	20	-	-
		22-16	24	-	-
		22-17	15	-	-
		22-18	15	-	-
		22-19	16	-	-
		22-20	15	19	-
		22-21	12	-	-
		22-22	17	-	-
		22-23	12	-	-
		22-24	3	-	-
60.0	28.0-60.0m: granodiorite light grey colored, fine to medium grained granodiorite (or quartz diorite), not altered, sometimes it shows porphyritic texture with big plagioclase phenocrysts total amount of pyrite and pyrrhotite = 0 % to 1%	22-25	100	-	-
		22-26	117	-	-
		22-27	56	-	-
		22-28	36	-	-
		22-29	25	-	-
		22-30	21	25	-
		22-31	8	-	-
		22-32	202	-	-
		22-33	52	-	-
		22-34	5	-	-
		22-35	57	-	-
		22-36	8	-	-
		22-37	10	-	-
		22-38	10	-	-
		22-39	186	-	-
		22-40	11	8	-
		22-41	85	-	-
		22-42	98	-	-
		22-43	18	-	-
		22-44	13	-	-
		22-45	10	-	-
		22-46	7	-	-
		22-47	16	-	-
		22-48	47	-	-
		22-49	55	-	-
		22-50	150	424	-
22-51	25	-	-		
22-52	54	-	-		
22-53	196	-	-		
22-54	17	-	-		
22-55	213	-	-		
22-56	91	-	-		
22-57	47	-	-		
22-58	173	-	-		
22-59	29	-	-		
22-60	6	11	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
2.0	0.0-2.0m: Carapace brown to reddish brown colored soil with iron nodules	23-1	185	-	-
		23-2	583	-	-
5	2.0-8.0m: Saprolite A yellow to brownish yellow colored saprolite	23-3	52	-	-
		23-4	24	-	-
		23-5	60	-	-
		23-6	40	-	-
		23-7	69	-	-
		23-8	48	-	-
8.0	8.0-25.0m: Saprolite B greenish grey to yellowish grey colored saprolite including chips of weathered sandstone	23-9	54	-	-
		23-10	42	46	-
		23-11	35	-	-
		23-12	20	-	-
		23-13	20	-	-
		23-14	32	-	-
		23-15	57	-	-
		23-16	66	-	-
		23-17	56	-	-
		23-18	65	-	-
		23-19	41	-	-
		23-20	50	50	-
		23-21	8	-	-
		23-22	30	-	-
		23-23	26	-	-
25.0	25.0-37.0m: Meta-sandstone dark grey to grey colored meta-greywacke including quartz grains (1mm of diameter) with pyrite and arsenopyrite dissemination total amount of sulfides = 1 % to 3%	23-24	12	-	-
		23-25	15	-	-
		23-26	9	-	-
		23-27	28	-	-
		23-28	35	-	-
		23-29	23	-	-
		23-30	17	22	-
		23-31	24	-	-
		23-32	25	-	-
		23-33	41	-	-
37.0	37.0-48.0m: Silicified zone ? slightly silicified meta-sandstone, with pyrite and arsenopyrite dissemination quartzite ??	23-34	17	-	-
		23-35	26	-	-
		23-36	87	-	-
		23-37	219	-	-
		23-38	558	-	-
		23-39	311	-	-
		23-40	400	300	-
		23-41	41	-	-
48.0	48.0-57.0m: Meta-sandstone dark grey to grey colored meta-greywacke including quartz grains (1mm of diameter) with pyrite dissemination	23-42	57	-	-
		23-43	285	-	-
		23-44	485	-	-
		23-45	52	-	-
		23-46	178	-	-
		23-47	720	-	-
		23-48	431	-	-
57.0		23-49	69	-	-
		23-50	33	35	-
		23-51	7	-	-
		23-52	8	-	-
		23-53	4	-	-
		23-54	13	-	-
		23-55	14	-	-
		23-56	4	-	-
		23-57	31	-	-

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
1.0 5 8.0 10 15 16.0	0.0-1.0m: Carapace brown colored soil with iron nodules	24-1	32	-	-
		24-2	44	-	-
		24-3	42	-	-
	2.0-8.0m: Saprolite A greenish grey to yellow colored, fine grained powder	24-4	34	-	-
		24-5	35	-	-
		24-6	20	-	-
		24-7	13	-	-
	8.0-16.0m: Saprolite B greenish grey colored, medium grained powder including chips of meta-sandstone	24-8	24	-	-
		24-9	33	-	-
		24-10	23	9	-
		24-11	24	-	-
		24-12	33	-	-
		24-13	22	-	-
		24-14	16	-	-
		24-15	163	-	-
		24-16	72	-	-
16.0-38.0m: Meta-sandstone dark grey colored, fine to medium grained sandstone (greywacke), including quartz grains, with sulfide dissemination, sulfide amount is 1% to 3 %, sometimes higher than 3 %	24-17	351	-	-	
	24-18	267	-	-	
	24-19	79	-	-	
	24-20	44	36	-	
	24-21	195	-	-	
	24-22	148	-	-	
	24-23	24	-	-	
	24-24	22	-	-	
	24-25	66	-	-	
	24-26	26	-	-	
	24-27	25	-	-	
	24-28	12	-	-	
	24-29	10	-	-	
	24-30	8	18	-	
	24-31	238	-	-	
	24-32	6	-	-	
	24-33	2	-	-	
	24-34	0	-	-	
	24-35	4	-	-	
	24-36	7	-	-	
	24-37	12	-	-	
24-38	12	-	-		
38.0-60.0m: Alternation beds of meta-sandstone and peritic schist dark grey colored, alternation beds of meta-sandstone (greywacke) and peritic schist, with sulfide dissemination, sulfide amount is 1% to 3 %, sometimes higher than 3 % rock chip of "24-48" contains quartz veinlets	24-39	5	-	-	
	24-40	15	13	-	
	24-41	9	-	-	
	24-42	11	-	-	
	24-43	16	-	-	
	24-44	12	-	-	
	24-45	25	-	-	
	24-46	4	-	-	
	24-47	24	-	-	
	24-48	6	-	-	
	24-49	15	-	-	
	24-50	25	22	-	
24-51	21	-	-		
24-52	13	-	-		
24-53	11	-	-		
24-54	25	-	-		
24-55	30	-	-		
24-56	34	-	-		
24-57	12	-	-		
24-58	12	-	-		
24-59	8	-	-		
24-60	15	20	-		

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5 10 11.0	0.0-11.0m: Carapace reddish brown colored carapace, medium grained, with a small amount of iron nodules	74-1	11	-	-
		74-2	2	-	-
		74-3	3	-	-
		74-4	8	-	-
		74-5	9	8	-
		74-6	17	-	-
		74-7	13	-	-
		74-8	14	-	-
		74-9	22	-	-
		74-10	28	-	-
		74-11	27	-	-
15 20 25 30 31.0	11.0-31.0m: Saprolite A reddish brown colored saprolite, medium -coarse grained, homogeneous, without rock chips	74-12	104	-	-
		74-13	33	-	-
		74-14	20	-	-
		74-15	226	27	16
		74-16	13	-	-
		74-17	33	-	-
		74-18	17	-	-
		74-19	17	-	-
		74-20	34	-	-
		74-21	51	-	-
		74-22	18	-	-
		74-23	27	-	-
		74-24	30	-	-
		74-25	34	23	-
		74-26	105	-	-
		74-27	462	-	-
		74-28	54	-	-
		74-29	400	-	-
		74-30	60	-	-
74-31	43	-	-		
74-32	83	-	-		
35 40 45 50 50.0	31.0-50.0m: Saprolite A greenish light gray to yellowish gray colored saprolite, very fine grained powder, without rock chips	74-33	46	-	-
		74-34	31	-	-
		74-35	4	18	-
		74-36	24	-	-
		74-37	33	-	-
		74-38	36	-	-
		74-39	26	-	-
		74-40	9	-	-
		74-41	12	-	-
		74-42	6	-	-
		74-43	8	-	-
		74-44	8	-	-
		74-45	194	130	-
		74-46	4	-	-
		74-47	3	-	-
		74-48	1	-	-
		74-49	3	-	-
55 60 60.0	50.0-60.0m: Saprolite B yellowish light gray colored saprolite, coarse grained, including a lot of chips of weathered sandstone, (diameter: 3-4mm, sometimes black colored, meta-graywacke and peritic schist can be observed, iron-oxides along fracture)	74-50	130	-	-
		74-51	140	-	-
		74-52	19	-	-
		74-53	16	-	-
		74-54	2	-	-
		74-55	4	7	-
		74-56	9	-	-
		74-57	0	-	-
		74-58	24	-	-
		74-59	3	-	-
		74-60	2	-	-
	59.0-60.0m: Oxide zone				

KRC-75

Coordination: N1500 E400

Elevation: 382m

Depth: 60m (vertical)

Column (m)	Description	Sample No.	Au		
			ppb	ppb	ppb
5	0.0-8.0m: Carapace red to brown colored hard carapace, including a lot of lateritic gravels	75-1	32	-	-
		75-2	6	-	-
		75-3	6	-	-
		75-4	4	-	-
		75-5	6	15	-
		75-6	121	-	-
		75-7	28	-	-
		75-8	25	-	-
10	8.0-15.0m: Mottled clay brown to reddish brown colored clay, including some lateritic gravels	75-9	51	-	-
		75-10	51	-	-
		75-11	34	-	-
		75-12	24	-	-
		75-13	21	-	-
		75-14	15	-	-
		75-15	205	99	-
15	15.0-54.0m: Saprolite A brownish to yellowish brown to greenish gray colored saprolite, fine grained, soft	75-16	239	-	-
		75-17	15	-	-
		75-18	459	-	-
		75-19	21	-	-
		75-20	58	-	-
		75-21	39	-	-
		75-22	8	-	-
		75-23	11	-	-
		75-24	10	-	-
		75-25	10	5	-
		75-26	7	-	-
		75-27	4	-	-
		75-28	19	-	-
		75-29	12	-	-
		75-30	45	-	-
		75-31	9	-	-
		75-32	49	-	-
		75-33	9	-	-
		75-34	7	-	-
		75-35	6	6	-
		75-36	16	-	-
		75-37	14	-	-
		75-38	11	-	-
		75-39	3	-	-
		75-40	6	-	-
		75-41	6	-	-
		75-42	5	-	-
		75-43	4	-	-
		75-44	9	-	-
		75-45	8	7	-
		75-46	10	-	-
		75-47	6	-	-
		75-48	1	-	-
75-49	6	-	-		
75-50	0	-	-		
75-51	6	-	-		
75-52	15	-	-		
75-53	11	-	-		
75-54	12	-	-		
55	54.0-60.0m: Saprolite B green colored saprolite, including chips of deeply weathered schistosed meta-sandstone	75-55	10	6	-
		75-56	0	-	-
		75-57	0	-	-
		75-58	0	-	-
		75-59	0	-	-
		75-60	54	-	-
60					