


LEGEND

- Short wavelength High
- Short wavelength Low
- ⊞ Large amplitude SW High
- ⊞ Large amplitude SW Low

- Existing Porphyry-Cu Deposits & Prospects
- 1: Rasario
- 2: Jamiralla
- 3: Dos Hermanas
- 4: Campanane
- 5: Tignamar
- 6: Camarones
- 7: Mocha
- 8: Cucho
- 9: Queen Elizabeth
- 10: Santa Rosa
- 11: Flor del Desierto
- 12: Cerro Colorado
- 13: Tigre-San Carlos
- 14: Sagasca
- 15: La Planada
- 16: Hundida
- 17: Arauco
- 18: Copaquire
- 19: Rasario(Collahuasi)
- 20: Venus
- 21: Ponderosa
- 22: Tarapaca
- 23: Ujina(Collahuasi)
- 24: Esperanza
- 25: Quebrada Blanca
- 26: Olga, Lorena, Caniqueta

Fig. 2-6-7
SW Magnetic Anomalies and Existing Porphyry-Cu Deposits & Prospects






LEGEND


- Medium wavelength High
- Medium wavelength Low
- ▨ Large amplitude MW High
- ▨ Large amplitude MW Low

- Existing Porphyry-Cu Deposits & Prospects**
- 1: Rasario
 - 2: Jamiralla
 - 3: Dos Hermanas
 - 4: Campanane
 - 5: Tignamar
 - 6: Camarones
 - 7: Mocha
 - 8: Cucho
 - 9: Queen Elizabeth
 - 10: Santa Rosa
 - 11: Flor del Desierto
 - 12: Cerro Colorado
 - 13: Tigre-San Carlos
 - 14: Sagasca
 - 15: La Planada
 - 16: Hundida
 - 17: Arauco
 - 18: Copaquire
 - 19: Rasario(Collahuasi)
 - 20: Venus
 - 21: Ponderosa
 - 22: Tarapaca
 - 23: Ujina(Collahuasi)
 - 24: Esperanza
 - 25: Quebrada Blanca
 - 26: Olga, Lorena, Caniqueta

Fig. 2-6-8
MW Magnetic Anomalies and Existing Porphyry-Cu Deposits & Prospects



— 565 ~ 566 —



LEGEND

- Longt wavelength High
- Long wavelength Low
- Large amplitude LW High
- Large amplitude LW Low

- Existing Porphyry-Cu Deposits & Prospects
- 1: Rasario
- 2: Jamiralla
- 3: Dos Hermanas
- 4: Campanane
- 5: Tignamar
- 6: Camarones
- 7: Mocha
- 8: Cucho
- 9: Queen Elizabeth
- 10: Santa Rosa
- 11: Flor del Desierto
- 12: Cerro Colorado
- 13: Tigre-San Carlos
- 14: Sagasca
- 15: La Planada
- 16: Hundida
- 17: Arauco
- 18: Copaquire
- 19: Rasario(Collahuasi)
- 20: Venus
- 21: Ponderosa
- 22: Tarapaca
- 23: Ujina(Collauasi)
- 24: Esperanza
- 25: Quebrada Blanca
- 26: Olga,Lorena,Caniqueta

Fig. 2-6-9
LW Magnetic Anomalies and Existing Porphyry-Cu Deposits & Prospects



— 567 ~ 568 —

6-2-2 Measurement of rock magnetism

(1) Results of magnetic susceptibility measurements of rocks

Magnetic susceptibility of rocks were measured at 873 outcrops near the center of short wavelength magnetic anomalies. The measurements were made at 10 sites within an area of several meters square for each outcrop, and the mean values, outcrop number, the coordinates of the outcrop, stratigraphic division, rock species, type of alteration if any, for all measurements are listed.

The above results were statistically processed such as the frequency of mean values by stratigraphic division and rock species, and alteration types were counted. The mean magnetic susceptibility by stratigraphic horizon and rock species are shown in Table 2-6-1 and the mean susceptibility of non alteration rock by rock species in Table 2-6-2. Histograms by stratigraphic horizon are shown in Figure 2-6-10 and those by stratigraphic horizon and alteration type in Figure 2-6-11.

The following feature were clarified regarding the magnetism of the rocks of the survey area.

- ① The intensity of magnetic susceptibility decreases in the order of intrusive rocks, volcanic rocks, sedimentary rocks.
- ② The magnetic susceptibility of volcanic rocks changes significantly by the lithofacies (basaltic/andesitic), but the variation due to age (Tertiary/Cretaceous) of similar lithofacies rock is small.
- ③ Regarding sedimentary rocks, Tertiary conglomerate (sand intercalation measured) showed higher susceptibility than pre-Tertiary rocks.
- ④ Variation by lithofacies or age are small among pyroclastic rocks.

The relation between the magnetic susceptibility, rock species and alteration type for Cretaceous and Tertiary intrusive rocks is shown in Table 2-6-3. It is clear from this table that for intrusive rocks the mafic nature increases the magnetic susceptibility, and that magnetic susceptibility decreases significantly with the intensity of phyllic alteration, and also that clear correlation does not exist between the intensity of propylitic alteration and magnetic susceptibility.

Regarding Cretaceous intrusive rocks, variation of magnetic susceptibility by composition is not apparent for those which appear fresh by unaided eyes. But those strongly affected by phyllic alteration has lower susceptibility, while strongly propylitized rocks do not have lower values.

(2) Rock susceptibility and magnetic anomaly

The magnetic susceptibility measured at 873 outcrops was divided into 7 ranks and they were given different colors, and were overlaid with short and medium wavelength magnetic anomalies on maps. They are shown in Figures 2-6-12 and 2-6-13.

Generally rock bodies with higher magnetic susceptibility show high anomalies, and

low magnetic anomalies are considered to occur in rock bodies with high magnetic susceptibility and reverse remanent magnetism. This relation can be observed in many magnetic anomalies of Figures 2-6-12 and 2-6-13.

In these two figures, however, there are significant number of magnetic anomalies corresponding to low magnetic susceptibility. These anomalies are surrounded by circles (broken lines) in these figures. It is difficult theoretically to conceive rock bodies with low susceptibility to show magnetic anomalies, and it would be natural to consider these anomalies to be caused by strong remanent magnetism. As will be mentioned later in this report, strong remanent magnetism was measured for tuffs with low magnetic susceptibility, and these rocks provably cause magnetic anomalies similar to those by intrusive rocks with high susceptibility.

(3) Magnetic susceptibility of drill holes

The results of magnetic susceptibility measurement of drill cuttings at every 20m interval are shown in Figure 2-6-14.

These measurements clarified the following features regarding the distribution of magnetic susceptibility to approximately 500m subsurface depth.

- ① There are parts of Quaternary gravel beds (Qcp(Grav)) and Tertiary conglomerate (Tc(cgl)) where high magnetic susceptibility close to 10^{-2} appear locally (MJC-2, 7, 8, 9).
- ② The magnetic susceptibility of Tertiary ignimbrite (Tig) is $0.5\sim 5.0 \times 10^{-3}$ with the exception of altered parts. These values agree well with the values measured on the surface.
- ③ The magnetic susceptibility of the intrusive rocks and pre-Tertiary rocks (MJC-1, 11, 12) are low at less than 10^{-3} for parts affected by phyllic alteration, but propylitized parts show values mostly around 10^{-2} . These values agree well with those in Table 2-6-3.

Table 2-6-1 Average Magnetic Susceptibility of Formation by Alteration Type (SI units)

Formation	Nos.	Non alteration	Alteration Type											
			Nos.	Oxidation	Nos.	Acid alt.	Nos.	Silicification	Nos.	Propylite alt.	Nos.	Phyllic alt.	Nos.	Potassic alt.
Qv	91	0.01225	3	0.00396	10	0.00055	2	0.00022	5	0.02565	14	0.00592		-
Qcp,Qvc	17	0.00443		-	1	0.00012		-		-		-		-
Qvr	63	0.00327		-	4	0.00054	7	0.00027	5	0.00060	3	0.00086		-
Tig	130	0.00230		-	1	0.00029	1	0.00012		-	8	0.00129		-
Tc	2	0.00328		-		-		-		-		-		-
Tgd	24	0.01742	1	0.02659	3	0.00193	5	0.00262	40	0.01772	44	0.00520	2	0.03523
kgd	20	0.01723		-	1	0.00028	3	0.00173	21	0.01636	15	0.00454		-
Kv	88	0.00677	3	0.00094	5	0.00021	28	0.01039	77	0.01004	28	0.00266		-
Kc	6	0.00075	2	0.00142		-	13	0.00041	6	0.00146		-		-
Jc,Jm	23	0.00134		-		-		-	6	0.00536	2	0.00006		-
Jv	5	0.00783	1	0.00007		-	2	0.00011	2	0.02150		-		-
P	8	0.01003		-		-		-	2	0.00016	1	0.00025		-

Table 2-6-2 Average Magnetic Susceptibility of Non Alteration Rocks (SI units)

Rock Name	Geologic Age									
	No.	Quaternary	No.	Tertiary	No.	Cretaceous	No.	Jurassic	No.	Paleozoic
Andesite	20	0.01111		-	24	0.00765	3	0.01270		-
Basalt	50	0.01509		-	7	0.01876		-		-
Dacite	4	0.01229		-	4	0.01181		-		-
Ignimbrite	25	0.00503	65	0.00221		-		-		-
Lap. tuff	2	0.00507	5	0.00119		-		-		-
Pum. tuff	23	0.00172	28	0.00122		-		-		-
Tuff	4	0.00078	11	0.00277	12	0.00177		-		-
Welded tuff	3	0.00826	11	0.00260		-		-		-
Sandstone	9	0.00706		-	8	0.00122	9	0.00033		-
Conglomerate	2	0.00211		-		-		-		-
Shale		-		-	2	0.00043	6	0.00091		-
Granite		-	3	0.01089	3	0.01089		-		-
granodiorite		-	11	0.01980	12	0.01730		-		-
Diorite		-	3	0.01908	3	0.04790		-		-
Schist		-		-		-		-	3	0.00026
Gneiss		-		-		-		-	2	0.00056




Fig. 2-6-10 Histogram of Susceptibility by Geologic Formation







Fig. 2-6-11 Histogram of Susceptibility by Alteration Type

Table 2-6-3 Variation of Magnetic Susceptibility in Tertiary Intrusive Rocks (Tgd) and Cretaceous Intrusive Rocks (Kgd)

Tgd-G,Qp Alteration Type	acidic / phyllic			propylitic			silicification		fresh
	h	m/m-s	s	h	m	s	m	s	
Maximum	0.33	13.24	2.35	-	-	5.66	-	0.56	28.4
Minimum	0.04	0.03	0.04	-	-	0.60	-	0.12	0.9
Average	0.15	3.61	0.56	-	-	2.52	-	0.34	10.0
Median	0.12	0.81	0.10	-	2.56	1.83	0.03	0.34	0.9
Number	11	7	5	-	1	5	1	2	3

Kgd-G,Qp Alteration Type	acidic / phyllic			propylitic			silicification		fresh
	h	m/m-s	s	h	m	s	m	s	
Maximum	8.24	-	-	-	-	7.66	-	-	40.0
Minimum	-	-	-	-	-	6.48	-	-	0.5
Average	-	-	-	-	-	7.25	-	-	18.2
Median	-	-	-	-	-	7.61	-	-	16.1
Number	1	-	-	-	-	3	-	-	4

Tgd-Gd,Da Alteration Type	acidic / phyllic			propylitic			silicification		fresh
	h	m/m-s	s	h	m/m-s	s	m	s	
Maximum	-	27.96	14.07	46.4	26.5	31.37	-	-	48.0
Minimum	-	0.12	12.59	20.0	0.8	1.84	-	-	0.1
Average	-	5.63	13.33	33.2	12.0	11.83	-	-	16.0
Median	-	0.58	13.33	33.2	11.6	7.57	-	7.75	16.6
Number	-	9	2	2	6	8	-	1	10

Kgd-Gd,Da Alteration Type	acidic / phyllic			propylitic			silicification		fresh
	h/h-m	m/m-s	s	h	m/m-s	s	m	s	
Maximum	-	16.68	-	27.6	15.1	26.81	-	-	26.8
Minimum	-	0.28	-	10.7	5.6	7.00	-	-	5.5
Average	-	3.64	-	21.5	9.8	15.09	-	-	17.7
Median	0.32	2.00	22.00	26.1	9.2	11.46	-	-	18.8
Number	1	10	1	3	4	3	-	-	11

Tgd-Di,Qdi Alteration Type	acidic / phyllic			propylitic			silicification		fresh
	h	m/m-s	s	h/h-m	m/m-s	s	m	s	
Maximum	28.09	10.69	46.66	41.4	50.1	57.64	-	-	50.4
Minimum	0.61	0.16	38.08	0.9	10.2	6.37	-	-	15.4
Average	14.35	3.57	42.37	23.1	32.1	25.26	-	-	24.9
Median	14.35	1.72	42.37	18.3	33.5	19.78	-	-	20.9
Number	2	4	2	5	6	10	-	-	6

Kgd-Di,Qdi Alteration Type	acidic / phyllic			propylitic			silicification		fresh
	h	m/m-s	s	h/h-m	m/m-s	s	m	s	
Maximum	-	-	-	-	-	-	-	-	-
Minimum	-	-	-	-	-	-	-	-	-
Average	-	-	-	-	-	-	-	-	-
Median	-	-	-	12.0	-	4.38	-	-	11.9
Number	-	-	-	1	-	1	-	-	1

Abbr. : h=high, m=moderate, s=small, Susceptibility= 10^{-3} SI

G=granite, Qp=quartz porphyry, Gd=granodiorite, Da=dacite, Di=diorite, Qd=quartz diorite




Fig. 2-6-12 Distribution of Classified Susceptibility (Northern Part)




Fig. 2-6-13 Distribution of Classified Susceptibility (Southern Part)




Fig.2-6-14
Susceptibility Profiles
of Drill hole

(4) Remanent magnetization measurements

The 14 sampling sites for remanent magnetism measurements are shown in Figure 3-6-15. Ten samples were measured for each site and the mean values are shown in Table 2-6-4. In this table, the initial remanent magnetic values (INIT), values after 30mT alternating field demagnetization, polarity of remanent magnetism, Q values, and magnetic susceptibility are shown. Magnetic susceptibility was measured on cylindrical (25mm diameter, 22mm high) test piece used for remanent magnetic measurement. Magnetic susceptibility meter Bartington MS-2 was used. Also the test piece is in a weak magnetic field during the measurement and thus it was done after remanent magnetization measurement so as not to affect the latter operation.

Q value is also called Königsberger ratio and is defined as follows (Nagata, 1961).

$$Q = M/kH$$

Q: Q value

M: remanent magnetization intensity (usually initial value)

k : magnetic susceptibility

H: geomagnetic field

The Q value indicates the ratio of induced magnetization (kH) to remanent magnetization (M), and where this value is greater than 1.0, the contribution of remanent magnetization is greater than induced magnetization to airborne magnetic maps.

It is known that magnetic susceptibility k of rocks is almost proportionate to the volume of magnetite contained. On the other hand, from the mechanism of acquiring remanent magnetism, the intensity increases with smaller grain size of magnetite and with larger amount of the mineral. In igneous rocks, it is also shown from the mechanism that the intensity increases with the cooling rate of magma (Nagata, 1961).

According to Table 2-6-4, the polarity of remanent magnetism is normal at 8 localities out of 14 and is reverse at 6 localities. Thus the occurrence of normal and reverse magnetization is about the same.

The magnetic declination of normal polarity is within $6^\circ \pm 20^\circ$ and those of reverse polarity are in two directions of $183^\circ \pm 15^\circ$ (4 localities) and $239^\circ \pm 17^\circ$ (2 localities).

The Q values of 3 tuff samples with low magnetic susceptibility are all high at 28, 15, and 10. This indicates that Tertiary and Quaternary ignimbrites which show low magnetic susceptibility in the order of 10^{-3} at outcrops and cuttings, if they have remanent magnetization, can show strong magnetic anomaly similar to those by intrusive and volcanic rocks with high susceptibility.

The Q values of pre-Tertiary to Tertiary intrusive rocks are 0.22 to 18. Of the 10 samples measured, 4 had Q values of less than 1.0 indicating a number of cases with lower remanent magnetization than induced magnetization. These 4 samples,

however, have very high magnetic susceptibility of $3\sim 6 \times 10^{-2}$, and this is considered to have caused the low Q below 1.0.

(5) Remanent magnetization and magnetic anomaly

The results of remanent magnetism measurements are plotted on short and medium wavelength magnetic anomalies (Fig. 2-6-16). The normal/reverse polarity of remanent magnetization coincides well with the high/low anomaly of the short and medium wavelength anomalies near the sampling sites. This fact indicates that the cause of the low magnetic anomalies is the reverse polarity of the remanent magnetism.

Regarding a medium wavelength low magnetic anomaly in the central part, remanent magnetism has been measured for 3 samples (LS-105, LS-120, LK-187), and 2 of them have normal polarity and the other reverse. Now, comparing the intensity of the remanent magnetism by "Q value \times magnetic susceptibility", the reverse polarity sample LK-187 is 0.3, while the normal LS-105 and LS-120 are 0.02 and 0.08 respectively. The reverse remanent magnetism is four times stronger than the normal value. These facts do not contradict the occurrence of low magnetism. This example shows clearly that one magnetic anomaly does not necessarily correspond to one magnetic rock body and can be formed by the interaction of magnetism of more than one rock body.

Table 2-6-4 Results of Remanent Magnetization Measurement

Outcrop No.	Sample No.	Rock facies			Lab. No.	INIT					After AFDEMAG (30 mT)					Po	Q	Susceptibility ($\times 10^{-3}$ SI unit)
		Formation/Intrusive	Rock Name	K-Ar age (Ma)		D°	I°	k	α	R	D°	I°	k	α	R			
LQ-268	Q-149	Qvr	Dacitic Tuff	-	8492	116	-57	37	8.1	9.756	353	-28	542	2.1	9.983	N	28	3.92
LQ-269	Q-150	Qvr	Pumice Tuff	-	9501	186	31	2271	1.0	9.996	190	25	1530	1.2	9.994	R	15	3.94
LS-133	S-047	Tig	Welded Tuff	-	0471	354	-50	260	3.0	9.965	355	-49	264	3.0	9.966	N	10	2.40
LK-056	K-016	Tgd	Granodiorite	44.6±1.1	5162	007	-55	161	3.8	9.944	009	-54	145	4.0	9.938	N	2.2	11.00
LS-052	S-010	Tgd	Granite	-	6010	247	-19	7	19.0	8.784	198	57	631	1.9	9.986	R	0.22	52.90
LK-187	K-084	Tgd	meta-Diorite porphyry	58.1±1.9	7842	183	27	129	4.3	9.930	173	32	137	4.1	9.935	R	15	21.00
LS-105	S-034	Tgd	Diorite porphyry	-	0341	040	-56	67	6.0	9.865	014	-24	229	3.2	9.961	N	0.39	51.90
LS-120	S-045	Tgd	Diorite	58.8±2	0452	025	-71	47	7.1	9.807	002	-48	300	2.8	9.970	N	5.3	15.00
LS-132	S-046	Tgd	Quartzdiorite	-	0462	003	-63	788	1.7	9.989	007	-61	1415	1.3	9.994	N	2.0	4.80
LT-212	T-080	Tgd	Porphyry	-	0801	231	49	4	29.7	7.501	170	45	156	3.9	9.942	R	0.27	38.40
LQ-048	Q-011	Kgd	Microdiorite	-	1111	065	-24	24	10.0	9.631	026	-4	6	21.9	8.458	N	16	7.90
LQ-056	Q-013	Kgd	Granite porphyry	-	2131	353	-69	64	6.1	9.860	004	-61	557	2.0	9.984	N	0.64	60.20
LT-077	T-020	Kv(i)	Sandstone	-	3202	225	-24	2	48.1	5.430	222	44	14	13.2	9.370	R	0.43	0.95
LK-023	K-011	Pzg	Diorite	300±7	4111	268	26	50	6.9	9.819	256	50	5	25.3	8.040	R	18	26.10

D°: Declination

I°: Inclination

k: Fisher's best estimation of precision (Fisher, 1953)

α : Half angle of the cone of confidence at p=0.95

R: Resultant of vector sum

Po: Polarity (N: Normal, R: Reverse)

Q: Q-value (Königsberger ratio)






Fig.2-6-15
Remanence Sample Location



— 589 —
— 590 —