

[References]

1. Akimoto, K. and Hasegawa, S. (1989): Depth distribution of recent benthic foraminifera of waters close to Japan - Implication for the paleoenvironmental scale of water depth. *Mem. Geol. Soc. Japan*, 32, 229-240 (in Japanese).
2. Anderson, O.R., Bennett, P. and Bryan, M. (1989) Experimental and observational studies of radiolarian physiological ecology: 3. Effects of temperature, salinity and light intensity on the growth and survival of *Spongaster tetras* maintained in laboratory culture. *Marine Micropaleontology*, 14, 275-282.
3. Barrett, T. (1982) Stratigraphy and sedimentology of Jurassic bedded chert overlying ophiolites in the North Apennines, Italy. *Sedimentology*, 29, 353-373.
4. Berggren, W. A., Kent, D. V., Swisher, III, C. C., and Aubry, M.-P. (1995) A revised Cenozoic geochronology and chronostratigraphy. SEPM Special Publication, No. 54, 129-212.
5. Craig, H., Chung, Y., and Fiadeiro, M. (1972) A benthic front in the South Pacific, *Earth and Planetary Sci. Letters*, 16, 50-55.
6. Cronan, D. S. (1984) South Pacific. *Mar. Geol. Notes*, 3, 1-17.
7. Cronan, D. S., Johnson, C. J. and Clark, A. L (1986) Manganese Nodules, Pacific mineral resources physical, economic, and legal issues, 161-179.
8. De Carlo, E. H. and McMurtry G. M. (1992) Rare earth element geochemistry of ferromanganese crusts from the Hawaiian Archipelago, central Pacific, *Chemical Geology*, 95, 235-250.
9. Doyle, P. S. and Riedel, W. R. (1979) Ichthyoliths: present status of taxonomy and stratigraphy of microscopic fish skeletal debris. *Scripps Institution of Oceanography Reference Series*, no. 79-16, pp231.
10. Doyle, P. S. and Riedel, W. R. (1985) Cenozoic and Late Cretaceous ichthyoliths. In Bolli, H. M. ed. *Plankton Stratigraphy*, 965-995.
11. Dumitrica, P. (1983) Systematics and evolution of the genus *Suttonium* Schauf (Radiolaria). *Revue de Micropaleontologie*, 26, 36-47.
12. Exon, N. F. (1983) Manganese nodule deposits in the Central Pacific Ocean and their variation with latitude. *Mar. Mining*, 4, 79-107.
13. Gowing, M. M. (1993) Seasonal radiolarian flux at the VERTEX North Pacific time-series site. *Deep-Sea Research*, 40, 517-545.
14. Griffin, J. J., Windom, H. and Goldberg, E. D. (1968) The distribution of clay minerals in the World Ocean, *Deep sea Res.*, 15, 433-459.

15. Halback, P., Friedrich G., and Stackelberg (Ed.)(1988) The manganese nodule belt of the Pacific Ocean, Ferdinand Enke, Stuttgart, pp 254
16. Haskin, L. A., Haskin M. A. Ferey F. A. and Wildman, T. R. (1968) Relative and absolute terrestrial abundance of the rare earth. In Ahrens L. H. (ed) Origin and distribution of the elements, Vol. 1 Pergamon, Oxford, 889-911.
17. Hein, J. R., Schulz, M. S. and Gein, L. M. (1992) Central Pacific cobalt-rich ferromanganese crusts: Historical perspective and regional variability, *Geology and offshore mineral resources of the central Pacific basin*, 261-283.
18. Inoue, Y. (1989) Northwest Pacific foraminifera as paleoenvironmental indicators. *Sci. Rep. Inst. Geosci., Univ. Tsukuba, Ser B*, vol. 10, 57-162.
19. Iwashita, M., Komaki, Y., Hoshino, T., Horibe, S., Masuzawa, J. (ed) (1972): *Chemistry of Sediments*. Tokai Univ. Press, 571pp (in Japanese)
20. JICA/MMAJ (1986): Ocean Resources Investigation in the Sea Area of CCOP/SOPAC Report on the Joint Basic Study for the Development of Resources (volume 1), Sea Area of Cook Islands
21. JICA/MMAJ (1987): Ocean Resources Investigation in the Sea Area of CCOP/SOPAC Report on the Joint Basic Study for the Development of Resources (volume 2), Sea Area of Cook Islands
22. JICA/MMAJ (1991): Ocean Resources Investigation in the Sea Area of SOPAC Report on the Joint Basic Study for the Development of Resources (volume 1-2), Sea Area of the Cook Islands.
23. JICA/MMAJ (2001): Ocean Resources Investigation in the Sea Area of SOPAC Report on the Joint Basic Study for the Development of Resources (volume 1), Sea Area of the Cook Islands.
24. Lombari, G. and Boden, G. (1985): Modern radiolarian global distributions. Cushman foundation for Foraminiferal Research, Special Publication, 16A, 125p.
25. Nigrini, C. (1967): Radiolaria in pelagic sediments from the Indian and Atlantic Oceans. *Bulletin of the Scripps Institution of Oceanography, University of California*, 11, 125p.
26. Pautot, G. and Melguen, M. (1979): Influence of deep water circulation and sea floor morphology on the abundance and grade of Central South Pacific manganese nodules, In Bischoff, J.L. and Piper, D.Z. (ed) *Marine Geology and Oceanography of the Pacific Manganese Nodule Province*, Plenum Press, New York and London, 621-649

27. Sanfilippo, A. and Nigrini, C. (1998): Code numbers for Cenozoic low latitude radiolarian biostratigraphic zones and GPTS conversion tables. *Marine Micropaleontology*, 33, 109-156.
28. Usui, A. and Moritani, T. (1992): Manganese nodule deposits in the Central Pacific Basin: Distribution, geochemistry, mineralogy and genesis. In Keating B. H. and Bolton B. R. (Ed.), *Geology and Offshore Mineral Resources of the Central Pacific Basin*, Springer, New York, 14, 205-223.
29. Usui A. and Someya M. (1997) Distribution and composition of marine hydrogenetic and hydrothermal deposits in the northwest Pacific, In Nicholson, K., Hein, J. R., Buhn, B. and Dasgupta (ed) *Manganese Mineralization: Geochemistry and Mineralogy of Terrestrial and Marine Deposits*, Geol. Soc. Special Pub. No.119, 177-198.
30. Wakita H., Rey, P., Schmitt R. A. (1971) Abundance of the 14 rare-earth elements and 12 other trace elements in Apollo 12 samples: five igneous and one breccia rocks and four soils. *Proc. 2nd Lunar. Sci. Conf.* Pergamon, Oxford, 1319-1329.